



## Corrigendum to “Simultaneous synthesis of lactic acid and hydrogen from sugars via capnophilic lactic fermentation by *Thermotoga neapolitana cf capnolactica*” [Biomass and Bioenergy 125 (2019) 17–22]



Nirakar Pradhan <sup>a,b,\*,1</sup>, Giuliana d’Ippolito <sup>a,\*\*</sup>, Laura Dipasquale <sup>a</sup>, Giovanni Esposito <sup>b</sup>, Antonio Panico <sup>c</sup>, Piet N.L. Lens <sup>d</sup>, Angelo Fontana <sup>a</sup>

<sup>a</sup> National Research Council of Italy, Institute of Biomolecular Chemistry, Via Campi Flegrei 34, 80078, Pozzuoli, Napoli, Italy

<sup>b</sup> Department of Civil and Mechanical Engineering, University of Cassino and Southern Lazio, Via Di Biasio, 43, 03043, Cassino, FR, Italy

<sup>c</sup> Telematic University Pegaso, Piazza Trieste e Trento, 48, 80132, Naples, Italy

<sup>d</sup> UNESCO-IHE Delft, Westvest 7, 2611AX, Delft, the Netherlands

The authors regret Fig. 3, the actual figure was inadvertently repeated as that of Fig. 2. The correct figure has been printed below.

The authors would like to apologise for any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.biombioe.2019.04.007>.

\* Corresponding author. Department of Civil and Mechanical Engineering, University of Cassino and Southern Lazio, Via Di Biasio, 43, 03043, Cassino, FR, Italy. , \*\* Corresponding author.

E-mail addresses: [nirakar.pradhan@gmail.com](mailto:nirakar.pradhan@gmail.com), [npradhan@ntu.edu.sg](mailto:npradhan@ntu.edu.sg) (N. Pradhan), [gdippolito@icb.cnr.it](mailto:gdippolito@icb.cnr.it) (Giuliana d’Ippolito), [ldipasquale@icb.cnr.it](mailto:ldipasquale@icb.cnr.it) (L. Dipasquale), [giovanni.esposito@unicas.it](mailto:giovanni.esposito@unicas.it) (G. Esposito), [antonio.panico@unipegaso.it](mailto:antonio.panico@unipegaso.it) (A. Panico), [p.lens@un-ihe.org](mailto:p.lens@un-ihe.org) (P.N.L. Lens), [afontana@icb.cnr.it](mailto:afontana@icb.cnr.it) (A. Fontana).

<sup>1</sup> Current address: Singapore Centre for Environmental Life Sciences Engineering, Nanyang Technological University, 60 Nanyang Dr, 637551, Singapore.

<https://doi.org/10.1016/j.biombioe.2021.105999>

Available online 15 February 2021

0961-9534/© 2021 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 3. Effect of carbon source ( $5 \text{ g L}^{-1}$ ) on the yield ( $\text{mol mol}^{-1}$  of sugar) of (A)  $\text{H}_2$ , (B) acetic acid and (C) lactic acid and (D) cell dry weight and biomass yield ( $\text{g g}^{-1}$  of sugar) of *Tn<sub>ef</sub>* via the CLF pathway after 72 h of incubation. Error bar = standard deviation ( $n = 3$ ). CMC =

Carboxymethyl cellulose. *p*-value was calculated and compared with experiments performed in standard culture medium with glucose as carbon source. ns = not significant; *p* value  $< 0.05$  (\*); *p*  $< 0.01$  (\*\*); *p* value  $< 0.001$  (\*\*\*)�.