

Manuscript Number: LWT-D-19-00894R1

Title: Impact of co-inoculation of *Saccharomyces cerevisiae*, *Hanseniaspora uvarum* and *Oenococcus oeni* autochthonous strains in controlled multi starter grape must fermentations

Article Type: Research paper

Keywords: wine; mixed-starter; *Hanseniaspora uvarum*; *Oenococcus oeni*; autochthonous yeast

Corresponding Author: Dr. Francesco Grieco,

Corresponding Author's Institution: CNR- Istituto di Scienze delle Produzioni Alimentari- Lecce

First Author: Vittorio Capozzi

Order of Authors: Vittorio Capozzi; Carmen Berbegal; Maria Tufariello; Francesco Grieco; Giuseppe Spano; Francesco Grieco

Abstract: The use of multi-species starter cultures is an approach of increasing significance for winemakers in order to improve the general quality and safety of the final product. As first step of the present study, we isolated and characterize two *Saccharomyces cerevisiae* yeast starter strains, denoted as ITEM 167292 and ITEM 17293, from natural must fermentations of "Negroamaro" grapes. As second step, we studied the interactions during grape must fermentation between these two strains, the *Hanseniaspora uvarum* strain ITEM 8785 and five autochthonous *Oenococcus oeni* strains, by microbial counts and by quantifying L-malic acid and ethanol concentrations. The best performing *O. oeni* strain, namely OT4, was used to create, with the *H. uvarum* strain ITEM 8785, two mixed starter formulations with the strains ITEM 167292 and ITEM 17293. The three microbial species showed to be compatible and to complete the fermentative processes producing wines denoted by reduced acetic acid concentrations. The performance of the mixed starter formulations were then validated by carrying pilot-scale vinifications. At the best of our knowledge, this report is the first study regarding the utilization of selected *H. uvarum*/*S. cerevisiae*/*O. oeni* autochthonous strains in a simultaneous multi-starter inoculation for the industrial production of regional typical wines.

- The first investigation on a non-*Saccharomyces/Saccharomyces/Oeni* starter culture is proposed.
- The compatibility among microbial species during fermentation was assessed.
- The mixed starter produced red wine with reduced acetic acid content.
- The results presented were validated by pilot-scale vinification trials
- The industrial application of the mixed starter formulation reported is a promising approach.

1     **Impact of co-inoculation of *Saccharomyces cerevisiae*, *Hanseniaspora***  
2     ***uvarum* and *Oenococcus oeni* autochthonous strains in controlled multi**  
3             **starter grape must fermentations**

4  
5     **Vittorio Capozzi <sup>a,§</sup>, Carmen Berbegal <sup>a,§</sup>, Maria Tufariello <sup>b</sup>, Francesco Grieco <sup>c</sup>,**  
6             **Giuseppe Spano <sup>b</sup>, Francesco Grieco <sup>b</sup>**

7  
8     <sup>a</sup> Department of the Sciences of Agriculture, Food and Environment, University of  
9     Foggia, Foggia, Italy

10    <sup>b</sup> CNR – Institute of Sciences of Food Production (ISPA), via Prov.le, Lecce-Monteroni,  
11    - 73100 Lecce, Italy

12    <sup>c</sup> CNR - Istituto di Scienze delle Produzioni Alimentari (ISPA), via G. Amendola,  
13    122/O, 70126 Bari, Italy

14  
15    §: Both authors equally contributed to this work

16  
17  
18    \*Corresponding author

19    Francesco Grieco, National Research Council - Institute of Sciences of Food Production  
20    (ISPA), via Prov.le Lecce-Monteroni, 165 - 73100 Lecce, Italy. Phone:  
21    +390832422612; Fax: +390832422620; Email: francesco.grieco@ispa.cnr.it

22

23 **Abstract**

24 The use of multi-species starter cultures is an approach of increasing significance for  
25 winemakers in order to improve the general quality and safety of the final product. As  
26 first step of the present study, we isolated and characterize two *Saccharomyces*  
27 *cerevisiae* yeast starter strains, denoted as ITEM 167292 and ITEM 17293, from natural  
28 must fermentations of “Negroamaro” grapes. As second step, we studied the  
29 interactions during grape must fermentation between these two strains, the  
30 *Hanseniaspora uvarum* strain ITEM 8785 and five autochthonous *Oenococcus oeni*  
31 strains, by microbial counts and by quantifying L-malic acid and ethanol concentrations.  
32 The best performing *O. oeni* strain, namely OT4, was used to create, with the *H.*  
33 *uvarum* strain ITEM 8785, two mixed starter formulations with the strains ITEM  
34 167292 and ITEM 17293. The three microbial species showed to be compatible and to  
35 complete the fermentative processes producing wines denoted by reduced acetic acid  
36 concentrations. The performance of the mixed starter formulations were then validated  
37 by carrying pilot-scale vinifications. At the best of our knowledge, this report is the first  
38 study regarding the utilization of selected *H. uvarum/S. cerevisiae/O. oeni*  
39 autochthonous strains in a simultaneous multi-starter inoculation for the industrial  
40 production of regional typical wines.

41

42 **Keywords:** wine; mixed-starter; *Hanseniaspora uvarum*; *Oenococcus oeni*;  
43 autochthonous yeast

44

## 45 **1. Introduction**

46 Traditionally, the vinification process of red wines includes two essential stages, i.e. the  
47 alcoholic fermentation (AF) and the malolactic fermentation (MLF). During the AF, the  
48 sugars of the must are transformed into ethanol and this process is carried out by the  
49 yeasts, principally by *Saccharomyces cerevisiae* (Garofalo, Tristezza, Grieco, Spano &  
50 Capozzi, 2016). However, non-*Saccharomyces* species have a role in the AF and they  
51 contribute to enhance the organoleptic properties of wine (Liu, Lu, Duan & Yan, 2016;  
52 Petruzzi et al., 2017). Several non-*Saccharomyces* species have been studied in mixed  
53 fermentations with the scope of adding peculiar features to the wine (Ciani, Beco &  
54 Comitini, 2006; Ciani, Comitini, Mannazzu & Domizio, 2009; Comitini et al., 2011;  
55 Suzzi et al., 2012, Tristezza et al., 2016b). These mixed cultures have an additional  
56 interest when they are formed by autochthonous selected yeasts, since they are adapted  
57 to the conditions of a specific wine-production area (Capozzi, Garofalo, Chiriatti,  
58 Grieco & Spano, 2015; Lopes, Rodriguez, Sangorrin, Querol & Caballero, 2007; Tofalo  
59 et al., 2016) and may ensure the maintenance of the typical oenological and sensory  
60 characteristics of wine (Rodríguez et al., 2010).

61 The development of efficient malolactic starter cultures is crucial for the oenological  
62 industry (Berbegal et al., 2016, Brizuela et al., 2017). Several are the strain-specific  
63 features requested for a malolactic starter culture, such as the capacity to stand low pH,  
64 high ethanol and SO<sub>2</sub> concentrations, the absence of biogenic amines production, the  
65 compatibility with yeast selected strains (Berbegal et al., 2016; Capozzi et al., 2010).

66 Besides, a critical step in the employment of MLF starters is the time of inoculation.  
67 **Lactic acid bacteria (LAB)** starters can be co-inoculated with yeast at the beginning of  
68 AF, or sequentially inoculated after the AF (Bartowsky, Costello & Chambers, 2015).  
69 Several recent investigations have indicated that when bacteria are directly inoculated

70 into the must they performed better than they when added after the end to the AF  
71 (Abrahamse & Bartowsky, 2011; Tristezza et al., 2016a).

72 In a previous study, the *H. uvarum* ITEM 8795 was selected because of its contribution  
73 in increasing the wine organoleptic quality and reducing the volatile acidity (De  
74 Benedictis, Bleve, Grieco, Tristezza & Tufariello, 2011). The oenological potential of  
75 this strain in co-inoculation and in a sequential inoculation with *S. cerevisiae* was also  
76 assessed by industrial wine production (Tristezza et al., 2016b).

77 In the present investigation, we report the selection of Apulian autochthonous *S.*  
78 *cerevisiae* and *O. oeni* strains to design of a mixed starter culture with *H. uvarum* ITEM  
79 8795 to simultaneously perform the AF and MLF. Furthermore, we evaluated the  
80 compatibility between the different microorganisms employed in the autochthonous  
81 mixed starter culture and the best inoculation time of *O. oeni* strains. At the best of our  
82 knowledge, this study described, for the first time the fermentative performance of a  
83 non-*Saccharomyces/Saccharomyces/O. oeni* mixed starter formulation.

84

85

## 86 **2. Material and methods**

87

### 88 *2.1 Microorganisms*

89 Yeast strains used in the present study are deposited in Agro- Food Microbial Culture  
90 Collection of ISPA (<http://www.ispacnr.it/collezioni-microbiche/>). All yeast strains  
91 were cultured in YPD (Sigma-Aldrich, USA) and incubated at 28°C 24-48 hours. *O.*  
92 *oeni* strains were previously isolated from Nero di Troia wine (Capozzi et al., 2014) and  
93 they are deposited in the collection of the Industrial Microbiology Laboratory

94 (University of Foggia). LAB strains were cultured in MRS broth (Sigma-Aldrich, USA)  
95 and incubated at 28 °C for 4-7 days.

96

### 97 *2.2 Yeast isolation and S. cerevisiae strains identification*

98 The enological selection was carried out according to Tufariello et al. (2019) from  
99 spontaneous fermentations of Negroamaro grapes collected in the “Brindisi” PDO/DOC  
100 area. Briefly, yeast isolates were firstly screened for their ability to produce hydrogen  
101 sulphide on Biggy agar (Sigma, USA). H<sub>2</sub>S-low producer isolates (i.e. white or light  
102 brown colonies) were selected for genetic characterization. The isolates were identified  
103 at species-level by PCR analysis of the ribosomal RNA region (Tufariello et al., 2019)  
104 and at strain-level by interdelta typing (Tristezza, Gerardi, Logrieco & Grieco, 2009).  
105 The amplified DNA products were visualized and analyzed by agarose gel  
106 electrophoresis (Hay et al., 1994).

107

### 108 *2.3 Lab-scale vinification*

109 The identified *S. cerevisiae* strains were tested by micro-fermentation assays conducted  
110 in Negroamaro grape must (21.5° Babo; 7.2 g/L total acidity; pH 3.4) added with 100  
111 mg/L potassium metabisulfite. One liter of treated must was inoculated with 10<sup>6</sup>  
112 CFU/mL of yeast culture. The vinifications were carried out in triplicate at 25°C and  
113 daily monitored by measuring the reducing sugars concentration. Wines were then  
114 filtered, separately bottled and stored at 18 °C for the sensorial analysis (Tufariello et  
115 al., 2019).

116

### 117 *2.4 Co-inoculation tests*

118 For the co-inoculation trials, yeast and bacteria starter cultures were prepared by  
119 growing strains in YPD or MRS medium as described above and then inoculating in  
120 triplicate the strains into 200 mL of Negroamaro grape must from (21° Babo; 7.2 g/L  
121 total acidity; 2.57 g/L malic acid; pH 3.78). Using the 2 selected *S. cerevisiae*, 1 *H.*  
122 *uvarum* and 5 *O. oeni* strains, a total of 10 different starter culture combinations were  
123 carried out. In the mixed starter cultures, the *H. uvarum* strain was simultaneously  
124 inoculated with *S. cerevisiae* in a 1:100 inoculum ratio (respectively 10<sup>4</sup> CFU/mL and  
125 10<sup>6</sup>CFU/mL). *O. oeni* strains were co-inoculated (ethanol content 0%) or sequentially  
126 inoculated during AF, when ethanol content was 2%, 4%, 6%, 8%, 10% or 12% (v/v)  
127 with a final concentration of 1x10<sup>6</sup> CFU/mL. The kinetics of the fermentations was  
128 monitored for 7 days. After AF, L-malic acid was determined by enzymatic kits  
129 (Biogamma, Italy).

130

### 131 2.5 Pilot-scale vinification

132 The vinification was carried out in an experimental cellar using sterile stainless steel  
133 100-L vessels by inoculating 90 L of Primitivo must (18.9° Babo; pH 3.22; nitrogen  
134 176.4 g/L), as single or mixed inoculum with 10<sup>6</sup> CFU/mL of *H. uvarum* ITEM 8795,  
135 10<sup>4</sup> CFU/mL of *S. cerevisiae* ITEM 17292 or ITEM 17293 and 10<sup>6</sup> CFU/mL of *O. oeni*  
136 OT4. The dynamics of the alcoholic fermentation process was daily monitored and  
137 samples of wines were stored at -20 °C for further analyses.

138

### 139 2.6 Analytical determinations

140 The main product components (ethanol, residual sugars, pH, glucose, fructose, malic  
141 acid, lactic acid, tartaric acid, citric acid, volatile acidity, total acidity, glycerol brix,  
142 density, SO<sub>2</sub>, total polyphenols, antocyanins, CO<sub>2</sub>, absorbance at 420, 520 and 620 nm) of



143 wine and must under fermentation were evaluated by Fourier Transform Infrared  
144 Spectroscopy the (FTIR) by employing the WineScan Flex (FOSS Analytical, DK).  
145 Samples were centrifuged at 8000 rpm for 10 min and then analyzed following the  
146 supplier's instructions. The major volatile constituents [acetaldehyde, ethyl acetate, 2-  
147 methyl-1-propanol, 1-propanol, higher alcohols, acetoin] were determined by gas-  
148 chromatography according to Di Toro et al. (2015). The internal standard solution used  
149 was 4-methyl-2-pentanol. Identification and quantification of the volatile compounds by  
150 GC-MS were carried out using an internal standard as already described (Tufariello et  
151 al., 2014). Volatile compounds were extracted in triplicate by solid phase extraction  
152 (SPE) technique (Garofalo et al., 2018). The samples were injected into a DB-WAX  
153 capillary column (60m×0.25mm I.D., 0.25 µm film thickness; Agilent, USA) and then  
154 analyzed with a 6890N series gas chromatograph (Agilent, USA) equipped with an  
155 Agilent 5973 mass spectrometer selective detector (MSD). The analysis was performed  
156 as previously reported (Tufariello et al., 2014). Technological parameters were obtained  
157 as previously described (Tufariello et al., 2019).

158

### 159 *2.7 Determination of microbial population*

160 The enumeration of viable yeast cells during the fermentations was carried out on WL  
161 agar medium (Sigma-Aldrich, USA), that allowed to discriminate *S. cerevisiae* (large  
162 white colonies) and *H. uvarum* (green colonies) after 48 h incubation at 28 °C for. The  
163 counting of LAB viable cells was made in MRS supplemented with 10 mg/L  
164 cycloheximide (Sigma-Aldrich, USA) to avoid yeast growth, after 7 days incubation at  
165 28 °C.

166

### 167 *2.8. Sensory analysis*

168 The sensory analysis was performed by a panel composed of 5 professional experts,  
169 chosen among oenologists and producers involved in Negroamaro wine production. The  
170 judges were asked to assign a score for different parameters of the wines, such as  
171 gustatory-intensity, balance, acidity, body and gustatory-persistence, using a sensory  
172 analysis-tasting sheet with a scale ranging from 0 (absence of perception) to 3  
173 (maximum perception). The mean scores of attributes were submitted to Quantitative  
174 Descriptive Analysis (QDA) according to Trani and Coworkers (2016).

175

## 176 *2.8 Statistical analysis*

177 Chemical data were subjected to One-Way factor analysis of variance (ANOVA).  
178 Significant differences were separated using the Duncan test. The level of significance  
179 was set at  $P < 0.05$ . The comparison of volatile classes of compounds during  
180 fermentation was achieved by principal component analysis (PCA). All statistical  
181 analyses were carried out using the STATISTICA7.0 software (StatSoft software  
182 package, USA).

183

184

## 185 **3. Results**

186

### 187 *3.1 Yeast isolation and identification*

188 The oenological selection of autochthonous yeasts associated with natural fermentations  
189 of Negroamaro grapes, collected in the “Brindisi” PDO/DOC area, started with the  
190 isolation of 1200 yeast isolates. To this scope, serial dilutions of must and lees collected  
191 at the end of spontaneous fermentation were spread after on BIGGY agar. This selective  
192 medium allowed the isolation of 145 yeast colonies no or low H<sub>2</sub>S producers. The above

193 145 isolates were identified by molecular analysis of yeast rDNA, and they confirmed  
194 to belong to the species *Saccharomyces cerevisiae*. Then, 36 isolates randomly selected  
195 were characterized at strain level using a PCR-based assay, relying on the amplification  
196 of interdelta regions. The molecular fingerprint allowed the identification of 15 different  
197 *S. cerevisiae* strains (not shown). One representative biotype for each strain/profile has  
198 been selected. For these strains (P1, P2, P5, P6, P9, P13, P14, P20, P25, P28, P26, P33,  
199 P34, P35 and P32) the fermentative performances in wine were further studied.

200

### 201 3.2 Lab-scale vinifications

202 These technological and oenological parameters were mainly considered for the  
203 selection of autochthonous yeast strains: (i) acetic acid <0.6 g/L, (ii) residual sugars <2  
204 g/L and (iii) absence of H<sub>2</sub>S production. The primary screening indicated that, among  
205 the 15 selected different biotypes, the P2, P5, P13, P20, P25, P26, P28, P33, P34 and  
206 P35 complied to the above criteria and they were further characterized. Table 1 and  
207 Table 2 describe their principal technological and chemical features of the obtained  
208 wines. The presence of higher alcohols produced by fermentation in must was evaluated  
209 (Table 3). The latter ranged from 51.79 mg/L (strain P34) to 59.66 mg/L (strain P26),  
210 indicating that all strains could positively contribute to the aromatic complexity of wine.  
211 The ethyl acetate values ranged between 12.54 mg/L for P13 and 22.79 mg/L for P20  
212 (Table 3). Acetaldehyde concentrations ranged from 14.91 mg/L (strain P5) to 22.79  
213 mg/L (strain P28). The amount of acetoin, produced by the tested strains, ranged from  
214 1.62 mg/L for P26 to 4.96 mg/L for P34 (Table 3).

215 The fermented musts were also subjected to sensory analysis and the strains P25 and  
216 P28 obtained the maximum score with 12 and 13 points out of 15. The global evaluation  
217 of obtained data indicated that the P25 (ITEM 17292) and P28 (ITEM 17293) strains

218 were those denoted by the best fermentative properties and they were chosen for the co-  
219 inoculation trials.

220

### 221 3.3 Malolactic activity of *O. oeni* strains in the mixture culture

222 The selected *S. cerevisiae* ITEM 17292 and ITEM 17293 strains were co-inoculated  
223 with *H. uvarum* ITEM 8795 in Negroamaro grape must and the five selected  
224 autochthonous *O. oeni* strains were further investigated for their ability to consume L-  
225 malic acid by co-inoculating (0%) or sequentially inoculating them during AF, when  
226 ethanol content was 2%, 4%, 6%, 8%, 10% or 12% (v/v) (Fig. 1). Results showed that  
227 ethanol level at the moment of bacterial inoculation was crucial for developing MLF.  
228 The strategy of co-inoculation with *S. cerevisiae* and *H. uvarum* was the best strategy  
229 for maintaining highest *O. oeni* populations and therefore for carrying out MLF in red  
230 must. Only OT3 *O. oeni* strain co-inoculated with *H. uvarum* ITEM 8795 and *S.*  
231 *cerevisiae* ITEM 17292 (Fig. 1A) and OT25 *O. oeni* strain co-inoculated with *H.*  
232 *uvarum* ITEM 8795 and *S. cerevisiae* ITEM 17292 (Fig. 1G) or ITEM 17293 (Fig. 1H)  
233 were not consuming all L-malic acid present in the red must after 21 days of the end of  
234 the AF. All strains of *O. oeni* exhibited malolactic activity when they were inoculate in  
235 an ethanol concentration up to 4%. We observed that *O. oeni* strains have more  
236 difficulties to initiate MLF with 6 -12 % of ethanol. Among the *O. oeni* strains, OT4  
237 presented the highest malolactic activity, consuming completely the L-malic acid in all  
238 the ethanol concentrations studied (Fig. 1C and Fig. 1D). When inoculated at ethanol  
239 concentrations up to 6% (v/v), *O. oeni* OT4 completed the MLF in less than 7 (Fig. 1C)  
240 or in 21 days (Fig. 1D), when the *S. cerevisiae* strains ITEM 17292 and ITEM 17293  
241 were respectively used in the mixed starter formulation.

242

### 243 3.4 Kinetics of alcoholic fermentation in the multi-strain fermentations

244 In order to evaluate the effect of the inoculated microorganisms on the AF, the  
245 formation of ethanol was followed for 4 days. Significant differences ( $P= 0.0020$ ) were  
246 found in ethanol formation depending on the *S. cerevisiae* stain used (Fig. 2). The  
247 ethanol concentration was 12.86 % (v/v) and 12.12 % (v/v), respectively when ITEM  
248 17292 and ITEM 17293 were used in the co-inoculation tests. There were not  
249 significant no differences on the final ethanol concentration depending on the time of  
250 inoculation of the *O. oeni* strain. The concentration of ethanol in the produced wines  
251 was not influenced by the procedure adopted for the *O. oeni* OT4 strain inoculation, i.e.  
252 co-inoculation with *H. uvarum* and *S. cerevisiae* or inoculation at the end of the AF  
253 (Fig. 3). These findings were observed with all *O. oeni* strains used in the study (data  
254 not shown). Taken together, the above results indicated that the OT4 strain was the best-  
255 performing and it was chosen for the further co-inoculation assays.

256

### 257 3.5 Dynamics of yeast and bacterial population

258 After 24 h of fermentation, *H. uvarum* underwent a slight yeast concentration decrease  
259 and then increased fast (Fig. 4). *S. cerevisiae* ITEM 17292 reached the maximum yeast  
260 population after 24 h of the inoculation, increasing from  $1.00 \times 10^6$  CFU/mL to  $5.55 \times 10^7$   
261 CFU/mL. In this case *H. uvarum* ITEM 8795 presented a maximum concentration of  
262  $1.00 \times 10^7$  CFU/mL after 60 h of the inoculation, however after 72 hours of incubation,  
263 the population of this yeast descended drastically (Fig. 4A). When the *H. uvarum* strain  
264 was co-inoculated with *S. cerevisiae* ITEM 17293, it reached its maximum  
265 concentration after 48 h with a population of  $3.30 \times 10^7$  CFU/mL. The strain of *O. oeni*  
266 OT4 showed a similar trend in both trials: reached a population higher than  $1.00 \times 10^7$   
267 CFU/mL after 168 h of the inoculation (Fig. 4C and 4D) and kept constant until the end

268 of the fermentation. Moreover, after 168 h of incubation *O. oeni* OT4 inoculated with  
269 6% (v/v) of ethanol showed a cell viability of  $2.40 \times 10^7$  CFU/mL in combination with *S.*  
270 *cerevisiae* ITEM 17292 while with *S. cerevisiae* ITEM 17293 was  $7.50 \times 10^5$  CFU/mL,  
271 indicating the connection of the cell viability with the malolactic activity. *O. oeni* OT3,  
272 OT5, OT25 and OM22 after 168 h of inoculation only presented populations above  
273  $1 \times 10^6$  CFU/mL when were inoculated simultaneously to *S. cerevisiae* and *H. uvarum*  
274 (data not shown), explaining the reduced malolactic activity of these strains when were  
275 inoculated from 2% (v/v) of ethanol up to 12 % (v/v).

276

### 277 3.6 Pilot-scale vinifications

278 In order to evaluate the fermentation performance and interactions of mixed cultures at  
279 winery-scale, selected yeast strains of *S. cerevisiae* ITEM 17292 and ITEM 17293, *H.*  
280 *uvarum* (ITEM 8795) and the selected bacteria *O. oeni* (OT4), the following pilot-scale  
281 vinifications were carried out: Trial A: ITEM 17292; Trial B: ITEM 17292 + OT4; Trial  
282 C: ITEM 17292 + ITEM 8795; Trial D: ITEM 17292 + ITEM 8795 + OT4; Trial E:  
283 ITEM 17293; Trial F: *S. cerevisiae* ITEM 17293 + OT4; Trial G: ITEM 17293 + ITEM  
284 8795; Trial H: ITEM 17293 + ITEM 8795 + OT4.

285 The principal chemical parameters were analyzed by FT-IR (Table 4). In all the  
286 obtained fermented musts, volatile acidity, expressed as acetic acid, was quite low  
287 ranging from 0.27 g/L (trial D) to 0.41 g/L (trial H). The lower values of VA were  
288 detected in trial D (ITEM 17292 + *H. uvarum* + *O. oeni*) and trial H (ITEM 17293 + *H.*  
289 *uvarum* + *O. oeni*). A decrease in malic acid concentration coupled to increase of lactic  
290 acid content was achieved, 0.16 g/L in trial B 0.19 g/L in trial D, 0.13 g/L in trial F and  
291 finally 0.18 g/L in trial H.. The values of total acidity, tartaric acid and glycerol did not  
292 differ in the eight fermentations, indicating that the technique of co-inoculation does not

293 adversely affect the chemistry of the wine compared to the classical inoculation  
294 procedures (Table 1).

295 The GC-MS assay allowed the identification and quantification of 22 different volatile  
296 compounds (Table 5). The higher concentrations of alcohols were detected in trial D  
297 (59.04 mg/L), trial B (46.59 mg/L), trial C (39.07 mg/L) and trial H (32.56 mg/L). The  
298 esters were detected in higher concentrations in the same samples (A-B-C-H), while the  
299 acids content ranged from 1.0 mg/L (trial G) to 2.72 mg/L (trial H). Among esters,  
300 isoamyl acetate, ethyl lactate, ethyl octanoate, ethyl decanoate, diethyl succinate and  
301 mono ethyl succinate showed significant differences among the wines analyzed. When  
302 compared with the other obtained wines, the concentrations of these molecules was  
303 higher in the samples B, C and H. Moreover, the wine samples B, D, and H showed the  
304 higher amounts of hexanoic (ranging from 0.35 to 0.40 mg/L), octanoic (ranging from  
305 0.54 to 0.60 mg/L) and decanoic (ranging from 0.27 to 0.36 mg/L), acids.

306 The Principal Component Analysis (PCA) was performed on the concentrations of  
307 molecules detected by GC-MS in the produced wines (Fig. 5). Indeed, the wines from  
308 the trials D and H, both obtained by employing the *Saccharomyces*/non-  
309 *Saccharomyces/O. oeni* mixed starter, were located in the third and in the fourth  
310 quadrant, both areas characterized by high concentrations of volatiles respect to the  
311 others trials (E-F-G) located in the first quadrant. The wine from trial D showed in  
312 particular high values of isoamylalcohols, phenylethanol and ethyl lactate, while wines  
313 from the vinification H showed high values of isoamylacetate, ethyl hexanoate, 2-  
314 methylpropanol and 1-hexanol.

315 Taken together, the obtained outcome indicated that the *Saccharomyces*/non-  
316 *Saccharomyces/O. oeni* mixed starter formulations, detained the technological and  
317 enological features required for their possible use as industrial starter.

318

#### 319 **4. Discussion**

320 Two autochthonous *S. cerevisiae* strains (ITEM 17292 and ITEM 17293) were selected  
321 using the procedure described by Tufariello et al (2019). The two selected *S. cerevisiae*  
322 strains were always able to dominate the fermentation process and **to obtain a final**  
323 **product with an adequate chemical composition.** These strains were used for the co-  
324 inoculation trials to develop a mixed starter culture with non-*Saccharomyces* yeasts and  
325 LAB.

326 The addition of non-*Saccharomyces* yeast species as part of mixed starter formulations,  
327 has been indicated as a way **to** simulate the spontaneous fermentations (Petruzzi et al.,  
328 2017; Suzzi et al., 2012, Tristezza et al., 2016b), **thus** conferring particular organoleptic  
329 characteristics to wines without increasing the risks for wine quality and safety often  
330 associated with uncontrolled vinifications (Bergal, Spano, Tristezza, Grieco, &  
331 Capozzi, 2017; Capozzi et al., 2015). The performance of MLF by LAB is highly  
332 affected by the physicochemical intrinsic properties of wine, such as pH, ethanol, SO<sub>2</sub>  
333 and by yeast metabolism (Petruzzi et al., 2017). Alcoholic fermentation in wine  
334 undergoes deep chemical changes enhanced by ethanol concentrations over 4% (v/v)  
335 and can inhibit the growth of most LAB (Balmaseda, Bordons, Reguant & Bautista-  
336 Gallego, 2018). In our study, all strains showed better malolactic activity when *O. oeni*  
337 were co-inoculated (0 % ethanol v/v) with the selected yeasts or inoculated up to 4% of  
338 ethanol. Indeed, only *O. oeni* OT4 consumed all L-malic acid when-inoculated with an  
339 ethanol concentration above 4% **v/v**. Moreover, the obtained evidences indicated that  
340 the duration of MLF was reduced by the co-inoculation of yeasts and all the *O. oeni*  
341 strains investigated. Interaction with yeasts can be from inhibitory, to neutral of  
342 stimulatory depending on the release of nutrients by yeasts, and on the ability of yeasts



343 to produce metabolites that can affect LAB (Alexandre, Costello, Remize, Guzzo &  
344 Guilloux-Benatier, 2004). One of the main strategies to mitigate the possible inhibitory  
345 interactions that have been proposed is the; co-inoculation of yeast and *O. oeni*  
346 (Izquierdo-Cañas, Pérez-Martín, Romero, Prieto & Herreros, 2012).

347 Our findings confirmed data of previous studies (Ciani, et al., 2016; Maturano et al.,  
348 2018; Tristezza, et al., 2016b), by showing that grape musts co-inoculated with the  
349 mixed starter cultures presented less ethanol content than when single cultures of *S.*  
350 *cerevisiae* were employed. Besides, *H. uvarum* ITEM 8795 grew better in combination  
351 with *S. cerevisiae* ITEM 17293 than with *S. cerevisiae* ITEM 17292. Contrariwise, the  
352 *O. oeni* OT4, with best malolactic activity in grape must, presented a higher L-malic  
353 consumption rate and cell viability when *S. cerevisiae* ITEM 17292 was used. Lactic  
354 acid bacteria have complex nutrient requirements and so, their development depends on  
355 the nutrients consumption by the yeasts during the AF (Ivey et al., 2013). In accordance  
356 with the results of Curiel, Morales, Gonzalez & Tronchoni (2017), *O. oeni* OT4 showed  
357 lower malolactic activity and growth in fermentation trials where *H. uvarum* ITEM  
358 8795 showed higher population. The outcome achieved by the lab-scale tests were  
359 validated by carrying out pilot-scale vinification trials. It is interesting to highlight that,  
360 the presence of fermentable sugars did not affect the values of the volatile acidity, as  
361 reported in previous studies (Liu, 2012; Tristezza et al., 2016a). Our findings confirmed  
362 the use of yeast/bacteria mixed inoculums for the management of the MLF, not affected  
363 by the addition of the non-*Saccharomyces* starter strain and it had a positive influence  
364 on fermentation length and on aroma composition of wine (Muñoz, Beccaria & Abreo,  
365 2014). In fact, highly considerable was the effect of the mixed starter formulation on the  
366 aroma pattern of produced wines, compared to those obtained by inoculation of the *S.*  
367 *cerevisiae* starter alone. Recent investigations have highlighted the variation of the

368 biochemical profile of wine produced by different LAB inoculation procedures  
369 (Abrahamse & Bartowsky, 2011; Izquierdo-Cañas et al., 2012). Our data suggested, in  
370 accordance to literature (Antalick, Perello & de Revel 2013), that yeast/LAB co-  
371 inoculation could enhance the fruity aroma, thereby increasing the level of esters.  
372 Among alcohols identified, other higher alcohols shows higher values standing out 2-  
373 isoamyl alcohols and 2-phenyletanol. The higher alcohols increase were significantly  
374 higher when the fermentation was carried out by mix composed by *Saccharomyces*/non-  
375 *Saccharomyces*/*O. oeni* strains and they were significant different when one of the two  
376 *S. cerevisiae* strains (ITEM 17292 or ITEM 17293) were used.  
377 The combination of the three different microbial starters was responsible for the high  
378 esters production, contributing to improve wine flavor with fruity notes. In fact, the  
379 wines obtained by the pilot-scale trials D and H showed a higher concentration of  
380 hexanoic-octanoic and decanoic acids, which during the storage or aging could undergo  
381 to the esterification with the higher alcohols, thus increasing the fruity aroma (Francis,  
382 & Newton, 2005). Total alcohol and acid concentrations were found to be higher in  
383 wines produced by *Saccharomyces*/non-*Saccharomyces*/*O. oeni* co-inoculation, these  
384 compounds being responsible for fruity, sweet, winery and acid sensory notes in wine.  
385 In conclusion, the proposed approach can be very effective for the preparation of mixed  
386 starter culture formed by *Saccharomyces*, non-*Saccharomyces* yeasts and LAB. These  
387 mixed starter cultures represent a value solution to improve the specific attributes of  
388 typical regional wines. At the best of our knowledge, this investigation firstly illustrates  
389 the preparation and validation of a non-*Saccharomyces*/*Saccharomyces*/*O. oeni* mixed  
390 starter formulation that could be successfully adopted for the industrial production of  
391 typical Apulian red wines.

392

393 **Acknowledgements**

394 This research was partially supported by the Apulia Region in the framework of the  
395 Project DOMINA APULIAE (POR Puglia FESR – FSE 2014-2020-Azione 1.6. –  
396 InnoNetwork; Project code AGBGUK2). The authors thank Mr. Giovanni Colella for  
397 technical assistance and Prof. H. Smith for text proofreading.

398

399 **References**

- 400 Abrahamse, C.E., & Bartowsky, E.J. (2011). Timing of malolactic fermentation  
401 inoculation in Shiraz grape must and wine: influence on chemical composition.  
402 *World Journal of Microbiology and Biotechnology*, 28, 255-265.
- 403 Alexandre, H., Costello, P.J., Remize, F., Guzzo, J., & Guilloux-Benatier, M. (2004).  
404 *Saccharomyces cerevisiae-Oenococcus oeni* interactions in wine: current knowledge  
405 and perspectives. *International Journal of Food Microbiology*, 93, 141-154.
- 406 Antalick, G., Perello, M. C., & de Revel, G. (2013). Co-inoculation with yeast and LAB  
407 under winery conditions: modification of the aromatic profile of merlot wines. *South*  
408 *African Journal for Enology and Viticulture*, 34, 223-232.
- 409 Balmaseda, A., Bordons, A., Reguant, C., & Bautista-Gallego, J. (2018). Non-  
410 *Saccharomyces* in wine: effect upon *Oenococcus oeni* and malolactic fermentation.  
411 *Frontiers in Microbiology*, 9, 534.
- 412 Bartowsky, E.J., Costello, P.J., & Chambers, P.J. (2015). Emerging trends in the  
413 application of malolactic fermentation. *Australian Journal of Grape and Wine*  
414 *Research*, 21, 663-669.
- 415 Berbegal, C., Peña, N., Russo, P., Grieco, F., Pardo, I., Ferrer, S., Spano, G., Capozzi,  
416 V., 2016. Technological properties of *Lactobacillus plantarum* strains isolated from  
417 grape must fermentation. *Food Microbiology*, 57, 187-194.

418 Berbegal, C., Spano, G., Tristezza, M., Grieco, F., & Capozzi, V. (2017). Microbial  
419 resources and innovation in the wine production sector. *South African Journal for*  
420 *Enology and Viticulture*, 38, 156-166.

421 Brizuela, N.S., Bravo-Ferrada, B.M., Curilen, Y., Delfederico, L., Caballero, A.,  
422 Semorile, L., Pozo-Bayon, M.A., & Tymczyszyn, E.E. (2018). Advantages of using  
423 blend cultures of native *L. plantarum* and *O. oeni* strains to induce malolactic  
424 fermentation of Patagonian Malbec wine. *Frontiers in Microbiology*, 9, 2109.

425 Capozzi, V., Garofalo, C., Chiriatti, M.A., Grieco, F., & Spano, G. (2015). Microbial  
426 terroir and food innovation: The case of yeast biodiversity in wine. *Microbiological*  
427 *Research*, 181, 75-83.

428 Capozzi, V., Russo, P., Beneduce, L., Weidmann, S., Grieco, F., Guzzo, J., & Spano, G.  
429 (2010). Technological properties of *Oenococcus oeni* strains isolated from typical  
430 southern Italian wines. *Letters in Applied Microbiology*, 50, 327-334.

431 Capozzi, V., Russo, P., Lamontanara, A., Orrù, L., Cattivelli, L., & Spano, G. (2014).  
432 Genome sequences of five *Oenococcus oeni* strains isolated from Nero di Troia wine  
433 from the same terroir in Apulia, Southern Italy. *Genome Announcement*, 2, e01077-  
434 14.

435 Ciani, M., Beco, L., & Comitini, F. (2006). Fermentation behaviour and metabolic  
436 interactions of multistarter wine yeast fermentations. *International Journal of Food*  
437 *Microbiology*, 108, 9, 239-245.

438 Ciani, M., Comitini, F., Mannazzu, I., & Domizio, P. (2009). Controlled mixed culture  
439 fermentation: a new perspective on the use of non *Saccharomyces* yeasts in  
440 winemaking. *FEMS Yeast Research*, 10, 123-133.

441 Ciani, M., Morales, P., Comitini, F., Tronchoni, J., Canonico, L., Curiel, J.A., Oro, L.,  
442 Rodrigues, A.J., & Gonzalez, R. (2016). Non-conventional yeast species for lowering  
443 ethanol content of wines. *Frontiers in Microbiology*, 7, 642.

444 Comitini, F., Gobbi, M., Domizio, P., Romani, C., Lencioni, L., Mannazzu, I., & Ciani,  
445 M. (2011). Selected non-*Saccharomyces* wine yeasts in controlled multistarter  
446 fermentations with *Saccharomyces cerevisiae*. *Food Microbiology*, 28, 873-882.

447 Curiel, J.A., Morales, P., Gonzalez, R., & Tronchoni, J. (2017). Different non-  
448 *Saccharomyces* yeast species stimulate nutrient consumption in *S. cerevisiae* mixed  
449 cultures. *Frontiers in Microbiology*, 8, 2121.

450 De Benedictis, M., Bleve, G., Grieco, F., Tristezza, M., & Tufariello, M. (2011). An  
451 optimized procedure for the enological selection of non-*Saccharomyces* starter  
452 cultures. *Antonie Van Leeuwenhoek*, 99, 189-200.

453 Di Toro, M. R., Capozzi, V., Beneduce, L., Alexandre, H., Tristezza, M., Durante, M.,  
454 Tufariello, M., Grieco, F., & Spano, G. (2015). Intraspecific biodiversity and  
455 ‘spoilage potential’ of *Brettanomyces bruxellensis* in Apulian wines. *LWT-Food*  
456 *Science and Technology*, 60, 102-108.

457 Francis, I. L. & Newton, J. L. (2005). Determining wine aroma from compositional  
458 data. *Australian Journal of Grape and Wine Research*, 11, 114–126.

459 Garofalo, C., Berbegal, C., Grieco, F., Tufariello, M., Spano, G., & Capozzi, V. (2018).  
460 Selection of indigenous yeast strains for the production of sparkling wines from  
461 native Apulian grape varieties. *International Journal of Food Microbiology*, 285, 7-  
462 17.

463 Garofalo, C., Tristezza, M., Grieco, F., Spano, G., & Capozzi, V. (2016). From grape  
464 berries to wine: population dynamics of cultivable yeasts associated to “Nero di

465 Troia” autochthonous grape cultivar. *World Journal of Microbiology*  
466 *and Biotechnology*, 32, 59.

467 Hay, J., Grieco, F., Druka, A., Pinner, M., Lee, S. C., & Hull, R. (1994). Detection of  
468 rice tungro bacilliform virus gene products in vivo. *Virology*, 205, 430-437

469 Ivey, M., Massel, M., & Phister, T.G. (2013). Microbial Interactions in Food  
470 Fermentations. *Annual Review of Food Science and Technology*, 4, 141-162.

471 Izquierdo-Cañas, P.M., Pérez-Martín, F., Romero, E.G., Prieto, S.S., & Herreros,  
472 M.d.l.L.P. (2012). Influence of inoculation time of an autochthonous selected  
473 malolactic bacterium on volatile and sensory profile of Tempranillo and Merlot  
474 wines. *International Journal of Food Microbiology*, 156, 245-254.

475 Liu, P.-T., Lu, L., Duan, C.-Q., & Yan, G.-L. (2016). The contribution of indigenous  
476 non-*Saccharomyces* wine yeast to improved aromatic quality of Cabernet Sauvignon  
477 wines by spontaneous fermentation. *LWT- Food Science and Technology*, 71, 356-  
478 363.

479 Lopes, C.A., Rodriguez, M.E., Sangorrin, M., Querol, A., Caballero, A.C., 2007.  
480 Patagonian wines: the selection of an indigenous yeast starter. *Journal of Industrial*  
481 *Microbiology & Biotechnology*, 34, 539-546.

482 Maturano, Y.P., Mestre, M.V., Kuchen, B., Toro, M.E., Mercado, L.A., Vazquez, F., &  
483 Combina, M. (2018). Optimization of fermentation-relevant factors: A strategy to  
484 reduce ethanol in red wine by sequential culture of native yeasts *International*  
485 *Journal of Food Microbiology*, 289, 40-48.

486 Muñoz, V., Beccaria, B., & Abreo, E. (2014). Simultaneous and successive inoculations  
487 of yeasts and lactic acid bacteria on the fermentation of an unsulfited Tannat grape  
488 must. *Brazilian Journal of Microbiology*, 45, 59-66.

489 Petruzzi, L., Capozzi, V., Berbegal, C., Corbo, M.R., Bevilacqua, A., Spano, G., &  
490 Sinigaglia, M. (2017). Microbial resources and enological significance: opportunities  
491 and benefits. *Frontiers in Microbiology*, 8, 995.

492 Rodríguez, M.E., Infante, J.J., Molina, M., Domínguez, M., Rebordinos, L., & Cantoral,  
493 J.M. (2010). Genomic characterization and selection of wine yeast to conduct  
494 industrial fermentations of a white wine produced in a SW Spain winery. *Journal of*  
495 *Applied Microbiology*, 108, 1292-1302.

496 Suzzi, G., Arfelli, G., Schirone, M., Corsetti, A., Perpetuini, G., & Tofalo, R. (2012).  
497 Effect of grape indigenous *Saccharomyces cerevisiae* strains on Montepulciano  
498 d'Abruzzo red wine quality. *Food Research International*, 46, 22-29.

499 Tofalo, R., Patrignani, F., Lanciotti, R., Perpetuini, G., Schirone, M., Di Gianvito, P.,  
500 Pizzoni, D., Arfelli, G., & Suzzi, G. (2016). Aroma profile of Montepulciano  
501 d'Abruzzo wine fermented by single and co-culture starters of autochthonous  
502 *Saccharomyces* and non-*Saccharomyces* yeasts. *Frontiers in Microbiology*, 7, 610.

503 Trani, A., Verrastro, V., Punzi, R., Faccia, M., & Gambacorta, G. (2016). Phenols,  
504 Volatiles and Sensory Properties of Primitivo Wines from the "Gioia Del Colle"  
505 PDO Area. *South African Journal of Enology and Viticulture*, 37, 139-148.

506 Tristezza, M., di Feo, L., Tufariello, M., Grieco, F., Capozzi, V., Spano, G., & Mita, G.  
507 (2016). Simultaneous inoculation of yeasts and lactic acid bacteria: Effects on  
508 fermentation dynamics and chemical composition of Negroamaro wine. *LWT- Food*  
509 *Science and Technology*, 66, 406-412.

510 Tristezza, M., Gerardi, C., Logrieco, A., Grieco, F., 2009. An optimized protocol for the  
511 production of interdelta markers in *Saccharomyces cerevisiae* by using capillary  
512 electrophoresis. *Journal of Microbiological Methods*, 78, 286-291.

513 Tristezza, M., Tufariello, M., Capozzi, V., Spano, G., Mita, G., Grieco, F., 2016b. The  
514 oenological potential of *Hanseniaspora uvarum* in simultaneous and sequential co-  
515 fermentation with *Saccharomyces cerevisiae* for industrial wine production.  
516 *Frontiers in Microbiology*, 7, 670.

517 Tufariello, M., Chiriatti, M.A., Grieco, F., Perrotta, C., Capone, S., Rampino, P.,  
518 Tristezza, M., & Mita, G. (2014). Influence of autochthonous *Saccharomyces*  
519 *cerevisiae* strains on volatile profile of Negroamaro wines. *LWT- Food Science and*  
520 *Technology*, 58, 35-48.

521 Tufariello, M., Maiorano, G., Rampino, P., Spano, G., Grieco, F., Perrotta, C., &  
522 Capozzi, V. (2019). Selection of an autochthonous yeast starter culture for industrial  
523 production of Primitivo “Gioia del Colle” PDO/DOC in Apulia (Southern Italy).  
524 *LWT- Food Science and Technology*, 99, 188-196.

525



526 **Figure legends**

527

528 **Figure 1.** L-malic acid consumption (g/L) by *O. oeni* strains (OT3, OT4, OT5, OT25,  
529 OM22) after AF, when were co-inoculated (●) or sequentially inoculated during AF,  
530 when ethanol content was 2% (■), 4% (▲), 6% (▼), 8% (◆), 10% (●) or 12% (■) (v/v).

531

532 **Figure 2.** Ethanol content (% , v/v) formation during the must fermentations carried out  
533 by the co-inoculation of *H. uvarum* ITEM 8795, *S. cerevisiae* ITEM 17292 (●) or *S.*  
534 *cerevisiae* ITEM 17293 (■), and A; *O. oeni* OT3, B; *O. oeni* OT4, C; *O. oeni* OT5, D;  
535 *O. oeni* OT25 and E; *O. oeni* OM22

536

537 **Figure 3.** Ethanol content produced during the must fermentations carried out by: A; *S.*  
538 *cerevisiae* ITEM 17292, *H. uvarum* ITEM 8795 and *O. oeni* OT4 co-inoculated (●) and  
539 sequentially inoculated when ethanol content was 12% (v:v) (■), and B; *S. cerevisiae*  
540 ITEM 17293, *H. uvarum* ITEM 8795 and *O. oeni* OT4 co-inoculated (●) and  
541 sequentially inoculated when ethanol content was 12% (v:v) (■).

542

543 **Figure 4.** Viable cell count (CFU/mL) of: **A**; *S. cerevisiae* ITEM 17292 (●) and *H.*  
544 *uvarum* ITEM 8795 (■), and **B**; *S. cerevisiae* ITEM 17293 (●) and *H. uvarum* ITEM  
545 8795 (■) co-inoculated with *O. oeni* OT4 in red must. **C**; *O. oeni* OT4 (●) co-inoculated  
546 with *S. cerevisiae* ITEM 17292 and *H. uvarum* ITEM 8795 and, **D**; *O. oeni* OT4 (●) co-  
547 inoculated with *S. cerevisiae* ITEM 17293 and *H. uvarum* ITEM 8795.

548

549 **Figure 5.** Principal Component Analysis (PCA) performed employing the data obtained  
550 by the GC-MS analysis of the wines obtained by the pilot-scale vinifications

**Table 1.** Main oenological and technological properties determined in 15 autochthonous *S. cerevisiae* strains

Isolate	FP	AYC	AC	H <sub>2</sub> S	Foam
P1	0.04	0.62	12.6	++	-
P2	0.03	0.64	13.7	-	-
P5	0.03	0.64	14.0	-	-
P6	0.03	0.63	13.6	+	-
P9	0.04	0.59	12.5	+	-
P13	0.03	0.63	13.6	-	-
P14	0.03	0.63	13.6	+	++
P20	0.03	0.64	13.9	-	-
P25	0.04	0.64	13.8	-	-
P26	0.03	0.65	14.1	-	-
P28	0.03	0.65	14.1	-	-
P32	0.05	0.64	14.0	-	+
P33	0.03	0.65	14.0	-	-
P34	0.03	0.64	13.8	-	-
P35	0.03	0.65	14.1	-	-
Control	0.04	0.63	13.3		

Data, measured at the end of fermentation, represent the average of three replicates. FP fermentation purity [volatile acidity (g/L)/ethanol (% v/v)], AYC alcohol yield coefficient [alcohol (% v/v)/initial sugars (%) -Final sugars (%)], AC alcohol content (% v/v). H<sub>2</sub>S and foam production: absent (-); low (+), high (++), very high (+++).

Table 2

Table 2. Concentration of major chemical compounds in wines obtained with 15 autochthonous strain of *S. cerevisiae*.

Strain	Ethanol	Sugars	TA	VA	pH	Malic	Lactic	Tartaric	Citric	Glycerol
P 1	13.2±0.15	4.94±0.95 <sup>b</sup>	6.26±0.05	0.41±0.11	3.39±0.55	1.41±0.16	0.25±0.05	2.04±0.44	0.47±0.11	8.21±0.67
P 2	13.68±0.45	3.40±0.66 <sup>a</sup>	5.78±0.31	0.44±0.16	3.39±0.47	1.26±0.13	0.14±0.07	1.89±0.28	0.47±0.13	7.99±1.11
P 5	14.05±0.87	1.92±0.24 <sup>a</sup>	5.99±0.65	0.44±0.07	3.41±0.38	1.41±0.24	0.07±0.03	1.99±0.65	0.47±0.07	8.66±0.94
P 6	13.74±0.55	3.11±0.43 <sup>a</sup>	6.19±0.16	0.42±0.16	3.37±0.31	1.49±0.33	0.12±0.04	1.88±0.48	0.48±0.14	7.55±0.55
P 9	13.07±0.92	7.26±2.35 <sup>b</sup>	6.21±0.35	0.58±0.21	3.37±0.37	1.45±0.27	0.05±0.02	1.68±0.33	0.43±0.19	7.67±0.07
P 13	13.70±0.40	1.87±0.34 <sup>a</sup>	6.57±0.95	0.41±0.15	3.39±0.62	1.59±0.34	0.12±0.04	2.01±0.07	0.48±0.15	8.64±0.27
P 14	13.60±1.05	2.05±0.07 <sup>a</sup>	6.51±0.44	0.42±0.11	3.39±0.38	1.58±0.37	0.14±0.03	1.98±0.27	0.51±0.08	8.16±0.18
P 20	13.92±0.88	2.15±0.12 <sup>a</sup>	5.81±0.27	0.45±0.22	3.39±0.17	1.29±0.28	0.08±0.03	1.50±0.37	0.45±0.15	8.78±0.05
P 25	14.08±0.27	2.11±0.44 <sup>a</sup>	6.35±0.65	0.33±0.08	3.45±0.73	1.36±0.54	0.31±0.07	1.52±0.27	0.47±0.12	10.27±0.77
P 26	14.12±0.84	2.24±0.23 <sup>a</sup>	6.84±0.38	0.47±0.23	3.38±0.37	1.70±0.17	0.23±0.11	1.47±0.65	0.50±0.20	8.76±0.93
P 28	14.31±0.11	1.76±0.28 <sup>a</sup>	6.98±0.48	0.32±0.08	3.39±0.51	1.72±0.52	0.26±0.08	2.04±0.12	0.51±0.14	9.20±3.10
P 32	14.01±0.41	1.49±0.33 <sup>a</sup>	7.50±0.38	0.66±0.12	3.38±0.93	1.94±0.17	0.25±0.11	1.34±0.26	0.44±0.18	8.18±0.66
P 33	14.02±0.60	2.69±0.76 <sup>a</sup>	5.99±0.95	0.48±0.07	3.39±0.45	1.40±0.66	0.09±0.03	1.91±0.54	0.47±0.15	8.48±0.10
P 34	13.87±0.76	2.36±0.27 <sup>a</sup>	6.06±0.55	0.47±0.08	3.42±0.61	1.25±0.27	0.13±0.04	2.24±0.38	0.43±0.12	8.37±0.65
P 35	14.26±0.36	3.17±0.94 <sup>a</sup>	6.33±0.95	0.44±0.11	3.41±0.75	1.41±0.52	0.09±0.03	2.27±0.25	0.46±0.25	8.36±0.05

TA, total acidity. VA, volatile acidity. Values are expressed in g/L. The ethanol concentration is expressed in g/100mL. Results are the mean of three injections of each replicate (n = 9); the standard deviation values (±) are indicated. Different letters in the column denote significant differences between yeast strains, at p < 0.05

**Table 3.** Concentration of selected volatile compounds determined by GC in wines obtained with the with 15 autochthonous strain of *S. cerevisiae*

Strain	acetaldehyde	ethyl acetate	1-propanol	2-metil-1-propanol	higher alcohols	acetoin
P2	17.58±0.55 <sup>a</sup>	15.27±.057 <sup>b</sup>	10.33±0.79 <sup>d</sup>	4.38±0.37 <sup>a</sup>	58.11±0.48 <sup>b</sup>	2.59±0.09 <sup>a</sup>
P5	14.91±0.61 <sup>a</sup>	21.33±0.69 <sup>d</sup>	12.54±0.53 <sup>e</sup>	4.19±0.62 <sup>a</sup>	55.80±1.41 <sup>a</sup>	2.10±0.17 <sup>a</sup>
P13	16.25±1.50 <sup>a</sup>	12.54±0.45 <sup>b</sup>	5.91±0.15 <sup>b</sup>	4.52±0.41 <sup>a</sup>	56.09±0.75 <sup>a</sup>	3.60±0.50 <sup>b</sup>
P20	16.89±0.12 <sup>a</sup>	22.79±0.25 <sup>d</sup>	12.91±0.25 <sup>e</sup>	3.70±0.24 <sup>a</sup>	61.20±1.07 <sup>b</sup>	1.92±0.16 <sup>a</sup>
P25	21.89±1.02 <sup>b</sup>	16.49±0.08 <sup>b</sup>	8.86±0.10 <sup>c</sup>	9.25±0.19 <sup>b</sup>	56.19±0.43 <sup>a</sup>	1.88±0.12 <sup>a</sup>
P26	15.72±0.53 <sup>a</sup>	22.03±0.91 <sup>d</sup>	11.21±0.33 <sup>d</sup>	3.91±0.25 <sup>a</sup>	59.66±0.12 <sup>b</sup>	1.62±0.15 <sup>a</sup>
P28	22.79±0.30 <sup>b</sup>	18.30±0.35 <sup>c</sup>	7.97±0.52 <sup>c</sup>	3.02±0.48 <sup>a</sup>	58.15±1.66 <sup>a</sup>	2.74±0.19 <sup>a</sup>
P33	15.09±0.83 <sup>a</sup>	3.28±0.67 <sup>a</sup>	5.36±0.41 <sup>b</sup>	4.61±0.32 <sup>a</sup>	55.83±0.54 <sup>a</sup>	4.17±0.12 <sup>b</sup>
P34	21.49±1.13 <sup>b</sup>	17.43±0.83 <sup>c</sup>	8.52±0.11 <sup>c</sup>	4.55±0.10 <sup>a</sup>	51.79±1.47 <sup>a</sup>	4.96±0.17 <sup>b</sup>
P35	24.97±0.53 <sup>c</sup>	19.49±0.64 <sup>c</sup>	3.55±0.06 <sup>a</sup>	2.44±0.29 <sup>a</sup>	58.56±0.44 <sup>b</sup>	2.39±0.24 <sup>a</sup>

Compound concentration (mg/L). Values are the mean of three injections of each replicate (n = 9); the standard deviation values (±) are indicated. Different letters in the row denote significant differences between yeast strains, at p < 0.05.

**Table 4.** Concentration of major chemical compounds in wines obtained by the pilot-scale vinifications

<b>Trial</b>	<b>Alcohol</b>	<b>Sugars</b>	<b>TA</b>	<b>VA</b>	<b>pH</b>	<b>Malic acid</b>	<b>Lactic acid</b>	<b>Tartaric acid</b>	<b>Glycerol</b>
A	11.89±0.56	1.25±0.12	7.56±0.56	0.36±0.07	3.19±0.36	2.75±0.56	0.26±0.06	4.19±0.10	9.50±0.60
B	11.80±0.10	1.24±0.07	7.17±0.10	0.35±0.05	3.24±0.26	0.16±0.05	1.83±0.14	4.19±0.25	9.05±0.87
C	11.97±0.22	1.23±0.16	7.41±0.76	0.35±0.10	3.19±0.24	2.7±0.76	0.19±0.05	4.2±0.14	9.61±0.87
D	12.26±0.84	1.29±0.23	7.02±0.48	0.27±0.08	3.27±0.26	0.19±0.04	1.97±0.16	4.14±0.15	9.12±0.56
E	11.70±0.17	1.19±0.17	7.82±0.86	0.33±0.08	3.20±0.28	2.78±0.55	0.26±0.05	4.12±0.26	9.31±0.67
F	11.85±0.54	1.19±0.07	7.52±0.66	0.41±0.06	3.31±0.15	0.13±0.04	1.99±0.07	3.93±0.24	9.25±0.38
G	11.99±0.11	1.22±0.34	7.35±0.10	0.41±0.12	3.27±0.18	2.76±0.85	0.04±0.02	4.14±0.20	9.42±0.33
H	12.56±0.10	1.29±0.41	7.62±0.77	0.29±0.07	3.21±0.16	0.18±0.04	1.89±0.15	4.11±0.33	9.03±0.94

TA; total acidity. VA; volatile acidity. Values are expressed in g/L. The ethanol concentration is expressed in g/100mL. Results are the mean of three injections of each replicate (n = 9); the standard deviation values (±) are indicated. No significant differences were detected at  $p < 0.05$ .

Table 5

Table 5: Concentration of selected volatile compounds determined by GC-MS in wine obtained by the pilot-scale vinifications

	<b>Trial A</b>	<b>Trial B</b>	<b>Trial C</b>	<b>Trial D</b>	<b>Trial E</b>	<b>Trial F</b>	<b>Trial G</b>	<b>Trial H</b>
	<i>mg/L±sd</i>							
<b>ALCOHOLS</b>								
2-Methyl-1-propanol	0.49±0.11 <sup>b</sup>	0.99±0.23 <sup>b</sup>	0.56±0.18 <sup>b</sup>	0.87±0.24 <sup>b</sup>	0.04±0.01 <sup>a</sup>	0.020±0.01 <sup>a</sup>	0.019±0.04 <sup>a</sup>	1.33±0.22 <sup>a</sup>
Isoamylalcohols	13.88±3.67 <sup>a</sup>	24.95±5.62 <sup>b</sup>	19.21±5.55 <sup>b</sup>	26.40±5.18 <sup>b</sup>	7.20±2.11 <sup>a</sup>	9.20±2.55 <sup>a</sup>	7.40±1.87 <sup>a</sup>	15.95±4.16 <sup>b</sup>
1-Hexanol	0.29±0.07 <sup>a</sup>	0.64±0.12 <sup>b</sup>	0.27±0.12 <sup>a</sup>	0.44±0.13 <sup>b</sup>	0.02±0.01 <sup>a</sup>	0.020±0.011 <sup>a</sup>	0.019±0.04 <sup>a</sup>	0.63±0.22 <sup>b</sup>
3-Hexen-1-ol (E)	0.54±0.11	0.67±0.22	0.74±0.21	0.81±0.25	0.46±0.18	0.94±0.26	0.22±0.08	0.67±0.21
3-Hexen-1-ol (Z)	0.011±0.06	0.03±0.01	0.014±0.04	0.02±0.01	nd	nd	nd	0.024±0.09
1-heptanol	nd	nd	0.28±0.10	0.56±0.17	nd	nd	nd	0.74±0.21
Methyionol	0.03±0.01 <sup>a</sup>	nd	0.033±0.011 <sup>a</sup>	0.76±0.23 <sup>a</sup>	nd	nd	nd	1.65±0.37 <sup>b</sup>
Phenylethanol	14.91±4.52 <sup>b</sup>	19.31±4.94 <sup>b</sup>	17.96±5.38 <sup>b</sup>	29.17±4.56 <sup>c</sup>	8.04±2.77 <sup>a</sup>	9.06±2.10 <sup>a</sup>	8.70±2.56 <sup>a</sup>	11.56±4.38 <sup>a</sup>
<b>TOTAL</b>	<b>30.15</b>	<b>46.59</b>	<b>39.07</b>	<b>59.04</b>	<b>15.77</b>	<b>19.24</b>	<b>16.34</b>	<b>32.56</b>
<b>ESTERS</b>								
Isoamyl-acetate	3.77±0.95 <sup>a</sup>	2.85±0.74 <sup>a</sup>	4.11±0.65 <sup>a</sup>	5.28±1.56 <sup>b</sup>	2.11±0.54 <sup>a</sup>	2.96±0.16 <sup>a</sup>	3.11±0.25 <sup>a</sup>	3.85±0.94 <sup>a</sup>
Ethyl-hexanoate	0.028±0.011	0.05±0.02	0.02±0.01	0.08±0.02	nd	nd	nd	0.09±0.03
Ethyl-lactate	0.14±0.05	1.38±0.17	0.11±0.03	1.18±0.44	0.22±0.08	2.76±0.94	0.01±0.01	2.13±0.76
Ethyl-octanoate	0.07±0.02	0.11±0.04	0.05±0.02	0.072±0.013	nd	nd	nd	0.13±0.03
3-Hydroxy-ethylbutanoate	nd	0.04±0.02	0.012±0.04	0.025±0.010	nd	nd	nd	nd
Ethyl-decanoate	0.94±0.34 <sup>a</sup>	0.83±0.14 <sup>a</sup>	0.77±0.26 <sup>a</sup>	2.67±0.34 <sup>c</sup>	0.95±0.26 <sup>a</sup>	1.76±0.38 <sup>b</sup>	1.88±0.25 <sup>b</sup>	2.46±0.84 <sup>c</sup>
Diethyl-succinate	0.45±0.12 <sup>a</sup>	0.07±0.02 <sup>a</sup>	0.40±0.17 <sup>a</sup>	0.67±0.28 <sup>b</sup>	0.22±0.07 <sup>a</sup>	0.32±0.08 <sup>a</sup>	0.21±0.06 <sup>a</sup>	0.69±0.19 <sup>b</sup>
Phenyl-acetate	0.18±0.06	0.24±0.10	0.21±0.06	0.22±0.06	nd	0.01±0.01	0.01±0.01	0.19±0.05
Monoethyl-succinate	2.37±0.94 <sup>b</sup>	3.84±0.84 <sup>b</sup>	4.42±1.45 <sup>b</sup>	6.04±2.67 <sup>c</sup>	1.09±0.27 <sup>a</sup>	1.13±0.16 <sup>a</sup>	1.07±0.16 <sup>a</sup>	3.11±0.83 <sup>b</sup>
<b>TOTAL</b>	<b>7.94</b>	<b>9.41</b>	<b>10.12</b>	<b>16.24</b>	<b>4.60</b>	<b>8.94</b>	<b>6.28</b>	<b>12.66</b>
<b>ACIDS</b>								
2-Methyl propanoic acid	0.52±0.18 <sup>a</sup>	0.45±0.17 <sup>a</sup>	0.77±0.23 <sup>a</sup>	0.95±0.34 <sup>a</sup>	0.77±0.15 <sup>a</sup>	0.65±0.18 <sup>a</sup>	0.47±0.12 <sup>a</sup>	1.56±0.27 <sup>b</sup>
Hexanoic acid	0.30±0.08 <sup>b</sup>	0.40±0.16 <sup>b</sup>	0.29±0.11 <sup>a</sup>	0.36±0.12 <sup>a</sup>	0.01±0.01 <sup>a</sup>	0.02±0.05 <sup>a</sup>	0.01±0.01 <sup>a</sup>	0.35±0.08 <sup>a</sup>

Octanoic acid	0.49±0.16	0.60±0.23	0.46±0.13	0.56±0.25	0.41±0.07	0.53±0.17	0.41±0.16	0.54±0.12
Decanoic acid	0.18±0.05	0.36±0.14	0.18±0.04	0.27±0.08	0.13±0.04	0.21±0.06	0.10±0.03	0.27±0.08
<b>TOTAL</b>	<b>1.49</b>	<b>1.89</b>	<b>1.70</b>	<b>2.14</b>	<b>1.31</b>	<b>1.41</b>	<b>1.00</b>	<b>2.72</b>

**TERPENS**

Citronellol	0.76±0.17	n.d.	n.d.	1.56±0.34	n.d.	n.d.	n.d.	0.73±0.21
-------------	-----------	------	------	-----------	------	------	------	-----------

Each value is expressed in mg/L. Results are the mean of three injections of each replicate (n = 9); the standard deviation values (±) are indicated. Different upper letters in row means significant differences at P < 0.05.

Figure 1  
[Click here to download high resolution image](#)






Figure 2  
[Click here to download high resolution image](#)




Figure 3  
[Click here to download high resolution image](#)





Figure 4  
[Click here to download high resolution image](#)







Figure 5  
[Click here to download high resolution image](#)

