

POSTER PRESENTATION

Open Access

# Critical connectivity for emergence of collective oscillations in strongly diluted neural networks

Stefano Luccioli<sup>1,2,3\*</sup>, Simona Olmi<sup>1,2,3</sup>, Antonio Politi<sup>4</sup>, Alessandro Torcini<sup>1,2,3</sup>

From Twenty Second Annual Computational Neuroscience Meeting: CNS\*2013  
Paris, France. 13-18 July 2013

Among the most relevant dynamical phenomena observed in brain circuits is the rhythmic collective behavior of neuronal populations [1]. In this work [2] we studied the dynamics of random neural networks models focusing on the role played by the (in-degree) connectivity  $K$  (i.e., the number of incoming connections per node) on the onset of collective oscillations. In modeling neural networks two classes of systems are generally considered [3]: massive networks, where  $K$  is proportional to the network size  $N$ ; *sparse* (or *strongly diluted*) networks, where  $K \ll N$ , and specifically  $K$  is independent on  $N$  as  $N \rightarrow \infty$ .

While it is not surprising to observe the onset of a collective motion in massive networks, it is less obvious to predict whether and when this can happen in sparse ones. Here, we showed that a *finite critical connectivity*  $K_c$  is able to sustain the emergence of collective oscillations and that this is a general and robust property of *sparse* networks. Since  $K_c$  turns out to be surprisingly of the order of a few tens in all models we have investigated, macroscopic motion appears to be rather ubiquitous and relevant in the context of neural dynamics. The existence of a critical connectivity separating asynchronous from coherent activity is similar to what experimentally observed in neuronal cultures [4].

Moreover, we showed that the microscopic evolution of sparse networks is extensive (i.e. the number of active degrees of freedom is proportional to the number of network elements) according to what observed for the  $\Theta$ -neuron model in Ref. [5]. This property is highly non-trivial, as the dynamics of a sparse network is intrinsically non additive [6] (it cannot be approximated with the juxtaposition of almost independent sub-structures).

We found all the above striking results to hold for networks of pulse-coupled leaky-integrate-and-fire neurons, among the most popular and yet simple models used in computational neuroscience, and more generally also for other kinds of networks (chaotic maps and Stuart-Landau oscillators).

#### Acknowledgements

The authors acknowledge funding support from the Joint Italian-Israeli Laboratory on Neuroscience funded by the Italian Ministry of Foreign Affairs and the European Commission through the Marie Curie Initial Training Network 'NETT', project 289146.

#### Author details

<sup>1</sup>CNR - Consiglio Nazionale delle Ricerche - Istituto dei Sistemi Complessi, via Madonna del Piano 10, I-50019 Sesto Fiorentino, Italy. <sup>2</sup>INFN Sez. Firenze, via Sansone, 1 - I-50019 Sesto Fiorentino, Italy. <sup>3</sup>Centro Interdipartimentale per lo Studio delle Dinamiche Complesse, via Sansone, 1 - I-50019 Sesto Fiorentino, Italy. <sup>4</sup>Institute for Complex Systems and Mathematical Biology and SUPA, University of Aberdeen, Aberdeen AB24 3UE, UK.

Published: 8 July 2013

#### References

1. Buzsáki G: *Rhythms of the Brain* Oxford University Press, New York; 2006.
2. Luccioli S, Olmi S, Politi A, Torcini A: **Collective dynamics in sparse networks.** *Phys Rev Lett* 2012, **109**:138103.
3. Golomb D, Hansel D, Mato G: *Handbook of Biological Physics* 2001, **4**:887, Elsevier Science.
4. Soriano J, Rodriguez Martinez M, Tlusty T, Moses E: **Development of input connections in neural cultures.** *Proc Natl Acad Sci USA* 2008, **105**:13758-13763.
5. Monteforte M, Wolf F: **Dynamical Entropy Production in Spiking Neuron Networks in the Balanced State.** *Phys Rev Lett* 2010, **105**:268104.
6. Liao W: **Graph-Bipartitioning Problem.** *Phys Rev Lett* 1987, **59**:1625-1628.

doi:10.1186/1471-2202-14-S1-P394

**Cite this article as:** Luccioli et al.: Critical connectivity for emergence of collective oscillations in strongly diluted neural networks. *BMC Neuroscience* 2013 **14**(Suppl 1):P394.

\* Correspondence: stefano.luccioli@fi.isc.cnr.it

<sup>1</sup>CNR - Consiglio Nazionale delle Ricerche - Istituto dei Sistemi Complessi, via Madonna del Piano 10, I-50019 Sesto Fiorentino, Italy

Full list of author information is available at the end of the article