

# Evaluation of the effects of different diets on microbiome diversity and fatty acid composition of rumen liquor in dairy goat

P. Cremonesi<sup>1†a</sup>, G. Conte<sup>2a</sup>, M. Severgnini<sup>3</sup>, F. Turri<sup>1</sup>, A. Monni<sup>1</sup>, E. Capra<sup>1</sup>, L. Rapetti<sup>4</sup>, S. Colombini<sup>4</sup>, S. Chessa<sup>1</sup>, G. Battelli<sup>5</sup>, S. P. Alves<sup>6</sup>, M. Mele<sup>2</sup> and B. Castiglioni<sup>1</sup>

<sup>1</sup>Istituto di Biologia e Biotecnologia Agraria, Consiglio Nazionale delle Ricerche, SS-Lodi, via Einstein, 26900 Lodi, Italy; <sup>2</sup>Dipartimento di Scienze Agrarie, Alimentari e Agro-ambientali (DISAAA-a), Università di Pisa, via del Borghetto, 80, 56124 Pisa, Italy; <sup>3</sup>Istituto di Tecnologie Biomediche, Consiglio Nazionale delle Ricerche, via Fratelli Cervi, 93, 20090 Segrate, Milano, Italy; <sup>4</sup>Dipartimento di Scienze Agrarie e Ambientali (DISAA) – Produzione, Territorio, Agroenergia, Università degli Studi di Milano, via Celoria, 2, 20133 Milano, Italy; <sup>5</sup>Istituto di Scienze delle Produzioni Alimentari, Consiglio Nazionale delle Ricerche, via Celoria, 2, 20133 Milano, Italy; <sup>6</sup>CIISA, Centro de Investigação interdisciplinar em sanidade animal, faculdade de medicina veterinaria, universidade de lisboa; avenida da universidade tecnica, 1300-477, Lisboa, Portugal

(Received 1 December 2016; Accepted 13 November 2017)

*Fat supplementation plays an important role in defining milk fatty acids (FA) composition of ruminant products. The use of sources rich in linoleic and  $\alpha$ -linolenic acid favors the accumulation of conjugated linoleic acids isomers, increasing the healthy properties of milk. Ruminal microbiota plays a pivotal role in defining milk FA composition, and its profile is affected by diet composition. The aim of this study was to investigate the responses of rumen FA production and microbial structure to hemp or linseed supplementation in diets of dairy goats. Ruminal microbiota composition was determined by 16S amplicon sequencing, whereas FA composition was obtained by gas-chromatography technique. In all, 18 pluriparous Alpine goats fed the same pre-treatment diet for  $40 \pm 7$  days were, then, arranged to three dietary treatments consisting of control, linseed and hemp seeds supplemented diets. Independently from sampling time and diets, bacterial community of ruminal fluid was dominated by Bacteroidetes (about 61.2%) and Firmicutes (24.2%) with a high abundance of Prevotellaceae (41.0%) and Veillonellaceae (9.4%) and a low presence of Ruminococcaceae (5.0%) and Lachnospiraceae (4.3%). Linseed supplementation affected ruminal bacteria population, with a significant reduction of biodiversity; in particular, relative abundance of Prevotella was reduced ( $-12.0\%$ ), whereas that of Succinivibrio and Fibrobacter was increased ( $+50.0\%$  and  $+75.0\%$ , respectively). No statistically significant differences were found among the average relative abundance of archaeal genera between each dietary group. Moreover, the addition of linseed and hemp seed induced significant changes in FA concentration in the rumen, as a consequence of shift from C18:2n-6 to C18:3n-3 biohydrogenation pathway. Furthermore, dimethylacetal composition was affected by fat supplementation, as consequence of ruminal bacteria population modification. Finally, the association study between the rumen FA profile and the bacterial microbiome revealed that Fibrobacteriaceae is the bacterial family showing the highest and significant correlation with FA involved in the biohydrogenation pathway of C18:3n-3.*

**Keywords:** metagenome, goat, rumen, fatty acid, dimethylacetal

## Implications

In this study, Next Generation Sequencing (NGS) and gas chromatography (GC) techniques assessed the effects of diets with fat supplementation on rumen microbial community and ruminal fatty acids (FA), respectively, in 18 Alpine goats. A treatment period of 100 days with supplemented feed showed a change in the composition of rumen microbial community with a significant effect on biodiversity of the ruminal bacterial community in goat and a relevant

repercussion on lipid metabolism of rumen, especially on biohydrogenation (BH) and plasmalogens synthesis.

## Introduction

In the last 20 years, the use of different polyunsaturated lipid source in the diet of dairy ruminant has been extensively adopted in order to modify the milk fatty acid composition toward a more desirable profile for human health (Mele, 2009). The efficiency of the FA transfer from diet to milk depends on several factors, including the nature of the fat supplementation and the interaction between fat source and

<sup>a</sup> These authors contributed equally to this work.

<sup>†</sup> E-mail: cremonesi@ibba.cnr.it

rumen microbes. Dietary polyunsaturated FA, in fact, are massively converted by lipolysis and BH processes into stearic acid (C18:0), after a previously lipolysis of triglycerides and phospholipids. Moreover, lipid supplements may interact with cellulolytic bacteria, interfering with fiber degradation (Chilliard *et al.*, 2003). Several *in vitro* and *in vivo* studies, aiming at evaluating the interaction between fat supplements and rumen microbes, demonstrated that FA composition of rumen liquor may be informative (Buccioni *et al.*, 2012). For instance, differences in C18:3, C18:2 and C18:1 isomers content of rumen liquor may be associated to alternative BH pathways (Shingfield *et al.*, 2010). At the same time, the content of specific FA originated by different metabolisms of rumen microbes (i.e. odd and branched chain FA) or the presence of plasmalogenic lipids that are components of cell membranes of microbes have been associated to specific dietary regimens (Vlaeminck *et al.*, 2006; Alves *et al.*, 2013). Fievez *et al.* (2012) proposed a predictive model to estimate ruminal activity, considering milk odd and branched chain FA. More recently, the introduction of high-throughput sequencing techniques, utilizing the 16S gene as universal target (Frey *et al.*, 2010), has led to many advances in rumen microbial communities characterization, as culture-dependent methods only detect around 11% of species present in this ecosystem (Fernando *et al.*, 2010). With these technologies, studies on: rumen metagenome composition (Cunha *et al.*, 2011) and microflora changes due to different diet and supplementation (Toral *et al.*, 2016) have been published for different species, mainly for cattle and sheep, whereas little information is available for goat.

The present study was designed to assess the responses and the associations between the microbiome structure and rumen FA production due to linoleic acid (hemp seeds) or  $\alpha$ -linolenic acid (linseed) supplemented diets in 18 Alpine goats, using NGS and GC techniques, respectively.

## Material and methods

### Animal selection and experimental design

The experiment was conducted at the Research Centre 'Cascina Baciocca' of the Department of Agricultural and Environmental Sciences, University of Milan in Cornaredo (Milan, Italy). Trial animals were handled as outlined by the guidelines of the Italian law on animal welfare for experimental animals (Legislative Decree 116/92) and of the University of Milan Ethics Committee for animal use and care.

In all, 18 Alpine goats were selected among 150 secondiparous goats reared at 'G.P. Guidobono Cavalchini' experimental farm of the University of Milan in Borgo Adorno (Cantalupo Ligure, Alessandria, Italy). All goats were unaffected by caprine arthritis-encephalitis virus and mastitis and were daughters of 12 bucks. The 18 goats were selected considering in sequence: production at their first lactation (only goats producing similar milk yield were considered); body condition evaluation (only goats with good to high evaluation were considered); pedigree (among the remaining

animals, goats were selected to be daughter of the highest number of bucks as possible, resulting in seven bucks) in order to minimize their total phenotypic diversity and maximize genetic diversity among the goats. Moreover, a genomic analysis was conducted using the Goat 50K SNP chip (Illumina, San Diego, CA, USA) for the assessment of genetic diversity (identity by state) to assign the goats to the three diet groups avoiding including too similar animals in the same group. Then, the three groups were assigned randomly to the three different dietary treatments.

The diets (Tables 1 and 2) were formulated to meet the protein and energy requirements of lactating goats weighing 65 kg and producing 2.8 kg of milk/day.

**Table 1** Composition of the experimental control (C), linseed (L) or hemp (H) diets (% on dry matter)

Ingredients	C	L	H
Meadow hay second cut <sup>1</sup>	24.4	24.3	24.3
Meadow hay first cut <sup>1</sup>	14.0	14.0	14.0
Alfalfa hay third cut <sup>1</sup>	10.5	10.5	10.5
Corn flakes <sup>2</sup>	18.6	13.1	13.1
Corn meal <sup>1</sup>	5.5	5.5	5.5
Soybean hulls <sup>2</sup>	13.8	13.8	13.8
Solvent Soybean meal <sup>2</sup>	6.6	2.9	2.9
Carob pulp <sup>1</sup>	4.5	4.5	4.5
Cane molasses <sup>3</sup>	0.4	0.4	0.4
Salts <sup>3,4</sup>	1.4	1.4	1.4
Vitamin mix (MecoVit-Metagro) <sup>3</sup>	0.3	0.3	0.3
Linseeds <sup>3</sup>	–	9.3	–
Hemp seeds <sup>3</sup>	–	–	9.3

<sup>1</sup>Ingredients of the basal ration.

<sup>2</sup>Ingredients of both basal ration and supplement concentrates.

<sup>3</sup>Ingredients of the concentrate supplements.

<sup>4</sup>Salts: 35% calcium carbonate, 32% sodium bicarbonate, 15% dicalcium phosphate, 13% sodium chloride, 5% magnesium oxide.

**Table 2** Chemical composition of the administered control (C), linseed (L) or hemp (H) diets

	C	L	H
Organic matter (OM) (%)	92.1	91.9	92.2
CP	13.4	13.4	13.7
Ether extract (EE) g/100 g DM	2.24	5.96	5.12
NDF g/100 g DM	39.3	39.9	41.6
NFC	37.1	32.7	31.9
Forage/concentrate ratio	49/51	49/51	49/51
C16:0	28.01	13.14	17.50
C18:0	16.89	7.34	8.64
C18:1 <i>cis</i> 9	18.25	21.58	18.77
C18:2n-6 LA	28.54	20.41	43.46
C18:3n-6 GLA	0.06	0.14	1.06
C18:3n-3 ALA	8.05	37.30	10.28
C18:4n-3 SDA	0.05	0.04	0.27
Other fatty acids	0.15	0.05	0.02
n-6/n-3	3.53	0.55	4.22

DM = dry matter; NFC = Non-Fiber Carbohydrates, calculated as follows: (NFC = OM – (CP + EE + NDF)); LA = linoleic acid; GLA =  $\gamma$ -linolenic acid; ALA =  $\alpha$ -linolenic acid; SDA = stearidonic acid.

The treatments were arranged to evaluate the main effects of two dietary lipid supplementations (linseed or hemp), on rumen population and its lipid metabolism as follows: (1) control diet (C), (2) diet supplemented with linseed (L), (3) diet supplemented with hemp seed (H). In order to balance for the fat supplementation in the two experimental diets, corn flakes and soybean meal were included in the C diet.

After calving, the 18 animals were divided into three different boxes and fed for the  $40 \pm 7$  (SD) days *postpartum* with a transition diet (after-calving diet; time point = T0), very similar to the control diet fed in T1, but with a slightly higher forage to concentrate ratio (51/49 v. 49/51, for T0 and T1, respectively), allowing management of the administration of independent rations. Thereafter, until  $140 \pm 7$  days of lactation (time point = T1), goats were fed a common basal ration plus three different concentrate supplements corresponding to different dietary treatments. The basal ration, containing the forages and some concentrate raw materials (Table 1), was prepared with a mixer wagon to allow a reduction of forage particle size. On dry matter (DM) basis, it accounted for 85.6% of total diet. The concentrate supplements (14.4% on total dietary DM) were administered individually twice daily (08.30 and 17.30 h) about half an hour before the basal ration. The goats were fed *ad libitum* and the orts were recorded once daily. The feeding rate was adjusted to yield orts on the basis of at least 5% of the amount supplied (on an as-fed basis). Basal ration was administered in the collective manger of each box. During the entire experiment, the goats had free access to water.

Individual rumen fluid was collected at the two time points (T0 and T1), using an oesophageal polyethylene probe (internal and external diameters of 8 and 12 mm, respectively; length: 1.8 m). Three diverse oesophageal probes connected with a glass vacuum pump were used for the each specific diet. The samples (0.2 l of rumen fluid) were collected 5 h after the morning feeding (13:30 h), stored at  $-20^{\circ}\text{C}$  and used for metagenomic analyses and FA composition characterization.

#### *Diet and milk chemical analysis*

The basal ration and the supplements were stored at  $-20^{\circ}\text{C}$  and analyzed in the next 48 h for the concentrations of ash (method 942.05; AOAC International, 1995), CP (method 984.13; AOAC International, 1995), ether extract (EE; method 920.29; AOAC International, 1995), and NDF corrected for insoluble ash and with the addition of  $\alpha$ -amylase (aNDFom; Mertens, 2002), using the Ankom 200 fiber apparatus (Ankom Technology Corp., Fairport, NY, USA).

Diet FA were analyzed according to Demirel *et al.* (2004). In brief, FA in the feed were obtained by hydrolysis with 5 M KOH in 1:1 (vol/vol) methanol:water at  $60^{\circ}\text{C}$  for 3 h. The samples were acidified to pH 1.0 and heated for an additional 1 h before extraction of the FA into light petroleum (boiling point:  $40^{\circ}\text{C}$  to  $60^{\circ}\text{C}$ ). The fatty acid methyl esters (FAMES) in hexane were injected into a GC Agilent 6890 GC system (Agilent Technologies, Santa Clara, CA, USA) equipped with autosampler, on-column injector, and a FID

detector. The separation was performed on a 100% dimethylpolysiloxane column (CP-Sil88 for FAME,  $100\text{ m} \times 0.25\text{ mm} \times 25\text{ }\mu\text{m}$ , Agilent Technologies, Santa Clara, CA, USA) adopting the conditions of Revello Chion *et al.* (2010), with some modification: the carrier gas was hydrogen at a constant pressure of 28 psi. Column temperature was set at  $60^{\circ}\text{C}$  for 6 minutes, and, then, increased to  $165^{\circ}\text{C}$  at a rate of  $15^{\circ}\text{C}/\text{min}$  and held for 1 min, then increased to  $225^{\circ}\text{C}$  at a rate of  $2^{\circ}\text{C}/\text{min}$  and held for 17 min. Fatty acid methyl esters were identified by comparison of their retention times with known standards (37-component FAME mix, Supelco 47885-U) and expressed as g/100 FA.

Milk samples were analyzed for fat using Gerber method (method 2446; International Organization for Standardization, 1976); CP and Lactose concentration were determined using a Fourier transform IR analyzer (MilkoScan FT6000; Foss Analytical A/S, Hillerød, Denmark). Milk urea concentration was determined by using a differential pH technique (method 14637; International Organization for Standardization, 2006).

#### *Deoxyribonucleic acid extraction, library preparation and sequencing*

Deoxyribonucleic acid extraction was performed from 0.25 g of lyophilized rumen fluid samples following the protocol described in literature (Yu and Morrison, 2004), and stored at  $-20^{\circ}\text{C}$  until use. DNA concentration and purity were measured using a NanoDrop ND-1000 spectrophotometer (Nano Drop Technologies, Inc., Wilmington, DE, USA).

For metagenomic analysis, bacterial and archaeal DNAs were amplified using the primers described in literature (Caporaso *et al.*, 2011), which target the V3 to V4 hypervariable regions of the 16S rRNA gene. All PCR amplifications were performed in 25- $\mu\text{l}$  volumes/sample; 12.5  $\mu\text{l}$  of Thermo Master Mix 2 $\times$  (Life Technologies, Italia, Monza, MB, Italy) and 0.2  $\mu\text{l}$  of each primer (100  $\mu\text{M}$ ) were added to 2  $\mu\text{l}$  of genomic DNA (5 ng/ $\mu\text{l}$ ). A first amplification step was performed in an Applied Biosystem 2700 thermal cycler (Applied Biosystems, Foster City, CA, USA), as follows: denaturation at  $98^{\circ}\text{C}$  for 30 s; 25 cycles with a denaturing step at  $98^{\circ}\text{C}$  for 30 s, annealing at  $56^{\circ}\text{C}$  for 1 min and an extension at  $72^{\circ}\text{C}$  for 1 min; final extension at  $72^{\circ}\text{C}$  for 7 min. Amplicons were cleaned-up with Agencourt<sup>®</sup> AMPure<sup>®</sup> XP (Beckman, Coulter Brea, CA, USA), following Illumina 16S Metagenomic Sequencing Library Preparation protocol ([http://supportres.illumina.com/documents/documentation/chemistry\\_documentation/16s/16s-metagenomic-library-prep-guide-15044223-b.pdf](http://supportres.illumina.com/documents/documentation/chemistry_documentation/16s/16s-metagenomic-library-prep-guide-15044223-b.pdf); Illumina) and checked for amplicon size with 2100 Bioanalyzer Instruments (Agilent Technologies, Santa Clara, CA, USA). Libraries were prepared by a second PCR amplification step using Nextera XT index 1 Primers (N7XX) and Nextera XT Index 2 Primers (S5XX) from the Nextera XT Index Kit (FC-131-1001 or FC-131-1002), following 16S Metagenomic Sequencing Library Preparation protocol. Libraries were quantified by Real Time PCR with KAPA Library Quantification Kits (KapaBiosystems, Inc., Wilmington MA, USA) pooled in equimolar proportion and sequenced in one Miseq (Illumina) run with 300-base paired-end reads.

### Metagenomic data analysis

The following procedure was applied for both the bacterial and the archaeal fraction of the rumen content. Raw sequencing reads were processed as follows: read pairs were, firstly, merged together by means of PandaSeq assembler ('PAired-eND Assembler for DNA sequences,' Masella *et al.*, 2012) which performs a local assembly between two overlapping pairs, thus creating a single fragment. Non-overlapping sequences, fragments shorter than 250 bases or longer than 900 bases were discarded. Then, sequences were quality-filtered using the 'split\_libraries\_fastq.py' utility of the QIIME suite (Caporaso *et al.*, 2010), which discards from further analyses any sequence having more than 25% low-quality nucleotides (i.e.: having a phred score of 3 or less). Quality-filtered reads were, then, analyzed with the standard QIIME pipeline. A random subset of 50 000 reads for each sample was used. Sequences were grouped into operational taxonomic units (OTU) by clustering together reads at 97% identity or higher and taxonomically classified against the Greengenes bacterial 16S rRNA database (release 13.8, <http://greengenes.lbl.gov>) by using the RDP classifier (Wang *et al.*, 2007) at 50% confidence. Singleton sequences were discarded as possible artifacts. Alpha diversities were calculated according to different microbial diversity metrics (i.e.: chao1, Shannon index, observed species and Faith's phylogenetic distance) and evaluated in order to determine whether the chosen subset was representative of the overall microbial diversity within each sample and if there were any differences in terms of microbial diversity between the groups of animals subjected to different diet supplementation. A non-parametric *t* test with random permutations was used to determine whether there were statistically significant differences in terms of microbial diversity.

Beta-diversity among samples was calculated using the UniFrac metric (Lozupone and Knight, 2005) and principal coordinates analyses (PCoAs) were conducted by evaluating both weighted and unweighted Unifrac distances, in order to highlight eventual clustering among samples depending on the diet supplementation. All other statistical evaluations and analyses were performed in Matlab (Natick, MA, USA).

### Rumen fatty acid composition analysis

Methyl esters of FA, dimethylacetal (DMA) and dimethyl esters (DME) were prepared by direct extraction and using an acid-catalyzed transmethylation procedure, as follows: 5 g of raw rumen fluid sample were added in a tube with 2 ml of 10% acetyl chloride methanolic solution. Samples were incubated at 50°C overnight, then 2 ml of 6% aqueous potassium carbonate solution were added to neutralize pH. Subsequently, methylated FA were extracted with 5 ml of hexane and transferred in a fresh tube. The extraction step with hexane was repeated twice. Samples were dried under nitrogen flow and, then, resuspended with 0.5 ml of hexane. Samples were collected in a vial for GC analysis.

This methodology directly produces FA methyl esters from the intact sample, reducing time and solvent, and avoiding the isomerization of conjugated FA (Christie, 1993).

Furthermore, acidic catalyzed transmethylation is necessary to release free FA, which were abundant in rumen liquor. As the high level of water in the samples may reduce the transesterification efficiency, to assure that reaction was complete, samples were examined by thin-layer chromatography (TLC) using hexane-diethyl ether (70 : 30, v/v) and the single spot representing fatty acid esters was visualized with chromo-sulphuric mix (Supplementary Figure S1).

The acidic transesterification method also produced DMA, which are derived from fatty aldehydes, released from plasmalogens.

Dimethylacetal and DME identification was performed on three samples by TLC separation and GC-MS as described by Alves *et al.* (2013), starting from an aliquot of esterified fraction. Mass spectrometry spectra of the single DMA were reported in the Supplementary Figures S2 and S3. Comparing total FA, FA methyl esters and DMA chromatograms, it is possible to identify the peaks.

The FA composition was determined by GC analysis using a GC2010 Shimadzu gas chromatograph (Shimadzu, Columbia, MD, USA) equipped with a flame-ionization detector and a high polar fused-silica capillary column (Chrompack CP-Sil88 Varian, 152 Middelburg, the Netherlands; 100 m, 0.25 mm i.d.; film thickness 0.20 µm). Hydrogen was used as the carrier gas at a flow of 1 ml/min. Split/splitless injector was used with a split ratio of 1 : 40. An aliquot of the sample was injected under the following GC conditions: oven temperature started at 40°C and was held at that level for 1 min; then, it was increased to 163°C at a rate of 2°C/min, and held at that level for 10 min, before being, once again, increased to 180°C at 1.5°C/min and held for 7 min, and, finally, to 187°C at a rate of 2°C/min; at last the temperature was increased to 220°C with a rate of 3°C/min and held for 25 min. Injector temperature was set at 270°C and detector temperature was set at 300°C. Individual FA methyl esters were identified by comparison with a standard mixture of 52 Component FAME Mix (Nu-Chek-Prep Inc., Elysian, MN, USA). Nonadecanoic acid was added as internal standard. The identification of isomers of 18 : 1 was based on commercial standard mixtures (Supelco, Bellefonte PA, USA) and published isomeric profiles (Kramer *et al.*, 2008). Total ruminal FA, DMA and DME fraction were expressed as mg/g DM, whereas the profile was expressed as g/100 g of the sum of FA DMA and DME, respectively.

### Statistical analysis

All data (i.e. milk yield and composition, FA, DMA and DME level) were processed using the following model, adjusted using the independent variable (i.e. FA composition) of the pre-trial period as covariates:

$$y_{ij} = \mu + D_i + \beta(x_{ij} - \bar{x}) + e_{ij}$$

where  $y_{ij}$  is the independent variable;  $\mu$  the overall mean;  $D_i$  the fixed effect of diet ( $i$  = Control, Hemp, Linseed diets);  $\beta$  the linear regression coefficient between  $y_{ij}$  and  $x_{ij}$ ;  $x_{ij}$  the value of covariate corresponding to  $y_{ij}$ ;  $\bar{x}$  the mean of  $x_{ij}$  and  $e_{ij}$  the residual error.

Correlations between FA, DMA and family-level microbiota profiles were evaluated calculating Pearson's coefficient between the raw relative abundance of each family and the ruminal FA composition (as assessed above); only correlations with a p-value of the linear model below 0.005 were considered as significant.

## Results

### Milk yield and composition

Milk yield resulted unaffected ( $P=0.985$ ) by the different dietary treatment, with an average production of 2.33 kg/d ( $SE=0.12$ ). Milk composition also resulted unchanged irrespective of the different supplementation, except for milk fat concentration. Goats fed the C diet showed a lower fat percentage than that fed with the fat supplemented diets (3.39, 3.73, and 3.69%, for C, L, and H, respectively;  $SE=0.10$ ;  $P=0.013$ ). Crude protein, lactose and milk urea concentrations were, on average, as follows: 3.27%, 4.54% and 28.1 mg/dl. The daily dry matter intake, determined on a group basis, was on average 1.9 kg.

### Metagenomic analyses

Raw sequencing data (bacteria and archaea) have been deposited into Short Reads Archive (<http://www.ncbi.nlm.nih.gov/sra/>) under Accession Number PRJNA316187.

Bacterial groups within the rumen generated an average of  $12\,461 \pm 1231$  OTUs per sample, which could be assigned to specific taxonomies down to the genus level. After singletons removal, we obtained  $38\,259 \pm 2684$  reads per sample. Independently from sampling time and diet, rumen composition was dominated by bacteria belonging to Bacteroidetes (summing up to about 61.2% on average relative abundance), followed by Firmicutes (24.2%), whereas subdominant phyla were Proteobacteria and Verrucomicrobia (4.1% and 3.3%, respectively). At family level (Figure 1a), Prevotellaceae resulted as the dominant group (average relative abundance, rel ab, 41.0%), followed by Veillonellaceae (9.4%), unclassified Bacteroidales (6.6%) and Bacteroidales RF16 family (5.3%), which corresponds to a group of unidentified rumen bacteria in SILVA 16S rRNA database (release 121, <http://www.arb-silva.de/browser/ssu-121/EU794092/>). Other components were within the families of Ruminococcaceae, Paraprevotellaceae and Lachnospiraceae (rel ab of 5.0%, 4.4%, 4.3%, respectively). The rarefaction curves analysis, which assesses species richness from sampling, suggested that our sampling was enough to sufficiently describe the main components of the rumen microbiota biodiversity. Moreover, in relation to the three different diets, a trend toward a reduction of the biodiversity was observed in the goats subjected to L diet (Figure 2, Supplementary Figure S4). No difference, on the other hand, was found on the basis of the inter-sample diversity on the basis of the PCoA evaluations (beta-diversity, data not shown).

The relative abundance of the most abundant bacterial genera of the three diet groups is reported in Table 3. Linseed supplementation induced a significantly reduction of *Prevotella* spp. (−12%), and an increase of *Succinivibrio* spp.

(+50%) and *Fibrobacter* spp. (+75%), compared with the samples of control group. On the contrary, hemp did not show significant differences on the relative abundance of phyla.

For archaeal diversity, the reads were clustered in an average of  $4616 \pm 958$  OTUs at 97% of identity. Operational Taxonomic Unit rarefaction curves were considered as having reached a plateau (approximating the saturation level), at about 35 000 reads (Supplementary Figure S5).

According to our data (Figure 1b), independently from the diets and the time points of rumen collection (T0 or T1), the goat rumen was dominated by the families of Methanobacteriaceae (average rel. ab  $36.8\% \pm 11.3\%$ ), Methanomassiliicoccaceae (average rel. ab.  $37.1\% \pm 10.4\%$ ), whereas about 26% of reads were found to be bacterial sequences of rumen origin by taxonomic classification by RDP classifier, specifically Prevotellaceae, Bacteroidales and Ruminococcaceae (about 8%, 3% and 1%, respectively). This was probably due to a lower representativeness of Archaea sequences within the Greengenes database which could result in non-specific matches for non-represented taxa or to a partial cross-match between the PCR primers used for Archaea amplification and bacterial 16S sequences.


The majority of the sequences were related to the genus vadin CA11 (within the family of Methanomassiliicoccaceae), representing an average of 33.2%, 33.2% and 39.0% of the sequences in the rumen fluid fraction of C, L and H diets, respectively. The genus *Methanobrevibacter* was the second most predominant, representing an average of 39.5%, 31.2% and 26.8% of the reads in C, L and H diets, respectively, followed by *Methanosphaera*. However, these differences in the average relative abundance of archaeal genera was found to be not statistically significant, due to the relatively high variability in the composition between the individual animals constituting each dietary group.

### Ruminal fatty acid composition

Fatty acid and DMA composition of rumen fluid is reported in Tables 4 and 5. Saturated FA represented the most abundant class (>70%), as a consequence of the high rumen BH rate.

Fatty acids amount was significantly higher in L (+30%) and H (+23%) groups than control group as consequence of lipid supplementation in the diet.

Saturated FA significantly increased with fat supplementation, being contained at higher amount in rumen liquor from L (+19%) and H diet (+15%). Diet H induced higher content of PUFA n-6 in rumen liquor, whereas L diet led to a significant increase of PUFA n-3. In particular, linseed supplementation increased  $\alpha$ -linolenic acid (+118%) and some FA that origin from the rumen BH of linolenic acid: C18:3c9t11c15 (+257%), C18:2t11c15 (+79%), C18:1t15 (+1369%) and C18:1t16 (+526%) ( $P < 0.01$ ). Diet supplemented with hemp seed increased the content of linoleic acid (+88%), C18:1t6 + t7 + t8 (+110%), C18:1t9 (+194%) and C18:1t10 (121%). Stearic acid (C18:0) content significantly increased regardless the type of lipid supplement. However, L diet led to the highest increase of stearic acid (+108%). The content of branched chain fatty acids (BCFA)



**Figure 1** Relative abundance of Bacterial (a) and Archaeal (b) compositions of different communities with the three experimental diets (C = control; H = hemp; L = linseed) at T0 and T1 sampling times.

(−20%) and odd chain fatty acid (OCFA) (−26%) decreased irrespective to the source of fat.

Nineteen DMA and two DME were identified in rumen samples (Table 5). In all, 12 individual DMA significantly changed in DMA profile as a consequence of lipid supplementation: DMA12:0, DMA13:0, DMA14:0, DMA15:0*iso*, DMA15:0*ante*, DMA15:0, DMA16:0, DMA17:0*iso*, DMA18:1*c*9, DMA18:1*c*11, DMA18:1*c*12 and DMA 18:2.


*Rumen fatty acid profile and bacterial microbiome association*

The FA and DMA that showed high and significant ( $P < 0.005$ ) Pearson’s correlations with bacterial microbiome composition at the family level are reported in Table 6.

Prevotellaceae showed a positive correlation with PUFA, PUFA n-6, whereas Clostridiales and Ruminococcaceae were negatively correlated with PUFA. Fibrobacteraceae was the only family that showed a significant correlation with PUFA n-3. Significant correlations were also found for C18:0 (Fibrobacteraceae), C18:1*t*10 (Prevotellaceae), C18:1*t*15 (Fibrobacteraceae), C18:1*t*12 (Paraprevotellaceae) and C18:1*t*16 (Fibrobacteraceae).

Dimethylacetal12:0, DMA17:0*iso* and DMA18:1*t*11 were the only DMA positively correlated with Prevotellaceae, (DMA12:0 and DMA18:1*t*11) and Veillonellaceae (DMA17:0*iso*).

Bacteroidales RF16 is the only family positively correlated with BCFA (mainly with C16:0*iso*) and OCFA.



**Figure 2** Bacterial rarefaction curves calculated according to chao1 (a) and Faith's phylogenetic metric (b) for all samples and represented for each different diet (C = control; H = hemp; L = linseed) at the two time points (T0 and T1). Curves were drawn using the least sequenced sample (~28 000 reads) as upper limit for the rarefactions.

**Table 3** Relative abundances of the 10 most abundant bacterial phyla revealed in the rumen with control (C), linseed (L) or hemp (H) diets

Headings	C	L	H	SEM	P-value
<i>Prevotella</i> spp.	0.405 <sup>a</sup>	0.355 <sup>b</sup>	0.452 <sup>a</sup>	0.023	0.003
<i>Succinicladium</i> spp.	0.059	0.052	0.053	0.011	0.483
<i>Ruminococcus</i> spp.	0.020	0.022	0.019	0.003	0.548
<i>Oscillospira</i> spp.	0.003	0.003	0.002	0.001	0.829
<i>Clostridium</i> spp.	0.004	0.005	0.003	0.001	0.738
<i>Butyrivibrio</i> spp.	0.008	0.007	0.006	0.001	0.333
<i>Coprococcus</i> spp.	0.007	0.007	0.006	0.001	0.651
<i>Pseudobutyrvibrio</i> spp.	0.003	0.002	0.002	0.001	0.932
<i>Selonomonas</i> spp.	0.005	0.005	0.004	0.001	0.665
<i>Succinivibrio</i> spp.	0.003 <sup>b</sup>	0.006 <sup>a</sup>	0.003 <sup>b</sup>	0.001	0.034
<i>Treponema</i> spp.	0.008	0.010	0.012	0.002	0.212
<i>Fibrobacter</i> spp.	0.006 <sup>b</sup>	0.014 <sup>a</sup>	0.009 <sup>b</sup>	0.002	0.021

SEM = standard error of the mean.

<sup>a,b</sup> Values within a row with different superscripts differ significantly at  $P < 0.05$ .

## Discussion

In this study, NGS and GC techniques were used to assess the effects of dietary fat supplementation on rumen microbial community and lipid composition of rumen liquor, respectively, in 18 Alpine goats. For the experimental plan, a period of 100 days was considered to verify a change in the composition of rumen microbial community.

Irrespective to the dietary treatment groups, Bacteroidetes and Firmicutes dominated the composition of phyla, with a high abundance of Prevotellaceae, Veillonellaceae and a low presence of Ruminococcaceae, Paraprevotellaceae and Lachnospiraceae. This result is consistent with a previous research published by Cunha *et al.* (2011), who characterized bacterial community in Moxotó goats by NGS technique. Moreover, similar results were described also by Min *et al.* (2014) where the percentages of Bacteroidetes and Firmicutes

in rumen sheep's samples were 53% and 36.7% of the total bacterial sequences respectively, and by Zened *et al.* (2013), for whom, more than 80% of total sequences in rumen samples of cow fed different diets, were represented by Firmicutes (49.6%) and Bacteroidetes (40.2%). In our study, Prevotellaceae was the dominant bacterial family in all the experimental diets at the both sampling times. Therefore, Prevotellaceae and Bacteroidaceae, were, apparently, the principal families involved in the degradation of the diet at the rumen level, though more investigation are needed to support these conclusions. As far as Archaea, which account for about 4% of ruminal microbes (Paul *et al.*, 2015), *Methanobrevibacter* was the main genus, confirming what previously reported in studies on rumen liquor from cattle (Sirohi *et al.*, 2013) and goat (Cunha *et al.*, 2011). *Methanobrevibacter* is a genus of Methanobacteriaceae, obligate anaerobes distinctly differentiated from other organisms, that produces methane as a major catabolic product. The variation in community composition may be due to the rumen environment and the diet (Buccioni *et al.*, 2012).

Rumen liquor contains several FA and DMA that are not present in the feedstuffs. Some of these compounds are the result of microbial BH, whereas DMA are known to be structural lipids of microbial membranes and cell walls, and are *de novo* synthesized by rumen microbes, participating in the maintenance of optimal membrane fluidity (Alves *et al.*, 2013). So, FA composition of rumen liquor may provide useful information about the effect of lipid supplementation on rumen bacterial community and its metabolism.

Several studies have investigated the effect of lipid supplementation on the rumen FA in ruminant species, with particular attention for BH pathway (Loor *et al.*, 2005; Toral *et al.*, 2012). The present study showed that two different lipid supplementations, hemp (linoleic acid source) or linseed ( $\alpha$ -linolenic acid source), promoted changes in rumen BH pathway in goat. The first effect observed was the dramatic

**Table 4** Effect of control (C), linseed (L) and hemp (H) diets in the total fatty acid (FA) (mg/g of dry matter) and FA composition (% of total FA) of goat rumen content

FAs	C	L	H	SEM	P-value
Total	23.502 <sup>B</sup>	30.528 <sup>A</sup>	28.880 <sup>A</sup>	1.503	0.023
SFA					
C12:0	0.741 <sup>A</sup>	0.296 <sup>B</sup>	0.374 <sup>B</sup>	0.033	0.003
C13:0iso	0.216	0.182	0.197	0.015	0.917
C13:0	0.269	0.232	0.202	0.021	0.779
C14:0iso + n.i. component	3.110 <sup>a</sup>	1.794 <sup>c</sup>	2.435 <sup>b</sup>	0.145	0.038
C14:0 + n.i. component	5.087 <sup>A</sup>	2.850 <sup>B</sup>	3.189 <sup>B</sup>	0.165	0.007
C15:0iso	1.968	1.728	1.581	0.066	0.568
C15:0ante	2.376	1.644	1.533	0.121	0.189
C15:0	1.495 <sup>A</sup>	1.002 <sup>B</sup>	0.931 <sup>B</sup>	0.036	0.006
C16:0iso	0.931 <sup>a</sup>	0.521 <sup>b</sup>	0.582 <sup>b</sup>	0.064	0.023
C16:0	26.272 <sup>A</sup>	19.035 <sup>b</sup>	21.858 <sup>b</sup>	0.767	0.045
C17:0iso	0.546 <sup>a</sup>	0.390 <sup>b</sup>	0.353 <sup>b</sup>	0.027	0.019
C17:0ante	1.334 <sup>A</sup>	0.600 <sup>B</sup>	0.717 <sup>B</sup>	0.033	<0.001
C17:0	0.976	0.724	0.725	0.052	0.588
C18:0iso	0.495 <sup>a</sup>	0.387 <sup>ab</sup>	0.316 <sup>b</sup>	0.017	0.039
C18:0	24.576 <sup>C</sup>	43.431 <sup>A</sup>	35.244 <sup>B</sup>	2.044	0.002
C20:0	0.323 <sup>B</sup>	0.279 <sup>B</sup>	0.439 <sup>A</sup>	0.017	0.004
C21:0	4.462	3.290	3.313	0.201	0.457
C23:0	1.382 <sup>A</sup>	0.344 <sup>B</sup>	0.345 <sup>B</sup>	0.071	<0.001
MUFA					
C18:1 $\omega$ -8	0.363 <sup>c</sup>	0.527 <sup>b</sup>	0.654 <sup>a</sup>	0.051	0.032
C18:1 $\omega$ 9	0.119 <sup>C</sup>	0.141 <sup>B</sup>	0.200 <sup>A</sup>	0.013	<0.001
C18:1 $\omega$ 10	0.300 <sup>C</sup>	0.305 <sup>B</sup>	0.571 <sup>A</sup>	0.018	<0.001
C18:1 $\omega$ 11	3.629 <sup>a</sup>	2.257 <sup>b</sup>	2.594 <sup>b</sup>	0.152	0.013
C18:1 $\omega$ 12	0.741 <sup>C</sup>	0.995 <sup>B</sup>	1.151 <sup>A</sup>	0.044	0.002
C18:1 $\omega$ 9 + $\omega$ 13 + $\omega$ 14	7.986	5.982	8.196	0.971	0.625
C18:1 $\omega$ 15	0.160 <sup>C</sup>	1.997 <sup>A</sup>	0.683 <sup>B</sup>	0.222	0.016
C18:1 $\omega$ 11	0.711	0.760	0.699	0.101	0.804
C18:1 $\omega$ 12	1.357	1.063	1.388	0.067	0.191
C18:1 $\omega$ 13	0.694	0.600	0.451	0.069	0.581
C18:1 $\omega$ 14	0.234	0.211	0.160	0.034	0.453
C18:1 $\omega$ 16	0.286 <sup>C</sup>	1.525 <sup>A</sup>	0.692 <sup>B</sup>	0.079	<0.001
C18:1 $\omega$ 15	0.564 <sup>a</sup>	0.463 <sup>a</sup>	0.253 <sup>b</sup>	0.041	0.003
PUFA					
C18:2 $n$ -6	3.298 <sup>B</sup>	1.380 <sup>B</sup>	5.401 <sup>A</sup>	0.367	0.003
C18:2 $n$ -7	0.500 <sup>B</sup>	0.760 <sup>A</sup>	0.265 <sup>B</sup>	0.044	0.002
C18:2 $c$ 9 $\omega$ 11	1.982 <sup>A</sup>	0.984 <sup>B</sup>	1.570 <sup>A</sup>	0.034	<0.001
C18:3 $n$ -3	0.365 <sup>B</sup>	0.680 <sup>A</sup>	0.396 <sup>B</sup>	0.019	<0.001
C18:3 $c$ 9 $\omega$ 11 $\omega$ 15	0.188 <sup>B</sup>	0.571 <sup>A</sup>	0.199 <sup>B</sup>	0.027	<0.001
Class of FA					
SFA	76.522 <sup>C</sup>	78.730 <sup>A</sup>	74.332 <sup>B</sup>	1.347	0.002
MUFA	17.145	16.685	17.835	0.821	0.412
PUFA	6.334 <sup>B</sup>	4.375 <sup>C</sup>	7.832 <sup>A</sup>	0.365	0.008
PUFAn-6	3.298 <sup>B</sup>	1.380 <sup>C</sup>	5.401 <sup>A</sup>	0.365	0.002
PUFAn-3	1.054 <sup>B</sup>	2.010 <sup>A</sup>	0.861 <sup>B</sup>	0.257	0.003
BCFA	10.976 <sup>A</sup>	7.247 <sup>B</sup>	7.715 <sup>B</sup>	0.288	0.029
BCFAiso	7.266 <sup>a</sup>	5.002 <sup>b</sup>	5.465 <sup>b</sup>	0.186	0.027
BCFAante	3.710 <sup>a</sup>	2.245 <sup>b</sup>	2.250 <sup>b</sup>	0.129	0.017
OCFA	14.988 <sup>A</sup>	10.137 <sup>B</sup>	9.896 <sup>B</sup>	0.215	0.003

SEM = standard error of the mean; n.i. = not identified; SFA = saturated fatty acids; MUFA = monounsaturated fatty acids; PUFA = polyunsaturated fatty acids; BCFA = branched chain fatty acids; OCFA = odd chain fatty acids. Different letters in the same row correspond to statistically different values: a,b,c:  $P < 0.05$ , A,B,C:  $P < 0.001$ .

increase of C18:0 in rumen fluid of animal fed with lipid supplemented diet, as a consequence of the extensive BH occurring in goat rumen, as proposed by Toral *et al.* (2016). During the BH process, unsaturated FA are isomerized and

**Table 5** Effect of control (C), linseed (L) and hemp (H) diets in the total fatty acid (FA) dimethylacetals (DMA) and dimethylesters (DME) (mg/g of dry matter) and DMA and DME composition (% of total DMA + DME) of goat rumen content

Dimethylacetals	C	L	H	SEM	P-value
Total	2.898	2.422	3.014	0.284	0.433
DMA12:0	1.468 <sup>b</sup>	2.004 <sup>a</sup>	1.506 <sup>b</sup>	0.142	0.045
DMA13:0iso	0.809	0.831	0.934	0.101	0.382
DMA13:0	3.224 <sup>a</sup>	3.583 <sup>a</sup>	2.210 <sup>b</sup>	0.371	0.021
DMA14:0iso	0.962	1.178	0.958	0.153	0.184
DMA14:0	7.206 <sup>A</sup>	6.193 <sup>B</sup>	7.024 <sup>A</sup>	0.522	0.007
DMA15:0iso	1.077 <sup>a</sup>	1.002 <sup>b</sup>	0.938 <sup>b</sup>	0.103	0.032
DMA15:0ante	5.376 <sup>b</sup>	8.304 <sup>a</sup>	7.179 <sup>a</sup>	0.686	0.989
DMA15:0	4.966 <sup>A</sup>	3.944 <sup>B</sup>	4.261 <sup>B</sup>	0.341	0.008
DMA16:0iso	2.341	2.069	2.499	0.337	0.077
DMA16:0	29.839 <sup>a</sup>	20.516 <sup>c</sup>	23.273 <sup>b</sup>	2.071	0.022
DMA16:1 $\omega$ 9	0.783	1.044	0.577	0.202	0.466
DMA17:0iso	1.378 <sup>a</sup>	0.674 <sup>b</sup>	0.745 <sup>b</sup>	0.206	0.037
DMA17:0ante	14.187	16.837	17.032	1.937	0.386
DMA18:0	0.694	0.761	0.543	0.085	0.127
DMA18:1 $\omega$ 11	0.895	1.336	0.801	0.268	0.467
DMA18:1 $\omega$ 9	2.555 <sup>a</sup>	2.647 <sup>a</sup>	1.918 <sup>b</sup>	0.306	0.012
DMA18:1 $\omega$ 11	2.061 <sup>A</sup>	1.427 <sup>B</sup>	1.511 <sup>B</sup>	0.147	<0.001
DMA18:1 $\omega$ 12	0.777 <sup>b</sup>	0.859 <sup>b</sup>	1.068 <sup>a</sup>	0.084	0.024
DME9:0	1.861	2.791	1.704	0.227	0.208
DMA17:1	14.104	16.431	17.631	1.331	0.503
DMA18:2	1.746 <sup>B</sup>	3.968 <sup>A</sup>	3.891 <sup>A</sup>	0.368	0.003
DME10:0	1.692	1.601	1.509	0.285	0.351
DMA_SFA	73.526 <sup>A</sup>	67.869 <sup>B</sup>	69.390 <sup>B</sup>	1.327	0.002
DMA_MUFA	21.175	23.744	23.506	1.374	0.681
DMA_PUFA	1.746 <sup>B</sup>	3.968 <sup>A</sup>	3.891 <sup>A</sup>	0.368	0.003
DMA_BCFA	26.130	30.895	30.573	2.341	0.429
DMA_OCFA	31.017 <sup>b</sup>	35.175 <sup>a</sup>	33.587 <sup>ab</sup>	2.629	0.023
DME	3.553 <sup>b</sup>	4.392 <sup>a</sup>	3.213 <sup>b</sup>	0.378	0.041

SEM = standard error of the mean; n.i. = not identified; SFA = saturated fatty acids; MUFA = monounsaturated fatty acids; PUFA = polyunsaturated fatty acids; BCFA = branched chain fatty acids; OCFA = odd chain fatty acids. Different letters in the same row correspond to statistically different values: a,b,c:  $P < 0.05$ , A,B:  $P < 0.001$ .

hydrogenated generating a complex system of isomeric unsaturated FA, mostly with *trans* double bonds. These intermediates of rumen BH may largely vary according to changes in diet composition, as reported by Chilliard *et al.* (2007). Linseed favors the accumulation of characteristic intermediate of C18:3 $n$ -3 BH (C18:3 $c$ 9 $\omega$ 11 $\omega$ 15, C18:2 $\omega$ 11 $\omega$ 15, C18:1 $\omega$ 15, C18:1 $\omega$ 15). On the other hand, hemp promotes an increase of C18:2 $n$ -6 BH intermediate (C18:1 $\omega$ -8, C18:1 $\omega$ -9, C18:1 $\omega$ -10, C18:1 $\omega$ -12).

Significantly, the higher level of C18:0 in the lipid supplemented diet is not related to a contemporary increase of vaccenic acid (VA, C18:1 $\omega$ -7), the principal BH intermediate. Probably, lipid supplementation shifts the BH pathway synthesizing other *trans* C18:1 isomers instead of VA. In fact, a relevant increase of all *trans* FA (~+50%) in H and L groups, except VA, which showed similar level in comparison to C group, was observed. Moreover, VA represents 67% of total *trans* FA in C group, whereas this percentage is lower than 50% in H and L ones. The change of  $\alpha$ -linolenic and linoleic acid ratio in the diet affect BH pathway, as well described by Chilliard and Ferlay (2004). This effect was



**Table 6** Rumen fatty acid (FA) and dimethylacetals (DMA) showing a significant ( $P < 0.005$ ) Pearson's correlation with bacterial microbiome composition at the family level

Family	FA and DMA															
	DMA 12:0	C12:0	C16 <sub>iso</sub>	DMA 17:0 <sub>iso</sub>	DMA 18:1 $\tau$ 11	C18:0	C18:1 $\tau$ 10	C18:1 $\tau$ 15	C18:1 $\tau$ 12	C18:1 $\tau$ 16	PUFA	n-6	n-3	BCFA	BCFA <sub>ante</sub>	OCFA
Prevotellaceae	–	0.58	–	–	0.52	–	0.51	–	–	–	0.56	0.63	–	–	–	–
Paraprevotellaceae	–	–	–	–	–	–	–	–	0.52	–	–	–	–	–	–	–
Fibrobacteraceae	–	–	–	–	–	0.55	–	0.58	–	0.50	–	–	0.60	–	–	–
Clostridiales	–	–	–	–	–	–	–	–	–	–	–0.54	–	–	–	–	–
Ruminococcaceae	–	–	–	–	–	–	–	–	–	–	–0.50	–	–	–	–	–
Veillonellaceae	–	–	–	0.52	–	–	–	–	–	–	–	–	–	–	–	–
Succinivibrionaceae	0.55	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Bacteroidales RF16	–	–	0.59	–	–	–	–	–	–	–	–	–	–	0.56	0.53	0.51

revealed in cows fed with linseed and sunflower supplemented diet, two sources of  $\alpha$ -linolenic and linoleic acid, respectively (Loor *et al.*, 2005).

In the present study, Fibrobacteriaceae is the bacterial family showing the highest and significant correlation with FA involved in the BH pathway of C18:3n-3. This correlation may be related to the production of C18:0, which is one of the most abundant SFA characterizing *F. succinogenes*, the principal bacterium of Fibrobacteriaceae (Shah and Collins, 1983), that, probably, synthesized C18:0 for requirements, starting from  $\alpha$ -linolenic acid. Maybe, *F. succinogenes* play a role in the detoxification of ruminal environment by the production of C18:0, necessary for its metabolism, starting from C18:3n-3. However, this hypothesis need more investigation.

Apart from the C18:1 $\tau$ 10 and C18:1 $\tau$ 11, several others C18:1 isomers were produced in the rumen which can be accumulated in milk and meat tissues. In fact, the BH pathways of C18:2n-6 and C18:3n-3 are characterized by 15 octadecenoic FA isomers, ranging from *trans*-6 to *cis*-16 (Chilliard *et al.*, 2007).

As proved up by this study, some intermediate compounds of rumen BH may be considered a marker of the most representative pathway. Particularly, C18:3 $\tau$ 9, $\tau$ 11, $\tau$ 15 and C18:2 $\tau$ 11,  $\tau$ 15 are well-known BH intermediates of the C18:3n-3 (Harfoot and Hazlewood, 1997); high level of these FA may be a good marker of a diet rich in C18:3n-3, like pasture or linseed supplementation.

Moreover, the significant and positive correlation between the relative abundance of the unidentified rumen bacteria RF16 and BCFA and OCFA, may indicate that this family plays a role in the rumen metabolism. BCFA and OCFA are synthesized in the rumen by cellulolytic and amilolytic bacteria (Buccioni *et al.*, 2012). RF16 is an unclassified family of the Bacteroidales (Jewell *et al.*, 2015), so an investigation is needed to understand its role on BCFA and OCFA synthesis.

Another important class of ruminal lipids is represented by plasmalogens, which are strictly related to bacteria metabolism. Plasmalogen lipids are present in large amounts in anaerobic bacteria membranes with different physical properties compared with other phospholipids (Alves *et al.*, 2013). Some of DMA are structural lipids of microbial

membranes and cell walls, regulating the optimal fluidity of membrane (Goldfine, 2010). Consequently, rumen microbes might have modified their plasmalogen content and composition in response to different environmental stimuli and microbial diversity. As reported by our results and confirmed by previous work (Alves *et al.*, 2013), rumen DMA composition was similar to the FA composition, presenting odd- and branched-chains with carbon lengths between C15 and C18.

From this study a significant reduction of BCFA<sub>iso</sub>, probably due to the relative decrease of cellulolytic bacteria, was revealed. Therefore, it is tempting to speculate that the reduction of DMA can be related with this group of bacteria. As BCFA have been described as a diagnostic tool of rumen fermentation pathway (Vlaeminck *et al.*, 2006), thus the DMA composition derived from plasmalogens may have the same role, as also proposed by Alves *et al.* (2013). Previous studies proposed that the vinyl ether origin by conversion of a FA via an alcohol intermediate, following a different way than the aerobic pattern in which the vinyl ether bond is formed at an early stage of the plasmalogen biosynthesis (Goldfine, 2010).

Ruminants harbor diverse and dense microbial populations in the rumen, which is essential for the bioconversion of feeds that are otherwise indigestible for the host digestive system. Microbial BH process may accumulate a wide range of intermediates (Palmquist *et al.*, 2005), including rumenic acid, which is reduced to VA and finally to 18:0.

The main bacterial species identified involved in the BH process belong to the *Butyrivibrio* group, which includes the genera *Butyrivibrio* and *Pseudobutyrvibrio* (Paillard *et al.*, 2007) and phylogenetically related bacteria (Devillard *et al.*, 2007). Both these genera were not affected by hemp and linseed supplementation. Despite this result, a different BH pathway between hemp and linseed group was observed. These findings confirm the recent *in vivo* studies (Belenguer *et al.*, 2010; Huws *et al.*, 2011; Shingfield *et al.*, 2012; Castro Carrera *et al.*, 2014), which have suggested a lower relevance of *Butyrivibrio* and *Pseudobutyrvibrio* species in ruminal BH than initially thought. Plasmalogens are found in many strictly anaerobic bacteria including most species of *Clostridium*, *Lachnospira*, and *Butyrivibrio* (Goldfine, 2010), but are not present in aerobic or facultative anaerobic bacteria.

In the present work, linseed supplementation reduced significantly the level of *Prevotella* spp. and DMA, so we speculate a possible role of this phylum on plasmalogens synthesis. To our knowledge, this is the first time that the effect of lipid supplementation on DMA synthesis has been reported in goat, so more investigations are needed to understand the role of *Prevotella* on this pathway of rumen lipid metabolism.

Data of Tables 4 and 5 showed that DMA and DME represent about the 15% of total FA + DMA + DME. This result was considerably higher than data reported by Alves *et al.* (2013) (0.86%). Similar result was observed by Saluzzi *et al.* (1995), which reported that DMA represent about 15% of total FA + DMA in microbial fraction isolated from the rumen. This large difference may be related to the type of rumen samples. In fact, it is important to consider that in whole rumen content DM, the microbial DMA concentration would be more diluted by the higher amounts of indigested feed than in liquid phase DM. Data reported by Alves *et al.* (2013) refer to the whole rumen content, whereas our results were obtained from very aqueous rumen fluid, collected by aesophageal probe. Unfortunately, few works on DMA exist in literature and not even one about goat, so more investigations are needed to understand this variability in rumen.

In conclusion, supplementation of feed with different lipid source had relevant effect on biodiversity of ruminal bacterial community in goat. These changes had relevant repercussion on lipid metabolism of rumen, especially on BH and plasmalogens synthesis. The effect of diet on ruminal population and its metabolism is very important to define milk composition by the modulation of FA with particular interest for milk quality.

## Acknowledgments

The authors are thankful to Dr Christian Mallo (IBBA-CNR) for his assistance in animal care. This research was supported by the Agreement Regione Lombardia – National Research Council (CNR), Project FilAgro 'Strategie Innovative e sostenibili per la filiera agroalimentare.'

## Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1751731117003433>

## References

Alves SP, Santos-Silva J, Cabrita ARJ, Fonseca AMJ and Bessa RJB 2013. Detailed dimethylacetal and fatty acid composition of rumen content from lambs fed lucerne or concentrate supplemented with soybean oil. *PLOS one* 3, e58386.

Association of Official Analytical Chemists International 1995. *Official Methods of Analysis*, 15th ed. AOAC International, Washington, DC, USA.

Belenguer A, Toral PG, Frutos P and Hervás G 2010. Changes in the rumen bacterial community in response to sunflower oil and fish oil supplements in the diet of dairy sheep. *Journal of Dairy Science* 93, 3275–3286.

Buccioni A, Decandia M, Minieri S, Molle G and Cabiddu A 2012. Lipid metabolism in the rumen: New insights on lipolysis and biohydrogenation with an emphasis on the role of endogenous plant factors. *Animal Feed Science and Technology* 174, 1–25.

Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J and Knight R 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* 7, 335–336.

Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D and Lozupone CA 2011. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proceedings of National Academy of Science* 108, 4516–4522.

Castro-Carrera T, Toral PG, Frutos P, Mcewan NR, Hervás G, Abecía L, Pinloche E, Girdwood SE and Belenguer A 2014. Rumen bacterial community evaluated by 454 pyrosequencing and terminal restriction fragment length polymorphism analyses in dairy sheep fed marine algae. *Journal of Dairy Science* 97, 1661–1669.

Chilliard Y, Ferlay A, Rouel J and Lamberet G 2003. A review of nutritional and physiological factors affecting goat milk lipid synthesis and lipolysis. *Journal of Dairy Science* 86, 1751–1770.

Chilliard Y and Ferlay A 2004. Dietary lipids and forages interactions on cow and goat milk fatty acid composition and sensory properties. *Reproduction Nutrition Development* 44, 467–492.

Chilliard Y, Glasser F, Ferlay A, Bernard L, Rouel J and Doreau M 2007. Diet, rumen biohydrogenation and nutritional quality of cow and goat milk fat. *European Journal of Lipid Science and Technology* 109, 828–855.

Christie WW 1993. *Advances in lipid methodology*. Oily Press, Dundee, UK. pp. 69–111.

Cunha IS, Barreto CC, Costa OY, Bomfim MA, Castro AP, Kruger RH and Quirino BF 2011. Bacteria and Archaea community structure in the rumen microbiome of goats (*Capra hircus*) from the semiarid region of Brazil. *Anaerobe* 17, 118–124.

Demirel G, Wachira AM, Sinclair LA, Wilkinson RG, Wood JD and Enser M 2004. Effects of dietary n-3 polyunsaturated fatty acids, breed and dietary vitamin E on the fatty acids of lamb muscle, liver and adipose tissue. *British Journal of Nutrition* 91, 551–565.

Devillard E, McIntosh FM, Duncan SH and Wallace RJ 2007. Metabolism of linoleic acid by human gut bacteria: different routes for biosynthesis of conjugated linoleic acid. *Journal of Bacteriology* 189, 2566–2570.

Fernando SC, Purvis HT, Najjar FZ, Sukharnikov LO, Krehbiel CR, Nagaraja TG, Roe BA and DeSilva U 2010. Rumen microbial population dynamics during adaptation to a high-grain diet. *Applied and Environmental Microbiology* 76, 7482–7490.

Fievez V, Colman E, Castro-Montoya JM, Stefanov I and Vlaeminck B 2012. Milk odd- and branched-chain fatty acids as biomarkers of rumen function – an update. *Animal Feed Science and Technology* 172, 51–65.

Frey JC, Pell AN, Berthiaume R, Lapierre H, Lee S, Ha JK, Mendell JE and Angert ER 2010. Comparative studies of microbial populations in the rumen, duodenum, ileum and faeces of lactating dairy cows. *Journal of Applied Microbiology* 108, 1982–1993.

Goldfine H 2010. The appearance, disappearance and reappearance of plasmalogens in evolution. *Progress in Lipid Research* 49, 493–498.

Harfoot CG and Hazlewood GP 1997. Lipid metabolism in the rumen. In *The rumen microbial ecosystem* (ed. PN Hobson), pp. 382–426. Elsevier, London.

Huws SA, Kim EJ, Lee MRF, Scott MB, Tweed JKS, Pinloche E, Wallace RJ and Scollan ND 2011. As yet uncultured bacteria phylogenetically classified as *Prevotella*, *Lachnospiraceae* incertae sedis and unclassified *Bacteroidales*, *Clostridiales* and *Ruminococcaceae* may play a predominant role in ruminal biohydrogenation. *Environmental Microbiology* 13, 1500–1512.

International Organization for Standardization 1976. *Milk: determination of fat content (Gerber method)*. ISO, Geneva, Switzerland.

International Organization for Standardization 2006. *Milk: Determination of urea content. Enzymatic method using difference in pH (Reference method)*. ISO, Geneva, Switzerland.

Jewell KA, McCormick CA, Odt CL, Weimer PJ and Suen G 2015. Rumen bacterial community composition in dairy cows is dynamic over the course of two lactation and correlates with feed efficiency. *Applied and Environmental Microbiology* 81, 4697–4710.

Kramer JKG, Hernandez M, Cruz-Hernandez C, Kraft J and Dugan MER 2008. Combining results of two GC separations partly achieves determination of all cis and trans 16:1, 18:1, 18:2 and 18:3 except CLA isomers of milk fat as demonstrated using Ag-Ion SPE Fractionation. *Lipids* 43, 259–273.

- Loor JJ, Ferlay A, Ollier A, Ueda K, Doreau M and Chilliard Y 2005. High-concentrate diets and polyunsaturated oils alter trans and conjugated isomers in bovine rumen, blood, and milk. *Journal of Dairy Science* 88, 3986–3999.
- Lozupone C and Knight R 2005. UniFrac: a new phylogenetic method for comparing microbial communities. *Applied and Environmental Microbiology* 71, 8228–8235.
- Masella AP, Bartram AK, Truszkowski JM, Brown DG and Neufeld JD 2012. PANDAseq: paired-end assembler for illumina sequences. *BMC Bioinformatics* 13, 31.
- Mele M 2009. Designing milk fat to improve healthfulness and functional properties of dairy products: from feeding strategies to a genetic approach. *Italian Journal of Animal Science* 8, 365–373.
- Mertens DR 2002. Gravimetric determination of amylase-treated neutral detergent fiber in feeds using refluxing in beakers or crucibles: collaborative study. *Journal of AOAC International* 85, 1217–1240.
- Min BR, Wright C, Ho P, Eun JS, Gurung N and Shange R 2014. The effect of phytochemical tannins-containing diet on rumen fermentation characteristics and microbial diversity dynamics in goats using 16S rDNA amplicon pyrosequencing. *Agricultural Food and Analytical Bacteriology* 4, 3–11.
- Paillard D, McKain N, Chaudhary LC, Walker ND, Pizette F, Koppova I, McEwan NR, Kopečný J, Vercoe PE, Louis P and Wallace RJ 2007. Relation between phylogenetic position, lipid metabolism and butyrate production by different *Butyrivibrio*-like bacteria from the rumen. *Antonie van Leeuwenhoek* 91, 417–422.
- Palmquist DL, Lock AL, Shingfield KJ and Bauman DE 2005. Biosynthesis of conjugated linoleic acid in ruminants and humans. *Advances in Food and Nutrition Research* 50, 179–217.
- Paul SS, Deb SM, Dey A, Somvanshi SP, Singh D, Rathore R and Stiverson J 2015. 16S rDNA analysis of archaea indicates dominance of *Methanobacterium* and high abundance of *Methanomassiliicoccaceae* in rumen of Nili-Ravi buffalo. *Anaerobe* 35, 3–10.
- Revello Chion A, Tabacco E, Giaccone D, Peiretti PG, Battelli G and Borreani G 2010. Variation of fatty acid and terpene profiles in mountain milk and 'Toma piemontese' cheese as affected by diet composition in different seasons. *Food Chemistry* 121, 393–399.
- Saluzzi L, Stewart CS, Flint HJ and Smith A 1995. Plasmalogens of microbial communities associated with barley straw and clover in the rumen. *FEMS Microbiology Ecology* 17, 47–56.
- Shah HN and Collins MD 1983. *Megamonas hypermegas* gen. nov., comb. nov. In Validation of the publication of new names and new combinations previously effectively published outside the IJSB, list no. 10. *International Journal of Systemic and Bacteriology* 33, 438–440.
- Shingfield KJ, Bernard L, Leroux C and Chilliard Y 2010. Role of trans fatty acids in the nutritional regulation of mammary lipogenesis in ruminants. *Animal* 7, 1140–1166.
- Shingfield KJ, Kairenius Äröla PA, Paillard D, Muetzel S, Ahvenjärvi S, Vanhatalo A, Huhtanen P, Toivonen V, Griinari JM and Wallace RJ 2012. Dietary fish oil supplements modify ruminal biohydrogenation, alter the flow of fatty acids at the oma- sum, and induce changes in the ruminal *Butyrivibrio* population in lactating cows. *Journal of Nutrition* 142, 1437–1448.
- Sirohi SK, Chaudhary PP, Singh N, Singh D and Puniya AK 2013. The 16S rRNA and mcrA gene based comparative diversity of methanogens in cattle fed on high fibre based diet. *Gene* 523, 161–166.
- Toral PG, Belenguer A, Shingfield KJ, Hervás G, Toivonen V and Frutos P 2012. Fatty acid composition and bacterial community changes in the rumen fluid of lactating sheep fed sunflower oil plus incremental levels of marine algae. *Journal of Dairy Science* 95, 794–806.
- Toral PG, Bernard L, Belenguer A, Rouel J, Hervás G, Chilliard Y and Frutos P 2016. Comparison of ruminal lipid metabolism in dairy cows and goats fed diets supplemented with starch, plant oil, or fish oil. *Journal of Dairy Science* 99, 301–316.
- Vlaeminck B, Fievez V, Cabrita ARJ, Fonseca AJM and Dewhurst RJ 2006. Factors affecting odd- and branched-chain fatty acids in milk: a review. *Animal Feed Science and Technology* 131, 389–417.
- Wang Q, Garrity GM, Tiedje JM and Cole JR 2007. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology* 73, 5261–5267.
- Yu Z and Morrison M 2004. Improved extraction of PCR-quality community DNA from digesta and fecal samples. *BioTechniques* 36, 808–812.
- Zened A, Combes S, Cauquil L, Mariette J, Klopp C, Bouchez O, Troegeler-Meynadier A and Enjalbert F 2013. Microbial ecology of the rumen evaluated by 454 GS FLX pyrosequencing is affected by starch and oil supplementation of diets. *FEMS Microbiology and Ecology* 83, 504–514.