


Journal Pre-proof

Integrated physiological and genetic data reveal key-traits for heat tolerance in tomato

S. Francesca , L. Vitale , S. Graci , M. Addonizio , A. Barone ,
M.M Rigano

PII: S2667-064X(24)00208-2
DOI: <https://doi.org/10.1016/j.stress.2024.100555>
Reference: STRESS 100555



To appear in: *Plant Stress*

Received date: 29 April 2024
Revised date: 12 July 2024
Accepted date: 29 July 2024

Please cite this article as: S. Francesca , L. Vitale , S. Graci , M. Addonizio , A. Barone , M.M Rigano , Integrated physiological and genetic data reveal key-traits for heat tolerance in tomato, *Plant Stress* (2024), doi: <https://doi.org/10.1016/j.stress.2024.100555>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 Published by Elsevier B.V.
This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Highlights

- Adult plants respond differently to prolonged heat stress than young plants
- Different physiological strategies can be employed in response to heat stress
- Efficient photosynthetic mechanisms are crucial for stress tolerance in tomato
- Key genes controlling heat tolerance were identified by GBS analysis

Journal Pre-proof

Integrated physiological and genetic data reveal key-traits for heat tolerance in tomato

Francesca S.^a, Vitale L.^b, Graci S.^a, Addonizio M.^a, Barone A.^a, Rigano M.M.^{a*}.

^aDepartment of Agricultural Science, University of Naples Federico II, Via Università 100, 80055 Portici, Italy

^bDepartment of Biology, Agriculture and Food Sciences, National Research Council, Institute for Agricultural and Forestry Systems in the Mediterranean, Portici, Italy

*corresponding: mrigano@unina.it

Abstract

As global temperatures continue to rise, high summer temperatures severely affect crop growth, reducing yield and quality. The projections of annual declines in crop yield require more in-depth multidisciplinary studies on plant tolerance to abiotic stresses. As tomato is a major crop in the Mediterranean region, its response to heat stress has become important to be addressed in order to identify those traits affecting stress tolerance. In this study, physiological and genomic analyses were performed on two heat-tolerant genotypes (LA3120 and E42) grown at high temperatures during the entire life cycle. The results showed that heat stress diversely affected gas exchange and fluorescence parameters in the two genotypes. In particular, E42 regulated the photosynthetic machinery under heat stress by modulating the electron transport chain, whereas LA3120 was less affected by the applied stress. Genotyping data obtained from a GBS (genotyping by sequencing) analysis were used to explore the genetic variability of both genotypes with the aim of identifying candidate genes that might regulate their stress response. These results further deepen our understanding of the physiological mechanisms activated in response to heat stress and allowed to select key traits that could be used in breeding program to select thermotolerant tomato genotypes.

Keywords: heat stress; novel genotypes; photosynthesis; Genotyping-by-sequencing; *Solanum lycopersicum*

1. Introduction

Mediterranean agriculture faces major challenges associated to climate change and sustainability. While climate change can improve temperature and rainfall patterns in some regions of the globe, the opposite is expected in the case of the Mediterranean region, identified as one of the most prominent hotspots in future climate-change projections, with 4-5°C rising temperatures and extreme rainfall patterns (www.fao.org/3/i3084e/i3084e16.pdf). In the European Mediterranean region, increases in the frequency of extreme climate events during specific plant developmental stages (e.g. heat stress during flowering period), together with higher rainfall intensity and longer dry spells, are likely to reduce yield of many plant species, including tomato (*Solanum lycopersicum* L.) (Alcamo et al., 2007). Indeed, the cultivation of tomato is deeply affected by high temperatures, since optimal reproduction and fruit set are limited in this species when the day's maximum temperatures exceed 32°C and the night's minimum falls below 21°C (Janni et al., 2020; Singh et al., 2022; Farinon et al., 2022; Graci & Barone, 2024). This is particularly worrisome considering that tomato is one of the most extensively cultivated species in temperate regions worldwide; indeed, according to FAOSTAT, globally 186.821 million tons of tomatoes were produced on 5,051,983 hectares in 2020 (FAO, 2022). Heat stress (HS) can cause different and often irreversible damages to plants altering biochemical, morphological, and physiological processes and leading to reductions in growth, early senescence, leaf abscission, damage to reproductive organs and pollen, fruit discoloration and reduced fruit quality (Singh et al., 2022; Gonzalo et al., 2022). Yield-related traits associated with the reproductive stages (such as total flower and fruit number, pollen viability, fruit set and final yield) are generally the main focus of the majority of the studies aimed at identifying elite tolerant lines (Gonzalo et al., 2021; Ruggieri et al., 2019). For this reason, high number of studies focusing on the genetic basis of these traits are reported (Graci & Barone, 2024). However, it is important to consider that heat tolerance is also dependent on the physiological status of vegetative organs. Indeed, physiological parameters are dramatically affected by HS and this can ultimately influence the final fruit production (Ahammed et al., 2018; Zahra et al., 2023). Plants subjected to high temperatures show a decrease in the rate of photosynthesis and in the accumulation of photosynthetic pigments in leaves (Shanmugam et al., 2013), water loss in aerial parts and alterations of membrane integrity (Li et al., 2015; Li et al., 2016). Both photosystems I and II (PSI and PSII) are affected by elevated temperatures due to the inhibition of light energy absorption, electron transport and energy distribution (Li et al., 2016), with PSII being more sensitive to HS (Yoshioka et al., 2006; Yan et al., 2013). However, the photosynthesis-related traits associated

to HS are still not known and very few studies focused on these traits. In the present study, in-depth physiological analyses were carried out on two different tolerant genotypes grown under prolonged high temperatures in order to identify key physiological traits related to HS tolerance. One genotype (LA3120, Malintka) is a well-known heat-tolerant genotype (Tomato Genetics Resource Centre, TGRC, University of California, CA, USA, www.tgrc-mvc.plantsciences.ucdavis.edu). The second used (coded E42) is a genotype that was selected in southern Italy and has been recently identified as heat-tolerant following many open field trials carried out over several years and in different locations (Ruggieri et al., 2019; Olivieri et al., 2020). The physiological responses to HS in E42 and LA3120 were previously investigated in a study from Francesca et al. (2022) that demonstrated the activation of different adaptive mechanisms in the two genotypes in young plants in response to a 3-week stress. However, it has been demonstrated that age-related eco-physiological mechanisms may be activated in response to abiotic stress that are deeply dependent on plant size and developmental stage (Zotz et al., 2001, Zhang et al., 2023). Therefore, in the present study a more in-depth physiological analysis was conducted on the same genotypes, which were here subjected to a prolonged HS over a period of 5-weeks. Our experimental design allowed us to identify different morpho-physiological traits related to HS tolerance. Moreover, these physiological data were combined with genotyping data in order to get further insights into the genetic basis of the identified tolerance-related traits. The emergence of Next Generation Sequencing (NGS) technologies able to identify millions of genome-wide single-nucleotide polymorphisms (SNPs) for high-throughput genotyping can allow the selection of candidate genes associated to specific phenotypic traits (Phan and Sim, 2017). Here the genomic analyses were conducted by using Reduced Representation Sequencing (RRS) data, a technology that uses one or more restriction enzymes to construct reduced representative libraries for the Illumina platform, resulting highly cost-effective (Peterson et al., 2012; Chung et al., 2017; Scheben et al., 2017).

Altogether, in this study we investigated the establishment of the different physiological and molecular mechanisms controlling stress adaptation in two genotypes known to be tolerant to elevated temperatures. The combination of physiological and genomic analyses was here able to provide key physiological parameters and related candidate genes, which could help breeders to select resilient genotypes that can cope with the current challenges of climate changes.

2. Materials and methods

2.1 Experimental Design and Plant Growth Conditions

This study was laid out at the Department of Agricultural Science of the University of Naples Federico II during the year 2023. The heat-tolerant tomato genotype LA3120 (Tomato Genetics Resource Centre, TGRC, University of California, CA, USA) and the tomato genotype E42 (University of Naples Federico II, Department of Agricultural Sciences) were used in the trial. The experiment was conducted with 10 biological replicates for each genotype per treatment. Plants were grown using a completely randomized block design. Seeds were sown in seed trays and after 20 days the seedlings were transferred to plastic pots (21 cm diameter) with commercial substrate. All pots were subjected to equivalent management throughout the experiment. One week after transplant, tomato seedlings were transferred to two growth chambers with a 16/8h photoperiod, PAR of $350 \mu\text{mol m}^{-2} \text{s}^{-1}$, RH of 65–75% and with climate settings of 26/18°C day/night temperatures in the control chamber and 35/18°C day/night temperatures in the hot chamber. The treatments were applied on 20-day-old seedlings and lasted for 5 weeks.

2.2 Assessment of Growth Parameters and Leaf Functional Traits

Five plants per genotype were randomly selected from each treatment to assess variability in growth traits. Leaf area (cm^2), specific leaf area (cm^2/g), fresh and dry shoot weight (g) were assessed. Leaf area was measured after scanning the leaves with the image analyzer ImageJ (Schneider et al., 2012). Specific leaf area (SLA) was measured as the ratio of leaf area to leaf dry mass. Leaf functional traits were determined on five leaves per genotype per treatment. The third leaf from the apex in each plant was chosen and the sampled leaves were in good condition, without wilting or disease. The tissues (leaves and shoot biomass) were dried in an oven at 80°C for 72 h to obtain dry biomass (DW).

2.3 Gas Exchange and Fluorescence Analyses

Gas exchange and fluorescence measurements were conducted on five healthy, young, fully expanded leaves (typically the 3rd leaf counting from the apical meristem) *per* genotype *per* treatment. Leaf gas exchange and chlorophyll fluorescence were simultaneously measured by using the Li6400 portable photosynthesis system (LiCor, Lincoln, NE, USA) integrated with Li6400-40 Leaf Chamber Fluorometer, which acts as both a leaf cuvette and light source/pulse-amplitude modulated fluorometer. Measurements were carried out in the morning with the following environmental parameters: photosynthetic photon flux density of 1200 $\mu\text{mol photons}$

$\text{m}^{-2} \text{s}^{-1}$, $400 \mu\text{mol CO}_2 \text{mol}^{-1}$, relative humidity 50–55%, and two fixed temperature regimens: 25 °C (considered as ambient control) and 35 °C (considered as heat treatment). Net photosynthesis (A_N), stomatal conductance (g_s), intercellular CO_2 concentration (C_i) and transpiration (T_T) were calculated according to von Caemmerer & Farquhar (1981) with the software of the Li6400. Steady-state fluorescence (F_s) and maximum fluorescence (F'_m) upon illumination were measured after a 0.8-s saturating flash of $7000 \mu\text{mol photons m}^{-2} \text{s}^{-1}$. Quantum yield of PSII electron transport (Φ_{PSII}) was calculated as reported in Maxwell & Johnson (2000). Intrinsic water use efficiency ($i\text{WUE}$) was calculated as A_N/g_s ratio ($\mu\text{mol/mol}$). Photochemical quenching (qP) was calculated according to Kramer et al. (2004) while the non-photochemical quenching of excitation in PSII-associated antenna complexes (NPQ_T) was obtained as proposed by Tietz et al. (2017). The electron transport rate (ETR) was calculated as: $\text{ETR} = 0.5 \times 0.84 \times I \times \Phi_{\text{PSII}}$, where 0.5 is the fraction of absorbed quanta used by PSII, 0.84 is the leaf absorbance, and I is the incident PPFD ($\mu\text{mol m}^{-2} \text{s}^{-1}$).

2.4 Determination of carotenoids and chlorophylls content

Carotenoid and chlorophyll assessment was performed following Wellburn (1994) and Zouari et al. (2014), as modified by Rigano et al. (2016). To obtain the lipophilic extract, 0.30 g of sample was extracted with 24 ml of acetone/hexane (40/60, v/v). The mixture was centrifuged at 20,000 g for 5 min at 4 °C. The supernatants were collected and stored at -20 °C until analysis. For the determination of carotenoids and chlorophyll a and b, the absorbance of the lipophilic extracts was read at 470, 663 and 645 nm, respectively. Results were expressed as $\text{mg } 100 \text{ g}^{-1} \text{ FW}$. Five separate biological replicates for each sample and three technical assays for each biological replicate were measured.

2.5 Hydrogen peroxide and malondialdehyde determination

The H_2O_2 content was quantified by a colorimetric method (Sergiev et al., 1997). Briefly, 500 mg of frozen leaf powder was extracted with 5 ml ice-cold 0.1% trichloroacetic acid (TCA), the mixture was incubated on ice for 15 min and centrifuged at 9,400 g for 15 min at 4 °C. To 500 μl of the supernatant, 500 μl of phosphate buffer 10 mM (pH 7.0) and 1 ml of potassium iodide (1 M) were added. The mixtures were then incubated in the dark for 40 min and measured at 525 nm using a nano photometer (Implen, Munich, Germany). Concentrations were expressed as $\text{mmol g}^{-1} \text{ FW}$. For the determination of malondialdehyde (MDA), 0.2 g of leaf sample was ground with 1 ml ice-cold 0.1% TCA. Samples were incubated on ice for 15

min and centrifuged at 9,400 g for 10 min at 4 °C. Then 0.25 ml supernatant was mixed with 1,250 ml reaction solution (TCA 20% + thiobarbituric acid [TBA] 0.5%), placed in a water bath at 95 °C for 30 min and measured at 532 and 600 nm with a Nano Photometer TM (Implen). The concentration was expressed as the amount of MDA-TBA complex (Zhang & Kirkham, 1996). For both analyses, five separate biological replicates were measured for each sample and three technical assays were measured for each biological replicate.

2.6 Leaf pigment index

Leaf pigment indexes such as chlorophyll content index, anthocyanin index, flavonol index, and nitrogen flavonol index (NFI, ratio of chlorophyll to flavonoids) were non-destructively measured on five fully expanded leaves per genotype per treatments using a portable instrument (multi-PigmentMeter MPM-100, Paris, France). This instrument uses ratio fluorescence to measure anthocyanin and flavonol content (ratios F660nm/ F525nm and F660nm/F325nm respectively) and leaf transmittance in the near and far infrared to measure chlorophyll content (T850/ T720). NFI was calculated as the ratio between chlorophyll (T850/T720) and flavonol (F660nm/F325nm) content.

2.7 Statistical analyses

Data were subjected to analysis of variance using a two-way ANOVA test. To separate means within each parameter, the student's t-test was performed. Differences at $p < 0.05$ were considered to be significant. To explore the overall data, we used the R environment for statistical computing and graphics R Core Team (2018). The PCA plot was obtained using R packages *factoMineR* version 2.9 (Le et al., 2008) and *factoextra* version 1.0.7 (Kassambara & Mundt, 2020).

2.8 Genome analyses

Genotyping analyses were conducted on the E42 and LA3120 genome data (available at <https://data.mendeley.com/datasets/hcr5fwtpf3/1>) obtained by using the Genotyping-By-Sequencing (GBS) platform through the double digest restriction-site associated (ddRAD) Illumina technology (Olivieri et al., 2020). Raw FASTQ files were processed in accordance with Graci et al. (2023). The number and distribution of homozygous variants for the alternative allele compared to Heinz were evaluated through a single-nucleotide polymorphism (SNP) density analysis by using the *snpcden* function of *VCFtools* (1 Mb non-overlapping

windows). Variants were annotated through the snpEff software (Cingolani et al., 2012) as reported by Graci et al. (2023). Finally, a list of genes potentially involved in the HS response of the two genotypes was prioritized by selecting those: I) showing HIGH and/or MODERATE impact on the translated protein and II) presenting Gene Ontology terms related to photosynthesis (GO:0015979), identified on the QuickGO database (Binns et al., 2009).

3. Results

3.1 Morpho-anatomical changes in tomato heat stressed plants

Morpho-anatomical changes in the two genotypes LA3120 and E42 were determined during exposure to HS in controlled chambers. Two-way ANOVA test evidenced that leaf area, specific leaf area and fresh weight biomass were significantly affected by the single factors heat stress and genotype, while the dry weight biomass was influenced only by the treatment factor (Table S1). The GxT interaction was not significant for all morpho-anatomical traits. Leaf area differed between genotypes under control condition, and the largest leaf area ($154.25 \pm 35.66 \text{ cm}^2$) was recorded in LA3120 (Table 1). In both genotypes a significant reduction in both leaf area and specific leaf area occurred under high temperatures. A strong decrease (approximately 50%) in fresh and dry biomass weight was also observed in both genotypes exposed to high temperatures compared to control plants.

3.2 Heat Stress modulate photochemical and fluorescence parameters in tomato plants

After exposure to thermal stress at 36 °C, both genotypes exhibited physiological and photochemical changes compared to non-treated plants (Fig. 1). ANOVA test indicated that genotypes, heat treatment and their interactions affected stomatal conductance (g_s), intrinsic water use efficiency ($iWUE$) and quantum yield of PSII electron transport (Φ_{PSII}). For the other photosynthetic parameters, there were no significant interactions between the factors, but only an effect of the single factor (genotype and treatment). On the contrary, internal CO_2 concentration (C_i) and non-photochemical quenching (NPQ_i) were only influenced by heat treatment and genotype, respectively (Table S1). Interestingly, in the absence of HS, significant differences were observed for all photosynthetic parameters between the two genotypes. The basal net photosynthetic values (A_N) under control conditions were significantly different between E42 and LA3120, presenting values of 27.58 and 14.20 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$, respectively (Fig. 1a). The CO_2 assimilation rate measured immediately after HS was significantly reduced (-20.34%) in E42, while in LA3120 no modifications after stress were

observed. Analyses of stomatal conductance (g_s) showed that it was different between genotypes (Fig. 1b) when no heating was applied, and g_s values were 0.79 and 0.26 mol m⁻² s⁻¹ in E42 and LA3120, respectively. After HS the stomatal conductance increased in both genotypes (+206% in E42 and +246% in LA3120). Under control conditions E42 displayed a higher leaf intercellular CO₂ concentration (C_i) compared to LA3120 (Fig. 1c); however, and consistent with g_s values, both genotypes showed a significant increase in C_i after HS. The intrinsic water use efficiency ($iWUE$) data are shown in Figure 1d and demonstrated that LA3120 had a 48% higher water use efficiency than E42 under control conditions. Nevertheless, $iWUE$ decreased significantly in both genotypes under HS, with LA3120 showing a more dramatic decrease compared to the control (-76.42%).

The parameters of Chl fluorescence were measured under control and HS conditions (Fig. 2). Interestingly, it was possible to note a constitutive difference in all the Chl fluorescence parameters between the two genotypes, with E42 presenting under control conditions higher values for Φ_{PSII} (quantum yield of PSII electron transport), qP (photochemical quenching) and ETR (electron transport rate) (Fig. 2a, b, d), and lower values for NPQ_t (non-photochemical quenching) (Fig. 2c) compared to LA3120. This confirmed the higher photosynthetic performance of E42 (see Fig. 1a). The parameters Φ_{PSII} and qP showed a significant decrease with the increasing temperatures only in E42 (Fig. 2 a,b). Moreover, as expected, HS caused a reduction in ETR that was statistically significant only in E42 (Fig. 2d), with a reduction of 30.41% relative to its basal value.

3.3 Effects of High Temperature Stress on the Photosynthetic Pigment Content, stress molecular markers and secondary metabolites.

Usually, HS alters chlorophylls and carotenoid contents in plants. In this study, carotenoid content decreased (-27.34%) under high temperature only in LA3120, while chlorophylls *a* and *b* content decreased under heat in both genotypes (Table 2) according with ANOVA test which indicated a significant effect only of the treatment (Table S1). That said, E42 subjected to heat showed a slight reduction of chlorophylls *a* and *b* (-18.52% and -18.87%, respectively) compared to the control, whereas a higher reduction in Chl *a* and *b* levels was observed in LA3120 treated plants (-28.96% and -33.66%, respectively). However, these differences in Chl *a* and *b* levels did not significantly affect the Chl *a/b* ratio in both genotypes. In order to check whether the stress conditions were responsible for oxidative damage in the two genotypes, the concentrations of H₂O₂ and MDA were determined. As expected, the levels of the two oxidative markers used were higher in heat-stressed plants than in control plants, being significantly

affected by the heat treatment (Table S1). No significant changes were observed in the production of flavonols and anthocyanins among treated and control plants in E42. On the contrary, an increase in flavonol content was observed in LA3120 after HS. Similar to the flavonol index, no significant differences in nitrogen-flavonol index (NFI) were found in E42 after HS. However, it was observed that LA3120 under HS exhibited lower NFI compared to control plants.

3.4 Principal components analysis

To provide a broad overview of the different analyses conducted on the two tomato genotypes in response to HS, a PCA was conducted. Based on our experimental data, three principal components (PCs) were associated with Eigenvalues >1 and accounted for 100% of the total variance, with PC1, PC2, PC3 accounting for 58.6%, 36.1%, and 5.4%, respectively (Table S2). PC1 was the primary drive for differences between treatment (Fig. 3 red light box) and the main parameters leading to this separation were oxidative markers and plant growth indices. There was also a genotype-dependent clustering, with the primary differences driven by PC2 (Fig. 3 blue light). The main parameters of PC2 were linked to primary and secondary metabolism.

3.6 Genome analyses

The VCF file of the two genotypes obtained from GBS sequencing data evidenced a final dataset of 30,046 and 9,562 variants for E42 and LA3120, respectively, respect to the reference genome of Heinz. The highest number of homozygous alternative polymorphisms compared to Heinz was detected in E42 (22,150), while LA3120 showed 7,474 homozygous polymorphisms. When looking at the distribution of the homozygous alternative variants compared to Heinz across the chromosomes (Table S3), E42 showed a high number of polymorphisms on chromosomes 1, 4, 7 and 12. On the other hand, LA3120 displayed a slightly higher concentration of variants on chromosome 12 compared to the others. In order to estimate the probable impact on proteins of homozygous SNPs and InDels detected in the two genotypes compared to Heinz, snpEff analysis was performed. Results showed that most of the variants had a putative MODIFIER impact (98% and 96.6%), followed by those with MODERATE (1.1% and 1.6%), LOW (0.8% and 1.5%) and HIGH (0.1% and 0.3%) impact in E42 and LA3120, respectively. E42 presented 31 HIGH and 288 MODERATE variants affecting 29 and 207 genes, respectively, while LA3120 showed 22 HIGH and 117 MODERATE polymorphisms affecting 21 and 94 genes, respectively.

A list of 101 GO terms (Table S4) was retrieved from the QuickGO database by selecting the GO:0015979 biological process term annotated for “photosynthesis” and the first 100 co-occurring GO terms, that were co-annotated to the GO:0015979 for the highest number of times. These features were combined and a list of 187 genes (Table S5) were prioritized since they showed 245 HIGH and/or MODERATE impact variants in the coding sequence of E42 and/or LA3120 and also exhibited annotation for GO terms related to photosynthesis. Among these genes, seven HIGH and 152 MODERATE polymorphisms affected 116 E42 genes, while five HIGH and 63 MODERATE mutations interested 54 LA3120 genes. In addition, both the genotypes displayed 17 genes exhibiting nine HIGH and nine MODERATE variants. Finally, from the list of 187 genes, a subset of 22 genes (Table 3) was highlighted, including those candidate genes that may explain the different physiological response mechanisms to HS in the two genotypes.

Interestingly, the highest number of variants (17 MODERATE) were exclusively found in the E42 genome, followed by one HIGH and ten MODERATE mutations affecting only LA3120, and two MODERATE polymorphisms common to both genotypes. Out of the 19 E42 variants, 14 mapped in the four most polymorphic chromosomes (1, 4, 7 and 12), while eight out of 13 LA3120 polymorphisms mapped on the other chromosomes.

4. Discussions

In this study, two tomato genotypes already identified as heat-tolerant based on yield-related traits were exposed to prolonged HS from the seedling stage up to the adult stage. Although high temperatures have been shown to affect all plant physiological processes, it remains unclear why high temperatures suppress plant growth and why some plants appear to be more heat tolerant than others. Over the past decade, several research studies focused on the effects of HS on specific physiological mechanisms, such as photosynthesis (Zahra et al., 2023). The majority of these studies were performed on plants at the seedling or vegetative stage, while little effort has been made to integrate the effects of high temperatures on these mechanisms in adult plants. A previous work (Francesca et al., 2021) carried out on the HS tolerant genotypes E42 and LA3120 on 20-day-old plants revealed the presence of valuable physiological traits for adaptation to stressful environments in both genotypes. However, since temperature is the main determinant of plant phenology, here we wondered how prolonged HS for almost the entire plant cycle might alter stress adaptation mechanisms. Looking at the overview provided by the PCA analysis, it can be observed that the experimental setup used in this study allowed

us to better identify the physiological traits affected by HS in both genotypes and, at the same time, to highlight the distinctive strategy adopted by them.

Generally, plants respond to changing environmental conditions by adjusting a suite of morphological traits, such as specific leaf area (SLA, the ratio of leaf area to leaf dry mass) and leaf size (Legner et al., 2014). In particular, given the direct relationship between SLA and light interception (Reich et al., 1998), SLA is intimately connected to resource acquisition and photosynthetic performance (Guo et al., 2022; Wright et al., 2004). In this work, we were able to decouple the effect of heat on leaf morphology from the genotype. Indeed, and contrary to what was observed previously on young plants (Francesca et al., 2021), here a general reduction in leaf area and SLA after HS was observed in both genotypes. This is in accordance with previous studies that demonstrated that leaves with lower SLA are better able to withstand and recover photosynthetic electron transport after high temperature stresses compared with leaves with greater SLA (Knight et al., 2003). It is generally recognized that the photosynthetic rate is associated with SLA; indeed, by increasing the light capture area per mass, high SLA improves photosynthesis (Goorman et al., 2011). In the present study both a higher SLA and a higher photosynthetic rate (A_N) were observed in E42 under control condition compared to LA3120. The A_N values observed in the present study were not observed previously when the same analyses were carried out on younger plants, stressing the fact that size and developmental stage deeply influence physiological traits. This observed high A_N could be the reason behind the high number of flowers and the high and stable final fruit yield observed in E42 in all the field trials previously conducted (Olivieri et al., 2020). In the present work, A_N was affected by the heat treatments in a genotype-dependent way, as it was negatively affected by stress only in E42, reaching the same A_N values observed in LA3120 both under stressful and non-stressful conditions. Under HS, generally, stomatal conductance (g_s) rises to increase intracellular CO_2 concentration and facilitate water loss by transpiration (Dos Santos et al., 2022). Evaporative cooling through open stomata mitigates the negative effects of supra-optimal temperatures and have a positive effect on photosynthesis, yield, and plant survival (Ameje et al., 2012; Urban et al., 2017). Accordingly, under HS an increase in stomatal conductance (g_s) and intracellular CO_2 concentration (C_i) was here observed, as expected. A stable photosynthetic rate combined with an increased stomatal conductivity, as shown here in LA3120, has been previously reported in heat-tolerant tomato and pepper cultivars (Zhou et al., 2015). On the contrary, the A_N decrease accompanied by the increases in g_s and C_i values observed in E42 after HS could indicate biochemical limitations (Zhou et al., 2015).

Fluorescence parameters measured under non-stressed condition confirmed the higher photosynthetic performances of E42, which showed higher levels of quantum yield of PSII (Φ_{PSII}), photochemical quenching (qP) and electron transport rate (ETR), and lower levels of non-photochemical quenching (NPQ) compared to LA3120. After HS, only in E42 a decrease in qP, Φ_{PSII} and ETR was demonstrated. This observation, which well fits with the gas exchange data previously discussed, confirmed that the photosynthetic apparatus of E42 was more sensible than LA3120 to HS.

Another consequence of abiotic stress in plants is the oxidative damage. This leads to the overproduction of reactive oxygen species (ROS), which are primarily directed against the lipids of the membrane (Francesca et al., 2022). Accordingly, an increase in H_2O_2 content and lipid peroxidation under HS was here observed in stressed plants. These changes were accompanied by an increase in flavonol content only in LA3120, while E42 did not alter its already high basal levels of flavonols and anthocyanins.

In order to identify the genetic basis of the identified physiological traits involved in the response mechanism to HS, a genomic analysis was here performed. Data obtained from GBS high-throughput genotyping technology were investigated to identify candidate genes involved in HS-tolerance mechanisms in E42 and LA3120. Francesca et al. (2021) already used the same dataset thus highlighting five genes showing E42 private variants with HIGH impact on the translated proteins, while no genes were previously discussed to explain the LA3120 response. Moreover, the analysis carried out in that study did not focus on all the MODERATE polymorphisms, which contribute to change the protein function by altering the amino acid sequence. Herein, attention was focused on all the genes showing HIGH and MODERATE polymorphisms known to be involved in the photosynthetic process by using their gene ontology (GO) annotation. This strategy allowed us to identify 22 candidate genes among which 13 presented variants exclusively in E42 and seven in LA3120. The genes Solyc01g056890 coding for the protein CURVATURE and the Solyc07g021200 coding for the Ribulose biphosphate carboxylase large chain were already highlighted by Francesca et al. (2021) in E42. The presence of these variants is in accordance with physiological data here observed. Indeed, the A_N decrease observed in E42 after HS could have been caused by modification in the gene Solyc07g021200.

Interestingly, the analyses carried out in the present study allowed the identification of another gene in E42, the gene Solyc02g090890, coding for a zeaxanthin epoxidase showing one missense variant. Zeaxanthin is a crucial carotenoid and, as a free pigment in thylakoids, plays a key role in antioxidation and stabilizing the membrane structure by inhibiting and scavenging

reactive oxygen species and suppressing lipid peroxidation (Tang et al., 2021). One hypothesis is that the MODERATE polymorphism in this gene may have blocked the zeaxanthin epoxidase activity, limiting the conversion of violaxanthin to zeaxanthin via the so-called xanthophyll cycle. Experiments have shown that NPQ cannot be activated when the xanthophyll cycle is blocked (Gilmore, 1997; Dreuw et al., 2003). Accordingly, our data showed lower NPQ values in E42 compared to LA3120 under all conditions. The block of the xanthophyll cycle could also have promoted in E42 the synthesis of the 9-cis-epoxycarotenoid dioxygenase (NCED) enzyme which is reported to positively regulate plant tolerance to abiotic stresses (Zhang et al., 2022). Considering LA3120, GBS analysis evidenced the presence of four missense variants in the Solyc03g124068 coding for the Ent-kaurene synthase. This is the key enzyme for the biosynthesis of gibberellins (GAs), which catalyze the formation of ent-kaurene, the precursor for GAs (Yang et al., 2021). This hormone is involved in flowering, pollen formation and pollen tube growth and promotes tomato fruit set and fruit initiation (Sponsel, 2016, Graci & Barone, 2024). It is also involved in leaf expansion, dry matter accumulation, photosynthesis, and transpiration rate. A clear correlation between this hormone and tolerance to drought and HS has also been found (Shohat et al., 2021). Recently, Guo et al. (2022) used gibberellic acid as a foliar treatment to alleviate HS in tomato plants. The polymorphisms present in the Ent-kaurene synthase gene may have had an impact on GA synthesis and accumulation. This well correlates with the physiological data observed here and in previous paper on LA3120 (Francesca et al., 2021). This is also in accordance with the LA3120 phenotypic data recorded in different years and environments, that showed a reduction of the time required in this genotype for flower and fruit formation (Ayenan et al., 2021; Olivieri et al., 2021). This could be a key mechanism in LA3120 that allows it to precociously form flower and fruit before the onset of the high summer temperatures typical of the Mediterranean area. Other two genes (Solyc12g014250 coding for the Phosphoenolpyruvate carboxylase and Solyc12g056940 coding for the Acetyl CoA carboxylase) were here identified that showed one and two missense variants in LA3120, respectively. Also, these two genes may play pivotal role in plants during carbon metabolism in photosynthesis, physicochemical processes, and tolerance to various abiotic stresses (Waseem & Ahmad, 2019; Upadhyay et al., 2020).

Altogether, in the present study a more in-depth physiological analysis was conducted on two thermotolerant genotypes, which were here subjected to a prolonged HS over a period of 5-weeks. The genotypes were able to employ different defense mechanisms in response to HS revealing key tolerance traits. The candidate genes here identified by combining the phenotypic

and genotypic analyses carried out in this work, helped to dissect these complex traits, and could explain the different physiological response to stress observed. These genes should be further studied and could be used in future breeding programs in order to improve tolerance to abiotic stress in modern cultivars.

Funding

This study was carried out within the Agritech National Research Center and received funding from the European Union Next-Generation EU (Piano Nazionale di Ripresa e Resilienza (PNRR)—Missione 4 Componente 2, Investimento 1.4—D.D. 1032 17/06/2022, CN00000022). This manuscript reflects only the authors' views and opinions. Neither the European Union nor the European Commission can be considered responsible for them.

Author contributions

S.F., S.G., M.M.R. and A.B. conceptualized the research. S.F., S.G., M.A. and L.V. performed the experiments. S.F., S.G., M.A. and L.V. analyzed the data. S.F., S.G., L.V., M.M.R. and A.B. wrote the manuscript. All authors reviewed and approved the final manuscript.

Data availability statement

All data are available in the manuscript or Supplementary files. The GBS raw data that support the findings of this study are available here: <https://data.mendeley.com/datasets/hcr5fwtpf3/1>

Conflict of interest statement

The authors have no conflicts of interest to declare.

References

Ahammed, G. J., Xu, W., Liu, A., & Chen, S. (2018). COMT1 silencing aggravates heat stress-induced reduction in photosynthesis by decreasing chlorophyll content, photosystem II activity, and electron transport efficiency in tomato. *Frontiers in Plant Science*, 9, 386258.

Alcamo, J., Flörke, M., & Märker, M. (2007). Future long-term changes in global water resources driven by socio-economic and climatic changes. *Hydrological Sciences Journal*, 52(2), 247-275.

Ameye M, Wertin TM, Bauweraerts I, McGuire MA, Teskey RO, Steppe K. (2012). The effect of induced heat waves on *Pinus taeda* and *Quercus rubra* seedlings in ambient and elevated CO₂ atmospheres. *The New Phytologist* 196, 448–461.

Ayenon, M. A. T., Danquah, A., Hanson, P., Asante, I. K., & Danquah, E. Y. (2021). Identification of new sources of heat tolerance in cultivated and wild tomatoes. *Euphytica*, 217(3), 33.

Binns, D., Dimmer, E., Huntley, R., Barrell, D., O'donovan, C., & Apweiler, R. (2009). QuickGO: a web-based tool for Gene Ontology searching. *Bioinformatics*, 25(22), 3045-3046.

Chung, Y. S., Choi, S. C., Jun, T.-H., & Kim, C. (2017). Genotyping-by-sequencing: a promising tool for plant genetics research and breeding. *Horticulture, Environment, and Biotechnology*, 58, 425–431.

Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., Land, S. J., Lu, X., & Ruden, D. M. (2012). A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly*, 6(2), 80–92.

Dreuw, A., Fleming, G. R., & Head-Gordon, M. (2003). Chlorophyll fluorescence quenching by xanthophylls. *Physical Chemistry Chemical Physics*, 5(15), 3247-3256.

Dos Santos, T. B., Ribas, A. F., de Souza, S. G. H., Budzinski, I. G. F., & Domingues, D. S. (2022). Physiological responses to drought, salinity, and heat stress in plants: a review. *Stresses*, 2(1), 113-135.

FAO (2022). Land and Water. Tomato. Available online at: <https://www.fao.org/land-water/databases-and-software/crop-information/tomato/en/>.

Farinon, B., Picarella, M. E., & Mazzucato, A. (2022). Dynamics of fertility-related traits in tomato landraces under mild and severe heat stress. *Plants*, 11(7), 881.

Francesca, S., Vitale, L., Arena, C., Raimondi, G., Olivieri, F., Cirillo, V., ... & Rigano, M. M. (2021). The efficient physiological strategy of a novel tomato genotype to adapt to chronic combined water and heat stress. *Plant Biology*, 24(1), 62-74.

Francesca, S., Najai, S., Zhou, R., Decros, G., Cassan, C., Delmas, F., ... & Rigano, M. M. (2022). Phenotyping to dissect the biostimulant action of a protein hydrolysate in tomato plants under combined abiotic stress. *Plant Physiology and Biochemistry*, 179, 32-43.

Gilmore, A. M. (1997). Mechanistic aspects of xanthophyll cycle-dependent photoprotection in higher plant chloroplasts and leaves. *Physiologia Plantarum*, 99(1), 197-209.

Gonzalo, M. J., Nájera, I., Baixauli, C., Gil, D., Montoro, T., Soriano, V., ... & Monforte, A. J. (2021). Identification of tomato accessions as source of new genes for improving heat tolerance: from controlled experiments to field. *BMC Plant Biology*, 21, 1-28.

Gonzalo, M. J., da Maia, L. C., Nájera, I., Baixauli, C., Giuliano, G., Ferrante, P., ... & Monforte, A. J. (2022). Genetic control of reproductive traits under different temperature regimes in inbred line populations derived from crosses between *S. pimpinellifolium* and *S. lycopersicum* accessions. *Plants*, 11(8), 1069.

Goorman, R., Bartual, A., Paula, S., & Ojeda, F. (2011). Enhancement of photosynthesis in post-disturbance resprouts of two co-occurring Mediterranean *Erica* species. *Plant Ecology*, 212, 2023-2033.

Graci, S., Ruggieri, V., Francesca, S., Rigano, M. M., & Barone, A. (2023). Genomic insights into the origin of a thermotolerant tomato line and identification of candidate genes for heat stress. *Genes*, 14(3), 535.

Graci, S., & Barone, A. (2024). Tomato plant response to heat stress: a focus on candidate genes for yield-related traits. *Frontiers in Plant Science*, 14, 1245661.

Guo Z., Yan Z., Majcher B.M., Lee C.K.F., Zhao Y., Song G., Wang B., Wang X., Deng Y., Michaletz S.T., Ryu Y., Ashton L.A., Lam H.M., Wong M.S., Liu L., Wu J. (2022) Dynamic

biotic controls of leaf thermoregulation across the diel timescale. *Agricultural and Forest Meteorology*, 315, 108827.

Janni, M., Gulli, M., Maestri, E., Marmioli, M., Valliyodan, B., Nguyen, H. T., & Marmioli, N. (2020). Molecular and genetic bases of heat stress responses in crop plants and breeding for increased resilience and productivity. *Journal of experimental botany*, 71(13), 3780-3802.

Kassambara A, Mundt F (2020). *factoextra: Extract and Visualize the Results of Multivariate Data Analyses*. R package version 1.0.7 <https://CRAN.R-project.org/package=factoextra>.

Knight, C. A., & Ackerly, D. D. (2003). Evolution and plasticity of photosynthetic thermal tolerance, specific leaf area and leaf size: congeneric species from desert and coastal environments. *New Phytologist*, 160(2), 337-347.

Kramer D.M., Avenson T.J., Edwards G.E. (2004) Dynamic flexibility in the light reactions of photosynthesis governed by both electron and proton transfer reactions. *Trends in Plant Science*, 9, 349–357.

Le, S., Josse, J. & Husson, F. (2008). *FactoMineR: An R Package for Multivariate Analysis*. *Journal of Statistical Software*. 25(1). pp. 1-18

Legner, N., Fleck, S., & Leuschner, C. (2014). Within-canopy variation in photosynthetic capacity, SLA and foliar N in temperate broad-leaved trees with contrasting shade tolerance. *Trees*, 28, 263-280.

Li, H., Ahammed, G. J., Zhou, G., Xia, X., Zhou, J., Shi, K., ... & Zhou, Y. (2016). Unraveling main limiting sites of photosynthesis under below-and above-ground heat stress in cucumber and the alleviatory role of luffa rootstock. *Frontiers in Plant Science*, 7, 183933.

Li, X., Ahammed, G. J., Zhang, Y. Q., Zhang, G. Q., Sun, Z. H., Zhou, J., ... & Shi, K. (2015). Carbon dioxide enrichment alleviates heat stress by improving cellular redox homeostasis through an ABA-independent process in tomato plants. *Plant Biology*, 17(1), 81-89.

Maxwell, K., & Johnson, G. N. (2000). Chlorophyll fluorescence—a practical guide. *Journal of experimental botany*, 51(345), 659-668.

Olivieri, F., Calafiore, R., Francesca, S., Schettini, C., Chiaiese, P., Rigano, M. M., & Barone, A. (2020). High-throughput genotyping of resilient tomato landraces to detect candidate genes involved in the response to high temperatures. *Genes*, 11(6), 626.

Olivieri, F., Graci, S., Francesca, S., Rigano, M. M., & Barone, A. (2021). Accelerating the development of heat tolerant tomato hybrids through a multi-traits evaluation of parental lines combining phenotypic and genotypic analysis. *Plants*, 10(10), 2168.

Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double digest RADseq: an inexpensive method for de novo SNP discovery and genotyping in model and non-model species. *PloS One*, 7(5), e37135.

Phan, N. T., & Sim, S.-C. (2017). Genomic tools and their implications for vegetable breeding. *Horticultural Science and Technology*, 35(2), 149–164.

Reich, P. B., Tjoelker, M. G., Walters, M. B., Vanderklein, D. W., & Buschena, C. (1998). Close association of RGR, leaf and root morphology, seed mass and shade tolerance in seedlings of nine boreal tree species grown in high and low light. *Functional ecology*, 12(3), 327-338.

Rigano, M. M., Arena, C., Di Matteo, A., Sellitto, S., Frusciante, L., & Barone, A. (2016). Eco-physiological response to water stress of drought-tolerant and drought-sensitive tomato genotypes. *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology*, 150(4), 682-691.

Ruggieri, V., Calafiore, R., Schettini, C., Rigano, M. M., Olivieri, F., Frusciante, L., & Barone, A. (2019). Exploiting genetic and genomic resources to enhance heat-tolerance in tomatoes. *Agronomy*, 9(1), 22.

Shanmugam, S., Kjaer, K. H., Ottosen, C. O., Rosenqvist, E., Kumari Sharma, D., & Wollenweber, B. (2013). The Alleviating Effect of Elevated CO₂ on Heat Stress Susceptibility

of Two Wheat (*Triticum aestivum* L.) Cultivars. *Journal of Agronomy and Crop Science*, 199(5), 340-350.

Scheben, A., Batley, J., & Edwards, D. (2017). Genotyping-by-sequencing approaches to characterize crop genomes: choosing the right tool for the right application. *Plant Biotechnology Journal*, 15(2), 149–161.

Schneider, C.A.; Rasband, W.S.; Eliceiri, K.W (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods*, 9, 671-675.

Sergiev, I., Alexieva, V., & Karanov, E. (1997). Effect of spermine, atrazine and combination between them on some endogenous protective systems and stress markers in plants. *Comptes rendus de l'Academie bulgare des Sciences*, 51(3), 121-124.

Shohat, H., Eliaz, N. I., & Weiss, D. (2021). Gibberellin in tomato: Metabolism, signaling and role in drought responses. *Molecular Horticulture*, 1(1), 15.

Singh, A. K., Mishra, P., Kashyap, S. P., Karkute, S. G., Singh, P. M., Bahadur, A., & Behera, T. K. (2022). Molecular insights into mechanisms underlying thermo-tolerance in tomato. *Frontiers in Plant Science*, 13, 1040532.

Sponsel, V. M. (2016). Signal achievements in gibberellin research: the second half-century. *Annual Plant Reviews, Volume 49: Gibberellins, The*, 1-36.

Tang, C., Xie, J., Lv, J., Li, J., Zhang, J., Wang, C., & Liang, G. (2021). Alleviating damage of photosystem and oxidative stress from chilling stress with exogenous zeaxanthin in pepper (*Capsicum annuum* L.) seedlings. *Plant Physiology and Biochemistry*, 162, 395-409.

Tietz, S., Hall, C. C., Cruz, J. A., & Kramer, D. M. (2017). NPQ (T): a chlorophyll fluorescence parameter for rapid estimation and imaging of non-photochemical quenching of excitons in photosystem-II-associated antenna complexes. *Plant, Cell & Environment*, 40, 8, 1243-1255.

Upadhyay, R. K., Fatima, T., Handa, A. K., & Mattoo, A. K. (2020). Polyamines and their biosynthesis/catabolism genes are differentially modulated in response to heat versus cold stress in tomato leaves (*Solanum lycopersicum* L.). *Cells*, 9(8), 1749.

Urban, J., Ingwers, M. W., McGuire, M. A., & Teskey, R. O. (2017). Increase in leaf temperature opens stomata and decouples net photosynthesis from stomatal conductance in *Pinus taeda* and *Populus deltoides* x *nigra*. *Journal of experimental botany*, 68(7), 1757-1767.

Waseem, M., & Ahmad, F. (2019). The phosphoenolpyruvate carboxylase gene family identification and expression analysis under abiotic and phytohormone stresses in *Solanum lycopersicum* L. *Gene*, 690, 11-20.

Wellburn, A. R. (1994). The spectral determination of chlorophylls a and b, as well as total carotenoids, using various solvents with spectrophotometers of different resolution. *Journal of plant physiology*, 144(3), 307-313.

Wright, I. J., Groom, P. K., Lamont, B. B., Poot, P., Prior, L. D., Reich, P. B., ... & Westoby, M. (2004). Leaf trait relationships in Australian plant species. *Functional plant biology*, 31(5), 551-558.

Yan, K., Chen, P., Shao, H., Shao, C., Zhao, S., & Brestic, M. (2013). Dissection of photosynthetic electron transport process in sweet sorghum under heat stress. *PLoS One*, 8(5), e62100.

Yang, M., Liu, G., Yamamura, Y., & Fu, J. (2021). Identification and functional characterization of ent-kaurene synthase gene in *Ilex latifolia*. *Beverage Plant Research*, 1(1), 1-7.

Yoshioka, M., Uchida, S., Mori, H., Komayama, K., Ohira, S., Morita, N., ... & Yamamoto, Y. (2006). Quality control of photosystem II: cleavage of reaction center D1 protein in spinach thylakoids by FtsH protease under moderate heat stress. *Journal of Biological Chemistry*, 281(31), 21660-21669.

Zahra, N., Hafeez, M. B., Ghaffar, A., Kausar, A., Al Zeidi, M., Siddique, K. H., & Farooq, M. (2023). Plant photosynthesis under heat stress: Effects and management. *Environmental and Experimental Botany*, 206, 105178.

Zhang J., Kirkham M. (1996) Antioxidant responses to drought in sunflower and sorghum seedlings. *New Phytologist*, 132, 361–373.

Zhang, W., Dong, X. M., Zhang, Y. W., Fan, Z. X., & Zhang, S. B. (2023). Age-related differences in physiological and metabolic responses of *Pleione aurita* (Orchidaceae) pseudobulbs to drought stress and recovery. *Plant Physiology and Biochemistry*, 197, 107655.

Zhang, Y., Liu, X., Su, R., Xiao, Y., Deng, H., Lu, X., ... & Zhang, G. (2022). 9-cis-epoxycarotenoid dioxygenase 1 confers heat stress tolerance in rice seedling plants. *Frontiers in Plant Science*, 13, 1092630.

Zotz, G., Hietz, P., & Schmidt, G. (2001). Small plants, large plants: the importance of plant size for the physiological ecology of vascular epiphytes. *Journal of experimental botany*, 52(363), 2051-2056.

Zhou R., Yu X., Kjær K.H., Rosenqvist E., Ottosen C.O., Wu Z. (2015) Screening and validation of tomato genotypes under heat stress using Fv/Fm to reveal the physiological mechanism of heat tolerance. *Environmental and Experimental Botany*, 118, 1–11.

Zouari, I., Salvioli, A., Chialva, M., Novero, M., Miozzi, L., Tenore, G. C., ... & Bonfante, P. (2014). From root to fruit: RNA-Seq analysis shows that arbuscular mycorrhizal symbiosis may affect tomato fruit metabolism. *Bmc Genomics*, 15, 1-19.

Table 1. Plant growth parameters and leaf traits (leaf area, specific leaf area, shoot and dry biomass weight) of two tomato genotypes under control (CTRL) and high temperatures conditions (HEAT). The data are mean \pm standard deviation of five biological replicates. Within each tomato genotypes, asterisks indicate significant differences between treatments based on Student's *t*-test ($p < 0.05^*$; $p < 0.01^{**}$; $p < 0.001^{***}$).

| Parameters | E42 | | LA3120 | |
|-------------------------------------------------------|--------------------|----------------------------------|--------------------|----------------------------------|
| | CTRL | HEAT | CTRL | HEAT |
| Leaf area (cm ²) | 87.83 \pm 11.42 | 44.47 \pm 15.76 ^{**} | 154.25 \pm 35.66 | 107.81 \pm 25.34 [*] |
| Specific leaf area (cm ² g ⁻¹) | 314.97 \pm 23.08 | 191.81 \pm 41.45 ^{**} | 257.53 \pm 41.84 | 142.62 \pm 27.73 ^{**} |
| Fresh weight biomass (g) | 436.57 \pm 86 | 218.8 \pm 34.9 ^{**} | 505.2 \pm 44.98 | 263.3 \pm 63.05 ^{**} |
| Dry weight biomass (g) | 84.25 \pm 24.36 | 39 \pm 4.50 [*] | 96.98 \pm 25.828 | 49.28 \pm 5.16 ^{**} |

Table 2. Pigments content, molecular stress markers and secondary metabolites of two tomato genotypes under control (CTRL) and high temperatures conditions (HEAT). Data are means \pm standard deviation (n = 5). Within each tomato line, asterisks indicate significant differences between treatments based on Student's *t*-test (p<0.05*; p<0.01**; p<0.001***).

| Parameters | E42 | | LA3120 | |
|-------------------------------------------------------|-------------------|----------------------|-------------------|----------------------|
| | CTRL | HEAT | CTRL | HEAT |
| Carotenoids (mg 100g FW ⁻¹) | 17.07 \pm 2.55 | 16.40 \pm 3.30 | 21.14 \pm 2.21 | 15.36 \pm 1.42*** |
| Chla (mg 100g FW ⁻¹) | 94.94 \pm 5.64 | 77.35 \pm 11.87** | 100.67 \pm 7.78 | 71.51 \pm 6.27*** |
| Chlb (mg 100g FW ⁻¹) | 48.57 \pm 5.95 | 39.40 \pm 5.11* | 53.85 \pm 10.21 | 35.72 \pm 3.90*** |
| Chla/b (mg 100g FW ⁻¹) | 1.92 \pm 0.10 | 1.95 \pm 0.07 | 1.96 \pm 0.19 | 2.00 \pm 0.05 |
| Hydrogen peroxide (mmol g ⁻¹) | 0.086 \pm 0.03 | 0.168 \pm 0.03* | 0.086 \pm 0.01 | 0.156 \pm 0.01** |
| Lipid peroxidation (nmol MDA-TBA g FW ⁻¹) | 0.032 \pm 0.002 | 0.052 \pm 0.004*** | 0.035 \pm 0.004 | 0.054 \pm 0.001*** |
| Flavonol content | 0.26 \pm 0.13 | 0.26 \pm 0.08 | 0.08 \pm 0.03 | 0.17 \pm 0.07* |
| Anthocyanins content | 0.073 \pm 0.005 | 0.076 \pm 0.010 | 0.050 \pm 0.020 | 0.053 \pm 0.02 |
| Nitrogen flavonol index (NFI) | 2.24 \pm 0.83 | 2.14 \pm 0.87 | 3.54 \pm 1.28 | 1.67 \pm 0.57** |

Table 3. List of the 22 E42 and LA3120 candidate genes showing HIGH and/or MODERATE variants in the gene coding sequences and also exhibiting GO terms related to photosynthesis.

| Gene | Position (SL4.0) | Impact | Effect | Genotype | Annotation ITAG4.1 |
|--------------------|------------------|--------------|----------------------|----------------|-----------------------------------------------------------------------------------------|
| Solyc01g05689 0 | 50,654,06 8 | MODERAT E | missense_varian t | E42 | protein CURVATURE |
| | 50,654,06 9 | MODERAT E | missense_varian t | E42 | THYLAKOID 1A, chloroplastic-like |
| Solyc01g09071 0 | 76,596,31 2 | MODERAT E | missense_varian t | E42 | Malate dehydrogenase |
| Solyc02g01190 0 | 12,636,27 1 | MODERAT E | missense_varian t | E42 | NAD(P)H-quinone oxidoreductase subunit 2, chloroplastic |
| Solyc02g09089 0 | 50,411,60 4 | MODERAT E | missense_varian t | E42 | zeaxanthin epoxidase |
| Solyc03g12406 8 | 65,296,48 2 | MODERAT E | missense_varian t | LA3120 | Ent-kaurene synthase |
| | 65,296,50 0 | MODERAT E | missense_varian t | LA3120 | |
| | 65,296,53 3 | MODERAT E | missense_varian t | LA3120 | |
| | 65,296,78 9 | MODERAT E | missense_varian t | LA3120 | |
| Solyc04g01421 0 | 4,534,561 E | MODERAT E | missense_varian t | E42 | RNA helicase DEAH-box12 |
| Solyc04g03982 0 | 12,288,60 3 | MODERAT E | missense_varian t | E42 | Acetyl-coenzyme A carboxylase carboxyl transferase subunit beta, chloroplastic |
| Solyc04g07687 0 | 59,787,25 8 | MODERAT E | missense_varian t | E42, LA3120 | Glutamyl-tRNA reductase |
| Solyc05g05500 0 | 64,162,77 4 | MODERAT E | missense_varian t | LA3120 | Cysteine desulfurase |
| Solyc07g02120 0 | 16,851,87 0 | MODERAT E | missense_varian t | E42 | Ribulose bisphosphate carboxylase large chain |
| Solyc07g02120 5 | 16,852,04 0 | MODERAT E | missense_varian t | E42 | DNA-directed RNA polymerase subunit beta |
| | 16,852,05 9 | MODERAT E | missense_varian t | E42 | |
| Solyc07g05602 0 | 63,838,95 5 | MODERAT E | missense_varian t | E42 | Translation initiation factor IF-2 |
| Solyc07g06293 0 | 65,398,59 3 | MODERAT E | missense_varian t | E42 | Ribosomal L.1 |

| | | | | | |
|--------------------|----------------|----------|------------------|----------------|--------------------------------------------------------------------------|
| Solyc08g07648 0 | 58,602,80 1 | HIGH | stop_gained | LA3120 | Plastid lipid-associated protein chloroplastic-like SEC12-like protein.1 |
| Solyc11g00778 0 | 2,058,238 | MODERATE | missense_variant | E42 | |
| Solyc11g01986 0 | 9,795,826 | MODERATE | missense_variant | LA3120 | Adenylate isopentenyltransferase |
| Solyc11g02050 7 | 11,424,15 2 | MODERATE | missense_variant | E42, LA3120 | Adenylate isopentenyltransferase |
| Solyc11g06593 0 | 49,569,43 8 | MODERATE | missense_variant | E42 | Xanthine dehydrogenase |
| Solyc12g00999 0 | 3,175,319 | MODERATE | missense_variant | E42 | Signal recognition particle receptor subunit alpha |
| | 3,175,321 | MODERATE | missense_variant | E42 | |
| | 3,175,323 | MODERATE | missense_variant | E42 | |
| Solyc12g01425 0 | 5,109,796 | MODERATE | missense_variant | LA3120 | Phosphoenolpyruvate carboxylase |
| Solyc12g04445 0 | 59,073,11 4 | MODERATE | missense_variant | LA3120 | Protein trichome birefringence-like 42 |
| Solyc12g05694 0 | 62,629,17 4 | MODERATE | missense_variant | LA3120 | Acetyl CoA carboxylase |
| | 62,629,53 2 | MODERATE | missense_variant | LA3120 | |

Figure legends

Figure 1. Gas exchange in leaves of two tomato genotypes under control (CTRL) and high temperatures conditions (HEAT). (a) Net photosynthesis (A_N), (b) stomatal conductance (g_s), (c) intracellular CO_2 concentration (C_i), (d) intrinsic water use efficiency (iWUE). Data are mean \pm standard error (SE) ($n = 5$). Asterisks indicate significant differences between genotypes and between treatments based on Student's t -test ($p < 0.05^*$; $p < 0.01^{**}$; $p < 0.001^{***}$).




Figure 2. Fluorescence measurements in leaves of two tomato genotypes under control (CTRL) and high temperatures conditions (HEAT). (a) quantum yield of PSII (Φ_{PSII}), (b) photochemical quenching (qP), (c) nonphotochemical quenching (NPQ_T), (d) electron transport rate (ETR). Data are mean \pm standard error (SE) (n = 5). Asterisks indicate significant differences between genotypes and between treatments based on Student's *t*-test (p < 0.05*; p < 0.01**; p < 0.001***).






Figure 3. Principal component loading plot and scores of principal components analysis (PCA) in two tomato genotypes under control (C) and heat (H) stress.



Graphical Abstract




Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The author is an Editorial Board Member/Editor-in-Chief/Associate Editor/Guest Editor for [Journal name] and was not involved in the editorial review or the decision to publish this article.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:



Journal Pre-proof