




**Multicomponent, Fragment-Based, Synthesis of New  
Natural-Based Polyphenols and their Inhibiting Activity on  
Beta-Amyloid Oligomerization**

Journal:	<i>Organic &amp; Biomolecular Chemistry</i>
Manuscript ID	OB-ART-08-2017-002182
Article Type:	Paper
Date Submitted by the Author:	31-Aug-2017
Complete List of Authors:	<p>Lambruschini, Chiara; University of Genova, Dipartimento di Chimica e Chimica Industriale  Galante, Denise; Consiglio Nazionale delle Ricerche, Istituto per lo Studio delle Macromolecole  Moni, Lisa; University of Genova, Dipartimento di Chimica e Chimica Industriale  Ferraro, Francesco; University of Genova, Dipartimento di Chimica e Chimica Industriale  Gancia, Giulio; Consiglio Nazionale delle Ricerche, Istituto per lo Studio delle Macromolecole  Riva, Renata; University of Genova, Dept. of Chemistry and Industrial Chemistry  Traverso, Alessia; University of Genova, Dipartimento di Chimica e Chimica Industriale  Banfi, Luca; University of Genova, Dipartimento di Chimica e Chimica Industriale  D'Arrigo, Cristina; Consiglio Nazionale delle Ricerche, Istituto per lo Studio delle Macromolecole</p>
<p>Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.</p>	
<p>Scheme_1.cdx  Scheme_2.cdx  Scheme_3.cdx  Scheme_4.cdx  Figure 1.cdx</p>	



# Organic & Biomolecular Chemistry (OBC)

Impact factor 3.562\*

OBC aims to provide authors with a quick decision on their manuscript. We would appreciate you assisting us in providing this service by submitting your review by the deadline in the invitation message.

**When assessing this manuscript please take the following into account:**

- OBC articles should report **new work** which makes a **highly-significant impact** in the field. Specific guidance on this is on the next page.
- Articles should be scientifically sound with the conclusions supported by the experimental data.
- Please comment on the **novelty and likely impact** of the work. **Routine and incremental work** should generally **not** be recommended for publication in OBC; even if competently researched and reported. If you rate the article as 'routine' yet recommend acceptance, please give specific reasons for this in your report.
- OBC has recently updated its [characterisation guidelines](#).

Best wishes,

Richard Kelly  
Editor, *Organic & Biomolecular Chemistry*

\* ISI Citation information 2014

## \*New\* Scope and Standards

OBC's scope includes total synthesis, synthetic methodology and bioorganic chemistry as well as organic and bioorganic aspects supramolecular and macromolecular chemistry, theoretical and computational chemistry, catalysis, medicinal chemistry and natural products, including biosynthetic aspects.

Research articles published in OBC must show a significant advance on previously published work, or bring new thinking or results which will potentially have a strong impact in their field. Specific guidance for some areas of our scope is given below:

**Organic synthesis** – we welcome research in all areas of organic synthesis, including studies on small organic molecules and biomolecules, and articles which report purely synthetic work without biological data. Total or multistep syntheses should report new or improved strategies or methods, or a more efficient route to the target compound. Methodology studies should show a significant improvement on known methods. Research which extends known methodology to a different class of compounds is generally not suitable unless that class is significantly different to those on which that methodology has been used previously.

**Medicinal chemistry** – we welcome studies which report significant synthetic or bioorganic research which is directed towards medicinal chemistry applications. Studies which show routine syntheses accompanied by biological testing are generally not suitable for OBC.

**Theoretical and computational studies** – we welcome studies that report new models of reactivity, selectivity, bonding or structure, or new computational methods, that have relevance for the design of subsequent experiments (and that relevance should be clearly justified in the paper). Relevance is perhaps most clearly demonstrated by the description of testable predictions derived from the results of the reported theoretical work; the tests of these predictions could be contained in the same paper in which the predictions are described. Computational research that merely reproduces experimental data is not suitable for OBC

**Natural products** – we welcome articles that report new and interesting syntheses of natural products (see Organic Synthesis guidelines above). Isolation or identification studies are welcome when the compound being reported:

- 1) Has a novel structural class with unreported carbon skeleton, unusual functional groups or unusual modifications and/or
- 2) Displays a potent or unexpected biological activity or an unexpected mechanism of action.

Routine isolation studies are not suitable for OBC.

**Sensors** - we welcome articles describing sensors for ions and/or molecules provided that (a) they address targets and situations of practical relevance, and (b) they represent significant and demonstrable improvements on previous methodology. In particular, sensors for species in artificial surroundings (e.g. hydrophilic ions in organic solvents) will not usually be acceptable for publication.



## Organic and Biomolecular Chemistry

## ARTICLE

## Multicomponent, Fragment-Based, Synthesis of New Natural-Based Polyphenols and their Inhibiting Activity on Beta-Amyloid Oligomerization

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Chiara Lambruschini,<sup>a†</sup> Denise Galante,<sup>b†</sup> Lisa Moni,<sup>a†</sup> Francesco Ferraro,<sup>a</sup> Giulio Gancia,<sup>b</sup> Renata Riva,<sup>a</sup> Alessia Traverso,<sup>a</sup> Luca Banfi<sup>a\*</sup> and Cristina D'Arrigo<sup>b\*</sup>

A new and concise fragment-based approach towards artificial (but "natural-based") complex polyphenols has been developed, exploiting the Ugi multicomponent reaction of phenol-containing simple substrates. The resulting library of compounds has been tested for the capacity to inhibit  $\beta$ -amyloid protein aggregation, as a possible strategy to develop new chemical entities to be used as a prevention or a therapy for Alzheimer's disease. Some of the members of the library have demonstrated, in Thioflavin assays, a highly promising activity in inhibiting aggregation for two  $\beta$ -amyloid peptides: A $\beta$ 1-42 and the truncated A $\beta$ pE3-42.

### Introduction

Alzheimer's Disease (AD) is the most prevalent neurodegenerative disorder. The hallmarks of AD are the extracellular plaques, derived from aggregation of  $\beta$ -amyloid peptides, and neurofibrillary tangles composed by hyperphosphorylated protein tau. Inhibition of  $\beta$ -amyloid protein aggregation represents one of the most promising targets in the development of pharmacological treatments for the prevention of Alzheimer's disease.<sup>1-3</sup> Moreover, substances that strongly bind to  $\beta$ -amyloid proteins may be very useful diagnostic tools for an early detection of this disease.<sup>4</sup> Among the various substances that have been found to bind to  $\beta$ -amyloids, natural polyphenols have emerged as a particularly promising class, being able to inhibit  $\beta$ -amyloid aggregation and disrupt preformed amyloid fibrils.<sup>5-11</sup> Hydrogen bonding, hydrophobic interactions, and aromatic stacking are suggested to be the driving forces of the anti-amyloidogenic role of polyphenols. In addition, antioxidant activity may also be involved in the anti-amyloidogenic role.<sup>12</sup> Figure 1 depicts some of the most active natural polyphenols.

However, these natural compounds have often poor pharmacokinetic properties. For example, pharmacokinetic results for curcumin and its metabolites suggested limited or very poor bioavailability; in particular, curcumin was present in




Figure 1 Some natural polyphenols with  $\beta$ -amyloid anti-aggregation properties.

very little amount in the cerebrospinal fluid.<sup>13</sup> In addition, several natural polyphenols contain a catechol or a pyrogallol type ring, making them highly susceptible to oxidation, thus strongly reducing their half-life in the body.<sup>12, 14</sup> Finally, chemical modifications of complex natural polyphenols is quite tricky,<sup>15</sup> hampering the systematic synthesis of analogues that

<sup>a</sup> Department of Chemistry and Industrial Chemistry, University of Genova, via Dodecaneso 31 - 16146 Genova, Italy. E-mail: banfi@chimica.unige.it.

<sup>b</sup> Istituto per lo Studio delle Macromolecole, Consiglio Nazionale delle Ricerche, via De Marini 6, 16149 Genova, Italy. E-mail: cristina.darrigo@ge.ismac.cnr.it.

<sup>c</sup> Address here.

† Contributed equally to this work

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x



## ARTICLE

## Organic and Biomolecular Chemistry

might overcome the above quoted limitations and/or be endowed of higher potency.

Therefore, we reasoned that a fragment-based synthesis of natural-derived polyphenols, obtained by joining simple, monocyclic, phenol containing, building blocks, would be a very useful tool to assembly a huge number of molecular entities, allowing: a) optimization of pharmacodynamic properties; b) optimization of pharmacokinetic properties; c) the synthesis of structures that include pharmacophores directed towards alternative AD-related targets, with the aim to develop drugs able to simultaneously interact with different targets (multi-target strategy).<sup>15</sup> If the synthetic sequence is smartly designed, in order to be quite short, and simple building blocks derived from renewable sources are exploited, the final optimized compounds could be easily accessible in an eco-friendly manner, thus making their potential use as nutraceuticals definitely feasible.

Multicomponent reactions have emerged in the last 20 years as a powerful tool in drug discovery. They are intrinsically endowed with very high step economy and operational simplicity and are thus perfectly suited for a rapid generation of libraries characterized by several diversity inputs. Among them, the isocyanide-based Ugi reaction (U-MCR)<sup>16</sup> is particularly useful, since it allows the simultaneous joining of 4 diversity inputs, represented by easily accessible compounds, as isocyanides, aldehydes, primary amines and carboxylic acids. Thus, also taking advantage of our previous experience both in the Ugi reaction,<sup>17-20</sup> and in the assembly of natural-based polyphenols,<sup>21</sup> we selected U-MCR at the key step in our fragment-based approach. The classical scaffold obtained by the Ugi reaction is a peptidomimetic structure. This is another added value of our approach, since also peptidomimetics are widely studied as potential inhibitors of  $\beta$ -amyloid aggregation.<sup>22</sup>


Using this synthetic methodology we were able to prepare a series of complex polyphenols containing 2 to 4 hydroxy-substituted aryl groups, most of them derived from renewable sources, and test them for their ability to inhibit *in vitro*  $\beta$ -amyloid aggregation. In this paper we report our preliminary observations, representing a first "proof of concept" for our approach.

## Results and discussion

### Synthesis

As depicted in Scheme 1, the Ugi reaction allows us to prepare a peptidomimetic structure **1** with 4 appendages. In our plan, from two to four of these appendages should contain a phenolic aryl group, tethered to the main scaffold through linkers of different lengths. In principle, our target compounds **1** could be accessed in just one step by employing, in the U-MCR, components containing the free phenols. However, preliminary investigation has shown that the presence of free phenols had a negative effect on the yield and cleanliness of the multicomponent reaction. Thus we shifted to a slightly

longer (2 steps) sequence, employing suitably protected building blocks.



Scheme 1 General synthetic strategy


Many efforts were devoted to the selection of the best protecting group. Initially, because of the high atom economy and the easiness of final deblocking, we opted for a simple acetyl group.<sup>21</sup> As shown in Scheme 2, a first Ugi reaction with only two phenol containing components (aniline **3**<sup>23, 24</sup> and aldehyde **4**<sup>25</sup>) worked well, affording compound **2a** in good yields. It was then smoothly deprotected by saponification to the diphenol **1a**. However, the use of the acetyl as protecting group was soon demonstrated to be far from general. For example, simply using the protected phenol containing isocyanide **5**, we failed to isolate any of the expected product **6**. Similarly, all attempted Ugi reaction using diacetyl protected caffeic acid **7** were unsuccessful. We think that the aryl acetates are somehow unstable under the Ugi reaction and that the liberated phenols and acetic acid promote unwanted side reactions. Thus, only with protected hydroxyanilines and reactive isocyanide/carboxylic acids the reaction turned out to be feasible, strongly limiting diversity exploration.

Looking for a more stable protection we shifted to the dimethyl-*tert*-butylsilyl group (TBDMS). In this case the group proved to be fully stable under the Ugi conditions. However, the reactions tend to be rather sluggish. An improvement can be obtained by performing the imine treating the aldehyde and the amine in CH<sub>2</sub>Cl<sub>2</sub> in the presence of dry MgSO<sub>4</sub>. Anyway, the isolated yield, in the case of compound **11**, was only 25%. Furthermore, the presence of more than two TBDMS groups renders the Ugi products rather insoluble in most solvents making their purification, as well as the assessment of purity by NMR or HPLC, troublesome. In particular, at NMR, broad signals due to slowly converting conformers are observed. Therefore, because of the poor atom economy, the slow reaction kinetics and the unsatisfactory chemico-physical properties, we decided to abandon this protecting group as well.

Our attention was then drawn by the allyl group for its high atom economy (similar to the acetyl), its expected stability

under the Ugi conditions, and the possibility to remove it under neutral conditions, thanks to palladium (0) catalysis. The Ugi reaction of substrates containing this group was thoroughly optimized. Scheme 3 shows a representative example. In particular we noticed that benzaldehydes bearing an allyloxy group in *para* position resulted less reactive (because of the electron-donating properties of allyloxy) than 4-acetoxybenzaldehyde **4** or benzaldehyde and that

degradation. Moreover, we lost some material during the work-up of the deprotection step. To avoid these problems, we decided to peracetylate the crude polyphenol, purify it by chromatography, and finally remove the acetyl groups. In this way, the crude polyphenols obtained after deacetylation, acid resin treatment, and filtration, were pure enough (HPLC and  $^1\text{H}$  NMR control) to be used as such for biophysical tests, without the need to extractive or chromatographic purifications that



Scheme 2 Synthesis of the first polyphenols using the Ac or TBDMS protecting groups.

substituted cinnamic acids such as protected ferulic and caffeic acids also brought about a slower kinetic. We found that the best solvent, in order to have reasonable reaction times and avoid the formation of Passerini side-products, was a 1:1 mixture of EtOH and trifluoroethanol. Moreover, it was advantageous to perform the imine in this solvent for 5 h, before adding the other reagents. The reactions typically last 48-72 h.

As for the deblocking step, we tried various Pd catalysts and scavengers. Eventually, the reaction was found to be more reproducible using a palladium (II) precursor, Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>, than with Pd(PPh<sub>3</sub>)<sub>4</sub>. As scavenger, ammonium formate was the best, allowing an easy removal of its excess by a simple extraction under neutral conditions. In the case of compound **1c**, which was initially used as model, chromatographic purification, followed by treatment with active coal, worked fine, affording a very pure product (procedure A). However, in other cases, especially when a catechol or a pyrogallol moiety were present, direct chromatographic purification of the final polyphenol was not fully satisfactory, due to partial

may be troublesome, because of the polyphenol polarity and/or for the possible partial degradation. This procedure (procedure B) is exemplified in Scheme 3 for the synthesis of compound **1d**. Apart from some products synthesized in the initial part of this research, we later routinely used procedure B.

Scheme 4 depicts all the polyphenols **1a-q** prepared so far, whereas Table 1 reports the procedure used in each case and the yields of various steps (except for compounds **1a,b** prepared as described in Scheme 2). When procedure B was used, full characterization was carried out only on the acetylated compounds **2**. As already stated above, the final phenols derived from deacetylation were pure enough for testing, as checked by  $^1\text{H}$  NMR and HPLC (HPLC purity  $\geq 96\%$  in nearly all cases, except for **1d**, **1l** and **1m** (92%).

Concerning the Ugi reactions, we found out that its efficiency depends on the nature of components used. For example, it worked poorer using substituted anilines than with benzylamines or other aliphatic amines. Also protected ferulic and caffeic acids were somehow less reactive than simple acids


## ARTICLE

## Organic and Biomolecular Chemistry

like propionic acid or benzoic acids. Allyl protected substituted benzaldehydes (*p*-hydroxybenzaldehyde or vanillin) were less reactive than benzaldehyde, probably because of the electron-donating properties of the allyloxy group in *para* position. Finally, aliphatic isocyanides behaved better than the aromatic ones. In particular, during the synthesis of compound **1k**, the combination of 4-allyloxybenzaldehyde, a bulky aromatic isocyanide and an aniline led to a poor yield in the Ugi reaction

(18%). In this case, we obtained a moderate improvement using 4-pivaloyloxybenzaldehyde (31%). We then used procedure B and isolated and characterized the mixed pivaloyl-acetyl compound **2k**.

Apart from the polyphenols listed in Scheme 4, other compounds targeted by us were not obtained in reasonable yields.



**Scheme 3** Representative procedures for the preparation of polyphenols **1** via allyl ethers.

**Table 1** Yields of polyphenol synthesis using the allyl protecting group

Entry	Polyphenol	Yield of <b>14</b>	Procedure used <sup>a</sup>	Yield of <b>2</b>	Yield of <b>1</b>
1	<b>1c</b>	69%	A	- <sup>b</sup>	80%
2	<b>1d</b>	57% (91%) <sup>c</sup>	B	76%	- <sup>c</sup>
3	<b>1e</b>	72%	A	- <sup>b</sup>	59%
4	<b>1f</b>	73%	B	83%	- <sup>d</sup>
5	<b>1g</b>	28%	B	59%	- <sup>d</sup>
6	<b>1h</b>	22%	A	- <sup>b</sup>	74%
7	<b>1i</b>	33%	A	- <sup>b</sup>	68%
8	<b>1j</b>	17%	B	64%	- <sup>d</sup>
9	<b>1k</b>	31% <sup>e</sup>	B	69%	- <sup>d</sup>
10	<b>1l</b>	75%	B	58%	- <sup>d</sup>
11	<b>1m</b>	59%	B	78%	- <sup>d</sup>
12	<b>1n</b>	63%	B	69%	- <sup>d</sup>
13	<b>1o</b>	69%	B	74%	- <sup>d</sup>
14	<b>1p</b>	71%	B	60%	- <sup>d</sup>
15	<b>1q</b>	72%	B	75%	- <sup>d</sup>

<sup>a</sup> See Scheme 3 and Experimental part. <sup>b</sup> In procedure A, **14** was directly converted to **1**. <sup>c</sup> In brackets the yield calculated taking into account the recovered aldehyde. <sup>d</sup> In procedure B, **2** was converted quantitatively into **1** by hydrolysis of acetates. <sup>e</sup> In this case 4-pivaloyloxybenzaldehyde was used, and Ugi product was not **14a**, but **15** (Scheme 4).




## Organic and Biomolecular Chemistry

### ARTICLE

For example, although triallylated gallic acid was a good substrate for the Ugi reaction, all attempts to deprotect it without extensive decomposition were unsuccessful. Similarly, when we saturated the double bond in caffeic acid derived polyacetate **2f**, deblocking of the acetyl group led to decomposition of the final product as well. We attribute this behaviour to the presence of a catechol or pyrogallol moiety,

which are prone to oxidation under basic conditions. In the case of caffeic acid adducts, the conjugation with the unsaturated amides makes the catechol less electron-rich and thus more stable to oxidation, but when the double bond is hydrogenated the catechol becomes too reactive.

**Biochemical and biophysical assays**




Scheme 4 Polyphenols prepared and their precursors.

The first biophysical analysis performed on the new polyphenols was the solubility in aqueous solution because the working condition is Phosphate Buffer Solution (PBS) at pH 7.4 to mimic the physiological environment. All compounds showed complete solubility at working concentration (25  $\mu$ M) in PBS containing 1% of DMSO (solubility-related data are reported in the S.I.). This percentage of DMSO does not damage cells and animals for future biological tests, and also

does not alter the aggregation of  $\beta$ -amyloids. In any case control blank experiments with samples containing 1% DMSO in the buffer were always performed in parallel.

To investigate their ability to inhibit the amyloid aggregation, kinetics assays monitored by thioflavin-T were used to follow the formation of  $\beta$ -sheet rich structures, visualized then by transmission electron microscopy (TEM).<sup>26, 27</sup>



**Figure 2** ThT Fluorescence in percentage respect to the control sample, after 24 h of aggregation at 37 °C, the concentration was 5  $\mu$ M for  $\beta$ -amyloids and 25  $\mu$ M for polyphenols in PBS + 1% DMSO.

We explored the interaction of the new complex polyphenols with two particular  $\beta$ -peptides, A $\beta$ 1-42 and A $\beta$ pE3-42. The full-length A $\beta$ 1-42 is one of the most abundantly identified in the brain deposits (together with A $\beta$ 1-40 and N-terminal truncated A $\beta$  peptides). A $\beta$ pE3-42 is a peptide N-terminal truncated at residue 3 (Glu) and further modified by cyclization of Glu (E) to pyroglutamic acid (pE). These structural modifications increase A $\beta$ pE3-42 aggregation propensity, its resistance to degradation of proteases, and display an enhanced cytotoxicity in comparison to A $\beta$ 1-42<sup>28, 29</sup> as well as the ability to unfold the full-length into toxic aggregates.<sup>30</sup>

Before aggregation kinetics experiments, we used a natural polyphenol as a positive control to compare the efficacy of the new synthetic polyphenols. From the literature,<sup>31</sup> epigallocatechin gallate (EGCG) was the most potent natural polyphenol in inhibiting (*in vitro*) aggregation of  $\beta$ -amyloid proteins, although it was also reported that *in vivo* has a poor stability.<sup>12</sup> After investigating the lowest concentration at which EGCG strongly inhibited amyloid aggregation, we chose to work at 25  $\mu$ M and to use this concentration also for our synthetic polyphenols.

We explored the interaction of the new complex polyphenols with A $\beta$ 1-42 and A $\beta$ pE3-42 to verify their ability to inhibit  $\beta$ -amyloid aggregation.  $\beta$ -sheet content and aggregation process

at 37 °C in PBS (150 mM, pH 7.4) and 1% DMSO were followed by fluorescence using Thioflavin T (ThT), a probe that detects the presence of  $\beta$ -sheets in the sample. Indeed, the aggregation of  $\beta$ -amyloids starts when they change their secondary structure from  $\alpha$ -helix (in the membrane environment) or coil (in basic environment) to  $\beta$ -sheet conformation.<sup>32</sup> Then the aggregation proceeds forming small aggregates, called oligomers, until reaching larger aggregates such as long fibrils, which will later precipitate.


We used as reference A $\beta$ 1-42 and A $\beta$ pE3-42 alone under the same conditions of the experiments carried out in presence of the new polyphenols. In Figure 2 it is reported, for all samples (except **1a** and **1b**), the ThT emission value after 24 h of aggregation at 37 °C. In the case of **1a** and **1b**, the first polyphenols prepared by us, we carried out the test on A $\beta$ 1-42 alone and at higher concentrations. Only **1b** showed moderate activity and we did not repeat the experiments at 25  $\mu$ M. The relative data are reported in the S.I.

As we can see in panel A, all new polyphenols have some effect on A $\beta$ 1-42 aggregation. **1m**, **1f** and especially **1j** even increase the  $\beta$ -sheets content of the peptide, so with these compounds also the fibrils formation grows. On the contrary, the best new polyphenol to inhibit the fibrillation process is **1c**. The degree of inhibition is also quite high for **1h**, **1i**, **1l** and **1n**, and moderate for **1d**, **1e**, **1k**, **1o**, **1p** and **1q**. However, with **1c** also the kinetic of aggregation slows down (see Figure 3).

For A $\beta$ pE3-42 (Figure 2, panel B) the results were different, the best inhibitor of the aggregation process being indeed **1f**, but also **1e**, **1i** and **1n** showed a good effect. Moderate inhibition was visible for **1l** and **1q**. In both cases, epigallocatechin gallate (EGCG), the most active natural polyphenol, was still more effective in preventing amyloid aggregation at the same concentration. It is however worth noting that the presence of two pyrogallol moieties in EGCG make its stability under physiological conditions troublesome, whereas we have obtained a similar, albeit somehow lower, activity with much more stable polyphenols derived from ferulic acid<sup>33</sup> (**1c**, **1i**, **1n**), which are much more stable to oxidation and more promising from the pharmacokinetic point of view. Is interesting to note that for A $\beta$ 1-42 there are different polyphenols with a good anti-aggregating effect (Inhibition>30%), while for A $\beta$ pE3-42, which is more prone to aggregation, more resistant to degradation and more toxic in comparison to A $\beta$ 1-42, polyphenols with a good anti-aggregating effect are fewer.

To better understand the different inhibition mechanism of the most active molecules, we determined also the aggregation kinetics curve of ThT Fluorescence emission over time for A $\beta$ 1-42 and A $\beta$ pE3-42 in presence of **1c** and **1f** (Figure 3). In the case of A $\beta$ 1-42, **1c** slows down the fibril growth phase and reduces the amount of fibrils, while **1f** slows down only the lag phase, but the amount of fibrils is similar to that of A $\beta$ 1-42 alone as reported by the plateau value.






**Figure 3** Kinetics of aggregation monitored by ThT Fluorescence emission. A) Kinetics curve for A $\beta$ 1-42; B) Kinetics curve for A $\beta$ pE3-42. The concentration was 5  $\mu$ M for A $\beta$  peptides and 25  $\mu$ M for polyphenols in PBS + 1% DMSO, in yellow are reported the standard error for the curves of A $\beta$  alone.

On the contrary, in the case of A $\beta$ pE3-42, **1c** has almost no effect on the aggregation inhibition, while **1f** inhibits the maximum A $\beta$  assembly. A $\beta$ pE3-42 is very fast to aggregate in the initial stage: in fact, in our conditions, it is never possible to see a lag phase for this peptide. The aggregation pathway is different from that of the full-length peptide and results in the enhancement of the seed production that speeds the aggregation into more fragmented and less structured species. We think that **1c** and **1f** act at different levels during aggregation, inhibiting the formation of different structural species. **1f** is able to inhibit the formation of oligomers that work as seed for the aggregation. In fact, it extends the lag phase in A $\beta$ 1-42 but does not inhibit the fibril formation, whereas is able to strongly reduce the A $\beta$ pE3-42 aggregation. So **1f** is effective on A $\beta$ pE3-42 because it inhibits the first phase of aggregation, the one forming the oligomers.

As shown in Figure 4, **1c** is even capable to disrupt fibrils, once they have formed. So, addition of **1c** after 24 h (when the aggregation process in the absence of inhibitors is already complete) provokes a significant decrease of ThT fluorescence emission.

To confirm these data, we studied the morphology of A $\beta$  peptides aggregates in presence of those polyphenols that behaved best from ThT test. They were observed after 24 h of incubation at 37°C at the same ratio of the fluorescence experiments (Figure 5). In Figure 5, panel A, the morphology of A $\beta$ 1-42 is reported, showing typical amyloid fibrils that appear as very entangled fibril bundles and also striated ribbons are visible. When **1c** is added to A $\beta$ 1-42 (panel B), fibrils bundles decrease and the thinner fibres appear less entangled. Moreover, the fibrils are somehow fragmented, due to the twist change along the fibre. When A $\beta$ 1-42 and **1f** are mixed (panel C), fibril bundles and little spheroidal aggregates appear even if the fibrils have smaller diameters than A $\beta$ 1-42 alone. The quantity of  $\beta$ -sheet in this sample is not different from that of full-length alone (as showed in the ThT assay). This

alternative morphology depends on amyloid multi-step assembly pathways that is altered from the slowdown of the lag phase by **1f**. Looking at panel D, the morphology of A $\beta$ pE3-42 is shown with few bundles and short fibrils. The addition of **1c** (panel E), results in the decrease in the amount of fibrils but the morphology is very similar to that of the A $\beta$ pE3-42 alone. As a matter of fact this polyphenol is not very effective in inhibiting the assembly of the pyroglutamate  $\beta$ -amyloid. Finally, in the presence of **1f** (panel F), in agreement with the ThT assay, the number of fibrils and their length strongly decrease and many dispersed small spheroidal aggregates appears. Moreover, globular aggregates attached along the fibrils are visible. Also the morphology confirms the structural changes induced by the addition of the best new polyphenols. Several studies have indicated that hydrophobic forces,



**Figure 4** Effect of addition of **1c** to aggregated fibrils (A $\beta$ 1-42, monitored by ThT Fluorescence emission).

aromatic stacking, and electrostatic interactions stabilize the A $\beta$  structure.<sup>34</sup> It was found that short fragments of A $\beta$  (QKLVFF) self-assemble and also bind specifically to full-length




Figure 5 Morphology of the species by TEM. A) A $\beta$ 1-42 alone; B) A $\beta$ 1-42 + 1c; C) A $\beta$ 1-42 + 1f; D) A $\beta$ pE3-42 alone; E) A $\beta$ pE3-42 + 1c; F) A $\beta$ pE3-42 + 1f

peptides, supporting the hypothesis that  $\pi$ - $\pi$  interactions may play a central role in the molecular recognition and A $\beta$  self-assembly process.<sup>35</sup> In the last years, various approaches to inhibit and reverse misfolding and aggregation of  $\beta$ -amyloid have been applied. One of these was the creation of short synthetic peptides capable of binding A $\beta$  but unable to become part of a  $\beta$ -sheet structure ( $\beta$ -sheet breaker peptides) that destabilize the amyloidogenic A $\beta$  conformer and hence preclude amyloid formation.<sup>36</sup> These  $\beta$ -sheet breaker peptides act by the binding of the central hydrophobic region of A $\beta$  protein (amino acids 17-21: LVFFA). Another approach was the use of polyphenolic compounds as  $\beta$ -sheet inhibitors. The possible mechanisms by which polyphenols destabilize  $\beta$ -amyloid aggregation still remain unclear, but several mechanisms have been proposed to date and structural similarities between various highly efficient inhibitors have been identified. It has become evident that the presence of phenolic rings with a few linkers and at least two hydroxy groups could favour effective non-covalent interactions with the fibril  $\beta$ -sheet structures and interfere with their elongation and/or assembly.<sup>33</sup> Both the number of hydroxy groups and the positioning of these groups on the polyphenolic structure is important, however there is no clear understanding of the link between phenol positional substitution and corresponding anti-aggregation activity. Moreover, the planarity of the inhibitor is essential for increasing surface contact with A $\beta$  peptides.<sup>5</sup>

All of our polyphenols have a peptidomimetic structure, more than two aromatic rings essential for  $\pi$ - $\pi$  stacking interactions

with hydrophobic amino acid residues of A $\beta$  and at least two hydroxyl groups to form hydrogen bonds with hydrophilic amino acid residues of A $\beta$ . The resonance structure of polyphenols provides enough planarity to penetrate the A $\beta$  fibril hydrophobic groove, thus disturbing the fibril structure.<sup>37</sup> On the basis of the results collected till now, we can try to correlate the observed activity with the various pharmacophores. As far as it concerns the groups derived from the isocyanide, we did not observe, for both peptides, a particular influence of the structure. It is noteworthy that the most active compounds so far have a *tert*-butyl group as the isocyanide derived one, suggesting that the presence of an aromatic ring in this position is not essential.

On the contrary, the structure of the residues derived from the carboxylic acid and the amine seems more important. Regarding the first one, best results have been obtained with cinnamic acid derivatives (caffeic and ferulic acid). For the pyroglutamate  $\beta$ -amyloid, we noted that polyphenols synthesized from caffeic acid (**1e** and **1f**) are able to strongly inhibit aggregation, whereas among polyphenols derived from ferulic acid, only **1i** and **1n** show good activity. On the other hand, for the full-length peptide we noticed in most cases (except for **1m**) a good inhibitory effect when the starting carboxylic acid is ferulic acid. The effect is good for **1c**, **1i**, **1l**, **1n** and moderate for **1d**, **1o**, **1p** and **1q**. On the other hand, polyphenols where carboxylic acid is a benzoic acid (**1g**, **1j** and **1k**) have no or little inhibitory effect on both peptides. A propionyl group (**1h**) resulted in no effect on the truncated peptide, but in a moderate activity on the full-length one. We



## ARTICLE

## Organic and Biomolecular Chemistry

can conclude that ferulic acid is best for A $\beta$ 1-42, whereas as caffeic acid is the best for A $\beta$ pE3-42, although, as shown by **1i** and **1n**, also ferulic acid derivatives may inhibit this peptide.

The nature of the group derived from the amine component in the Ugi seems important for both peptides. Best results have been obtained with benzylamines or anilines, whereas a drop of activity was observed in the case of **1m**, having just one more carbon atom, or for **1d**, where the benzyl/aryl group is replaced by a simple butyl. The benzyl groups seem better than the aryl ones (the most promising compounds, **1c** and **1f** have indeed a simple benzyl group), although it is remarkable that 4-hydroxyphenyl containing **1i** is more active than **1c** for the truncated peptide.

Finally, the group derived from the aldehyde has been so far less explored by us. However, in particular for the pyroglutamate  $\beta$ -amyloid peptide, we have noticed a remarkable influence of an additional methoxy group (compare compound **1c**, containing the 4-hydroxyphenyl group, and **1n**, containing the 4-hydroxy-3-methoxyphenyl group).

It is interesting to note that only **1n** and **1i** are able to inhibit the aggregation of both A $\beta$ 1-42 and A $\beta$ pE3-42. This great variation of substituent effects on the two  $\beta$ -amyloid peptides likely depends on their intrinsic differences. Probably our polyphenols bind in different region of the chain by interacting with diverse residues and/or at distinct levels in the assembly mechanism that brings to the aggregation.

## Conclusions

To the best of our knowledge, this paper represents one of the first reports, to our knowledge, on the combinatorial synthesis of complex artificial (but "natural-based") polyphenols using a fragment-based approach and on the demonstration that some of these compounds are indeed able to inhibit or even disrupt  $\beta$ -amyloid aggregation. In fact, we tested the anti-aggregation activity on two different  $\beta$ -amyloid peptides (A $\beta$ 1-42 and A $\beta$ pE3-42), normally present in AD brains, that have a different assembly pathway. For this reason, some polyphenols are more prone to inhibit the aggregation process of A $\beta$ 1-42 than that of A $\beta$ pE3-42 and vice versa. This approach could allow the formulation of mixtures of active polyphenols to inhibit simultaneously the aggregation of both peptides and avoid the formation of more neurotoxic co-aggregates.

Clearly more insight into the mechanism by which our systems inhibit  $\beta$ -amyloid protein aggregation is needed, for example by using NMR spectroscopy or computational models. However, notwithstanding the still limited number of molecules tested, the results depicted in Figure 3 indicates that subtle variation in the structure of the appendages may have a strong impact on activity. Thus, the smart synthetic approach (based on Ugi MCR), that allows to assemble these polyphenols in 2 steps, by varying up to 4 diversity inputs, will strongly facilitate the fine tuning of the pharmacophores in order to increase potency and/or selectively target different sub-species of  $\beta$ -amyloid proteins. The incorporation of fragments with known anti-oxidant activity, such as ferulic

acid, may have other kind of beneficial effects on AD patients, as pointed out in a recent paper.<sup>38</sup> Compared to the most active natural compounds (e.g. epigallocatechin gallate), our systems are expected to be metabolically much more stable (especially those, like **1c**, not containing a catechol system), thus overcoming the main drawback of some natural polyphenols and making them better suited for *in vivo* experiments, that will soon be carried out.

## Experimental

NMR spectra were taken at r.t. in CDCl<sub>3</sub> or in d<sub>6</sub>-DMSO at 300 MHz (<sup>1</sup>H), and 75 MHz (<sup>13</sup>C), using, as internal standard, TMS (<sup>1</sup>H NMR in CDCl<sub>3</sub>; 0.000 ppm) or the central peak of DMSO (<sup>1</sup>H NMR in d<sub>6</sub>-DMSO; 2.506 ppm) or the central peak of CDCl<sub>3</sub> (<sup>13</sup>C in CDCl<sub>3</sub>; 77.02 ppm), or the central peak of DMSO (<sup>13</sup>C in d<sub>6</sub>-DMSO; 39.43 ppm). Chemical shifts are reported in ppm ( $\delta$  scale). Peak assignments were made with the aid of gCOSY and gHSQC experiments. In ABX system, the proton A is considered upfield and B downfield. [ $\alpha$ ]<sub>D</sub> values are given in 10<sup>-1</sup> deg cm<sup>2</sup> g<sup>-1</sup>. IR spectra were recorded as solid, oil, or foamy samples, with the ATR (attenuated total reflectance) technique. TLC analyses were carried out on silica gel plates and viewed at UV ( $\lambda$ =254 nm or 360 nm) and developed with Hanessian stain (dipping into a solution of (NH<sub>4</sub>)<sub>4</sub>MoO<sub>4</sub>·4H<sub>2</sub>O (21 g) and Ce(SO<sub>4</sub>)<sub>2</sub>·4H<sub>2</sub>O (1 g) in H<sub>2</sub>SO<sub>4</sub> (31 mL) and H<sub>2</sub>O (469 mL) and warming). R<sub>f</sub> values were measured after an elution of 7–9 cm. GC-MS analysis were recorded on HP-5890 series II HEWLETT PACKARD equipped with a HP-1 column (12 m,  $\phi$  = 0.2 mm) using He as carrier gas. MS were recorded on an electronic impact (EI, 70 eV) HP-5971A detector. Chromatography condition: flow 1.0 mL/min, injector temperature 250 °C, method 1 (initial temperature 100 °C, initial time 2 min, rate 20 °C/min, final temperature 290 °C); method 2 (initial temperature 70 °C, initial time 2 min, rate 20 °C/min, final temperature 260 °C). The data are reported as follow: retention time (Rt, min), m/z values and the abundance relative. Only m/z > 5 are reported. HRMS: samples were analysed with a Synapt G2 QToF mass spectrometer. MS signals were acquired from 50 to 1200 m/z in either ESI positive or negative ionization mode. Column chromatography was done with the "flash" methodology by using 220–400 mesh silica. Petroleum ether (40–60 °C) is abbreviated as PE. All reactions employing dry solvents were carried out under nitrogen. Extractions were always repeated three times and organic extracts were always dried over Na<sub>2</sub>SO<sub>4</sub> and filtered before evaporation to dryness.

Compounds **3**,<sup>23, 24</sup> **4**,<sup>25</sup> **5**,<sup>39</sup> **10**,<sup>40</sup> **12**,<sup>41, 42</sup> 4-allyloxybenzoic acid,<sup>43</sup> 4-allyloxy-3-methoxybenzaldehyde,<sup>44</sup> E-3,4-bis(allyloxy)phenylpropenoic acid,<sup>45</sup> and 4-pivaloyloxybenzaldehyde<sup>46</sup> were prepared by the reported methods.

**4-((tert-Butyldimethylsilyl)oxy)aniline 9.** A solution of *p*-aminophenol (1.00 g, 9.16 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (30 mL) was treated with imidazole (1.25 g, 18.3 mmol) and *tert*-butyldimethylsilyl chloride (2.07 g, 13.7 mmol). After stirring for 3 h at r.t. a purple suspension was obtained. It was treated

with saturated aqueous NaHCO<sub>3</sub> and extracted with CH<sub>2</sub>Cl<sub>2</sub>. After evaporation and chromatography (PE / AcOEt 8:2), pure **9** (1.880 g, 92%) was obtained as a colorless liquid. The spectroscopic and analytical data were in agreement with those reported.<sup>47</sup>

**4-((tert-Butyldimethylsilyloxy)phenyl isocyanide 8.** A solution of 4-((tert-butyldimethylsilyloxy)aniline **9** (1.00 g, 4.48 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (45 mL) was treated with formic acid (203 μL, 5.38 mmol), 4-dimethylaminopyridine (DMAP) (101 mg, 0.90 mmol) and, finally, dicyclohexylcarbodiimide (DCC) (1.017 g, 4.93 mmol). After 3.5 h at r.t., the resulting suspension was treated with additional formic acid (51 μL, 1.34 mmol) and DCC (185 mg, 0.90 mmol). After further 1.5 h, the suspension was filtered through a celite cake washing with CH<sub>2</sub>Cl<sub>2</sub>. After evaporation and chromatography (CH<sub>2</sub>Cl<sub>2</sub> / acetone 95:5), *N*-(4-((tert-butyldimethylsilyloxy)phenyl)formamide<sup>47</sup> was obtained (83%). 172 mg (0.65 mmol) of this formamide was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (2.5 mL), cooled to -15 °C, and treated with Et<sub>3</sub>N (272 μL, 1.95 mmol) and trichloromethyl chloroformate (diphosgene) (46 μL, 0.39 mmol). The temperature was allowed to reach 0 °C during 1 h and the mixture further stirred for 30 min. Then, the reaction was quenched with saturated aqueous NaHCO<sub>3</sub> and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic extracts were washed with saturated brine, evaporated and chromatographed (PE / CH<sub>2</sub>Cl<sub>2</sub> 70:30) to give pure **8** as a pearlescent oil (152 mg, 95% from formamide). The spectroscopic and analytical data were in agreement with those reported.<sup>47</sup>

**(E)-3-(4-(allyloxy)-3-methoxyphenyl)acrylic acid 13.** A solution of *trans*-ferulic acid (6.00 g, 30.9 mmol) in dry MeCN (100 mL) was treated with K<sub>2</sub>CO<sub>3</sub> (10.3 g, 74.16 mmol) and allylbromide (8.8 mL, 102 mmol). After stirring for 18 h at 70 °C, the resulting suspension was filtered through a celite cake washing with MeCN. After evaporation, the obtained allyl (*E*)-3-(4-(allyloxy)-3-methoxyphenyl)acrylate was directly dissolved in MeOH (150 mL) and treated with 1 N KOH aqueous solution (62 mL, 61.8 mmol). After stirring for 24 h at 60 °C and evaporation to reduced volume, the crude was treated with 1 N NaOH aqueous solution and extracted with AcOEt. The aqueous phase was acidified with 12 N HCl (final pH = 3) and extracted with AcOEt. The organic extracts were washed with saturated brine and evaporated. The resulting crude was triturated with Et<sub>2</sub>O to give pure (*E*)-3-(4-(allyloxy)-3-methoxyphenyl)acrylic acid as white solid (6.65 g, 96%). The spectroscopic and analytical data were in agreement with those reported.<sup>45</sup>

**4-allyloxyaniline.** A solution of 4-nitrophenol (5.00 g, 35.9 mmol) in dry MeCN (70 mL) was treated with K<sub>2</sub>CO<sub>3</sub> (12.4 g, 89.7 mmol) and allylbromide (4.7 mL, 53.9 mmol). After stirring for 18 h at 70 °C, the resulting suspension was filtered through a celite cake washing with MeOH. After evaporation, the crude was treated with saturated aqueous NH<sub>4</sub>Cl and extracted with Et<sub>2</sub>O in order to completely remove the salts. The organic extracts were washed with saturated brine and evaporated. The obtained 1-(allyloxy)-4-nitrobenzene was directly dissolved in EtOH (70 mL) and treated with Fe powder (16.0 g, 287.2 mmol) and a solution of NH<sub>4</sub>Cl (7.68 g, 143.6

mmol) in deionized H<sub>2</sub>O (28 mL). After stirring for 18 h at 75 °C, the resulting black suspension was filtered through a celite cake washing with MeOH. After evaporation, the crude was treated with saturated aqueous NaHCO<sub>3</sub> and extracted with AcOEt. The organic extracts were washed with saturated brine and evaporated. The resulting crude 4-allyloxyaniline (light brown oil) was used in the next step without further purification. The spectroscopic and analytical data were in agreement with those reported.<sup>48</sup>

**4-allyloxyphenethylamine.** A solution of tyramine (1.50 g, 11.00 mmol) in dioxane / H<sub>2</sub>O (22 mL, 3:1) was treated at r.t. with triethylamine (1.53 mL, 11.00 mmol) and di-*tert*-butyl dicarbonate (2.40 g, 11.00 mmol). After 2 h the solution was concentrated under vacuum, and the residue was poured into a mixture of 5% aq (NH<sub>4</sub>)<sub>2</sub>PO<sub>4</sub> and 1 M HCl and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic extracts were washed with saturated brine and evaporated. The resulting *N*-Boc-tyramine was diluted in dry DMF (26 mL) and treated with Cs<sub>2</sub>CO<sub>3</sub> (4.70 g, 14.4 mmol) and allylbromide (1.3 mL, 14.1 mmol). After stirring for 4 h at 50 °C, the mixture was poured in saturated aqueous NH<sub>4</sub>Cl and extracted with Et<sub>2</sub>O. The organic extracts were washed with saturated brine and evaporated. Then, the crude was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (10 mL) at 0 °C and treated with trifluoroacetic acid (5 mL). After stirring for 3 h at r.t., the solution was evaporated to dryness, taken up with 1 M aqueous NaOH and extracted with CH<sub>2</sub>Cl<sub>2</sub> to give pure 4-allyloxyphenethylamine as yellow oil (1.72 g, 88% from tyramine). R<sub>f</sub> = 0.33 (CH<sub>2</sub>Cl<sub>2</sub> / MeOH 15:1 + 1% of Et<sub>3</sub>N). δ<sub>c</sub>(75 MHz, CDCl<sub>3</sub>, 25 °C): 156.8, 133.1, 131.4, 129.4 (x2), 117.1, 114.4 (x2), 68.5, 49.3, 43.1, 38.4. IR: ν<sub>max</sub>/cm<sup>-1</sup> 3373, 3029, 2926, 2857, 1715, 1648, 1610, 1582, 1509, 1457, 1424, 1382, 1362, 1297, 1237, 1221, 1177, 1154, 1111, 1069, 1021, 996, 924, 818, 752, 644, 617. GC-MS (method 1) t<sub>R</sub> 7.17 min, m/z (%) 148 ([M-CH<sub>2</sub>NH<sub>2</sub>]<sup>+</sup>, 7.4), 107 (17), 91 (7.2), 79 (6.4), 78 (7.4), 77 (13), 55 (7.6), 52 (7.5), 51 (9.7), 42 (5.0), 41 (100), 39 (35). The other spectroscopic and analytical data were in agreement with those reported.<sup>49</sup>

**3-allyloxybenzylamine.** A solution of 3-allyloxybenzyl alcohol<sup>50</sup> (2.66 g, 16.1 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (55 mL) was cooled at -15 °C and treated with Et<sub>3</sub>N (2.9 mL, 20.9 mmol) and mesyl chloride (1.5 mL, 19.3 mmol). After stirring for 4 h at -15 °C, the solvent was evaporated and the obtained mesylate was directly dissolved in dry DMF (23 mL) and treated NaN<sub>3</sub> (2.30 g, 33.8 mmol). After stirring for 3 days at r.t., the mixture was poured in H<sub>2</sub>O and extracted with Et<sub>2</sub>O. The organic extracts were washed with saturated brine (× 5) and evaporated. The crude was purified by chromatography (PE / CH<sub>2</sub>Cl<sub>2</sub> from 8:2 to 7:3) to give pure 3-allyloxybenzyl azide as pale yellow oil (2.63 g, 86% from 3-allyloxybenzyl alcohol). R<sub>f</sub> = 0.29 (PE / CH<sub>2</sub>Cl<sub>2</sub> 8:2). δ<sub>H</sub>(300 MHz, CDCl<sub>3</sub>, 25 °C): 7.29 (m) (1 H, m, ArCH); 6.93-6.86 (3 H, m, ArCH); 6.06 (1 H, ddt, J 10.5, 17.2, 5.3 (t), CH=CH<sub>2</sub>); 5.42 (1 H, dq, J 17.2(d), 1.5 (q), CH=CHH); 5.30 (1 H, dq, J 10.5 (d), 1.5 (q), CH=CHH); 4.55 (2 H, dt, J 5.3 (d), 1.5 (t), CH<sub>2</sub>CH=CH<sub>2</sub>); 4.30 (2 H, s, CH<sub>2</sub>N<sub>3</sub>). δ<sub>c</sub> (75 MHz, CDCl<sub>3</sub>, 25 °C): 158.9, 136.9 (arom. quat.), 133.1 (CH=CH<sub>2</sub>), 129.9, 120.6, 114.6, 114.5 (ArCH), 117.8 (CH=CH<sub>2</sub>), 68.8 (CH<sub>2</sub>O), 54.7 (CH<sub>2</sub>N). IR: ν<sub>max</sub>/cm<sup>-1</sup> 3064, 2925, 2870, 2094, 1649, 1599, 1586, 1489, 1448, 1424, 1342, 1263, 1157, 1098, 1027, 994, 927, 878, 854,

783, 762, 695, 650. GC-MS (method 1)  $t_R$  5.02 min,  $m/z$  (%) 189 (1.2)  $[M]^+$ , 120 (5.5), 92 (6.2), 91 (6.7), 79 (5.5), 78 (9.2), 77 (8.1), 65 (20), 64 (6.8), 63 (10), 51 (9.5), 50 (6.2), 41 (100), 39 (46), 38 (7.0). A solution of this azide (2.63 g, 13.9 mmol) in dry DMF (40 mL) cooled to 0 °C was treated with  $PMe_3$  (1 M in toluene, 15.3 mL, 15.3 mmol). When the gas evolution ceased, the mixture was warmed up to r.t. and stirred for 2 h. Then  $H_2O$  (1 mL, 55.6 mmol) was added and the reaction was further stirred for 2 h at r.t. After evaporation, the mixture was treated with saturated aqueous  $Na_2CO_3$  and extracted with AcOEt. The organic extracts were washed with saturated brine and evaporated to give crude 3-allyloxybenzylamine, that was not purified, but used as such for the Ugi reaction. It was just controlled at  $^1H$  NMR, that showed a purity > 95%.  $\delta_H$ (300 MHz,  $CDCl_3$ , 25 °C): 7.24 (1 H, t, J 8.1); 6.92-6.87 (2 H, m); 6.85-6.77 (1 H, m); 6.06 (1 H, ddt, J 10.5, 17.3 (d), 5.3 (t),  $CH=CH_2$ ); 5.45 (1 H, dq, J 17.3 (d), 1.5 (q),  $CH=CHH$ ); 5.28 (1 H, dq, J 10.5 (d), 1.5 (q),  $CH=CHH$ ); 4.55 (2 H, dt, J 5.3 (d), 1.5 (t),  $CH_2CH=CH_2$ ); 3.84 (2 H, s,  $CH_2NH_2$ ).

***N*-(4-allyloxyphenyl)formamide.** A solution of 4-allyloxyaniline<sup>51</sup> (499 mg, 3.35 mmol) in dry  $CH_2Cl_2$  (17 mL) at 0 °C was treated with formic acid (152  $\mu$ L, 4.02 mmol), 4-dimethylaminopyridine (DMAP) (82 mg, 0.67 mmol) and, finally, dicyclohexylcarbodiimide (DCC) (760 mg, 3.69 mmol). After 2 h at r.t., the resulting suspension was filtered through a celite cake washing with  $Et_2O$  + 2% of  $CH_2Cl_2$ . After evaporation and chromatography ( $CH_2Cl_2$  / AcOEt 7:1), *N*-(4-allyloxyphenyl)formamide was obtained (564 mg, 95%) as yellow solid. M.p.: 50.9–52.1 °C ( $CH_2Cl_2$ ).  $R_f$  = 0.45 (PE / AcOEt 1:1).  $\delta_H$ (300 MHz,  $CDCl_3$ , 25 °C)(two conformers in about 1:1 ratio are visible): 8.50 (0.5 H, d, J 11.6, *CHO* of 1 conformer); 8.34 (0.5 H, d, J 1.8, 0.5 H, *CHO* of 1 conformer); 7.59 (0.5 H, broad s, *NH* of 1 conformer); 7.44 (1 H, d, J 9.0, *ArCH* of 1 conformer); 7.12 (0.5 H, broad s, *NH* of 1 conformer); 7.03 (1 H, d, J 9.0, *ArCH* of 1 conformer); 6.91 (1 H, d, J 9.0, *ArCH* of 1 conformer); 6.89 (1 H, d, J 9.0, *ArCH* of 1 conformer); 6.05 (1 H, ddt, J 10.5 17.2 (d), 5.3 (t),  $CH=CH_2$ ); 5.46-5.36 (1 H, m,  $CH=CHH$ ); 5.33-5.26 (1 H, m,  $CH=CHH$ ); 4.55-4.50 (2 H, m,  $CH_2CH=CH_2$ ).  $\delta_C$  (75 MHz,  $CDCl_3$ , 25 °C): 163.3, 159.3 (C=O), 156.4, 155.5, 130.2, 129.8 (arom. quat.), 133.0, 132.9 ( $CH=CH_2$ ), 121.7, 121.2, 115.6, 114.9 (*ArCH*), 117.8, 117.6 ( $CH=CH_2$ ), 69.0, 68.9 ( $CH_2O$ ). IR:  $\nu_{max}/cm^{-1}$  3297, 3269, 3208, 3144, 3106, 3084, 3020, 2977, 2941, 2925, 2869, 2803, 2771, 1657, 1644, 1612, 1547, 1507, 1465, 1426, 1409, 1391, 1370, 1340, 1327, 1303, 1254, 1229, 1177, 1151, 1123, 1111, 1062, 1013, 1002, 940, 931, 873, 839, 822, 749, 737, 709, 648, 633. GC-MS (method 2)  $t_R$  7.88 min,  $m/z$  (%) 177 (43)  $[M]^+$ , 137 (8.5), 136 (100), 109 (11), 108 (99), 81 (6.4), 80 (50), 65 (8.6), 63 (5.2), 54 (5.5), 53 (26), 52 (16), 41 (37), 39 (27).  $m/z$  (ESI+) 178.0867 (M +  $H^+$ ).  $C_{10}H_{12}O_2N$  requires 178.0868.

**4-allyloxyphenyl isocyanide.** A solution of *N*-(4-allyloxyphenyl)formamide (130 mg, 0.734 mmol) in dry  $CH_2Cl_2$  (7 mL) was treated with  $Et_3N$  (470  $\mu$ L, 3.37 mmol) and cooled at -30 °C. Then  $POCl_3$  (103  $\mu$ L, 1.10 mmol) was added dropwise. After stirring for 1 h at -30 °C, the cold mixture was poured in saturated aqueous  $NaHCO_3$  and extracted with  $Et_2O$ . The organic extracts were washed with saturated brine and

evaporated. The crude was purified by chromatography (PE /  $Et_2O$  15:1) to give pure 4-allyloxyphenyl isocyanide as green oil (107 mg, 91%).  $R_f$  = 0.30 (PE /  $Et_2O$  15:1).  $\delta_H$ (300 MHz,  $CDCl_3$ , 25 °C): 7.30 (2 H, d, J 8.9, *ArCH*); 6.88 (2 H, d, J 8.9, *ArCH*); 6.03 (1 H, ddt, J 10.5, 17.3 (d), 5.3 (t),  $CH=CH_2$ ); 5.41 (1 H, dq, J 17.3 (d), 1.5 (q),  $CH=CHH$ ); 5.32 (1 H, dq, J 10.5 (d), 1.5 (q),  $CH=CHH$ ); 4.55 (2 H, dt, J 5.3 (d), 1.5 (t),  $CH_2CH=CH_2$ ).  $\delta_C$ (75 MHz,  $CDCl_3$ , 25 °C): 162.5 (NC), 158.8, 119.6 (broad) (arom. quat.), 132.3 ( $CH=CH_2$ ), 127.7, 115.3 (*ArCH*), 118.3 ( $CH=CH_2$ ), 69.0 ( $CH_2O$ ). IR:  $\nu_{max}/cm^{-1}$  3675, 3082, 2986, 2901, 2123, 1735, 1648, 1605, 1584, 1502, 1456, 1423, 1409, 1383, 1298, 1247, 1230, 1192, 1164, 1109, 1067, 1048, 1015, 995, 928, 830, 739, 700, 647, 618. GC-MS (method 2)  $t_R$  5.67 min,  $m/z$  (%) 159 (92)  $[M]^+$ , 158 (19), 144 (19), 132 (7.0), 131 (8.0), 130 (19), 119 (29), 103 (5.9), 102 (11), 91 (11), 90 (12), 76 (7.7), 75 (8.6), 64 (19), 63 (13), 41 (100), 39 (19).  $m/z$  (ESI+) 160.0769 (M +  $H^+$ ).  $C_{10}H_{10}ON$  requires 160.0762.

***N*-(4-(2-(2-(allyloxy)ethoxy)ethoxy)phenyl)formamide.** Known 1-(2-(2-(allyloxy)ethoxy)ethoxy)-4-nitrobenzene was prepared according to literature procedures.<sup>52, 53</sup> This compound (760 mg, 2.84 mmol) was dissolved in EtOH (33 mL) and treated with Fe powder (1.27 g, 22.7 mmol) and a solution of  $NH_4Cl$  (607 mg, 11.4 mmol) in deionized  $H_2O$  (6 mL). After stirring for 2 h at 75 °C, the resulting black suspension was filtered through a celite cake washing with MeOH. After evaporation, the crude was treated with saturated aqueous  $NaHCO_3$  and extracted with AcOEt. The organic extracts were washed with saturated brine and evaporated. The resulting crude 3-(2-(2-(allyloxy)ethoxy)ethoxy)aniline was directly treated with ethyl formate (4 mL) and stirred at 60 °C for 6 days. After evaporation, the crude was purified by chromatography (PE / AcOEt 1:1) to give pure *N*-(4-(2-(2-(allyloxy)ethoxy)ethoxy)phenyl)formamide as brown oil (506 mg, 67% from 1-(2-(2-(allyloxy)ethoxy)ethoxy)-4-nitrobenzene).  $R_f$  = 0.24 (PE / AcOEt 1:1).  $\delta_H$ (300 MHz,  $CDCl_3$ , 25 °C)(two conformers in about 1:1 ratio are visible): 8.50 (0.5 H, d, J 11.6, *CHO* of 1 conformer); 8.33 (0.5 H, d, J 1.8, *CHO* of 1 conformer); 7.46 (0.5 H, broad s, *NH* of 1 conformer); 7.43 (1 H, d, J 9.0, *ArCH* of 1 conformer); 7.12 (0.5 H, broad s, *NH* of 1 conformer); 7.01 (1 H, d, J 9.0, *ArCH* of 1 conformer); 6.91 (1 H, d, J 9.0, *ArCH* of 1 conformer); 6.89 (1 H, d, J 9.0, *ArCH* of 1 conformer); 5.92 (1 H, ddt, J 10.4, 17.2 (d), 5.7 (t),  $CH=CH_2$ ); 5.28 (1 H, dq, J 17.2 (d), 1.5 (q),  $CH=CHH$ ); 5.19 (1 H, dq, J 10.5 (d), 1.5 (q),  $CH=CHH$ ); 4.15-4.09 (2 H, m,  $CH_2O$ ); 4.04 (2 H, dt, J 5.7(d), 1.5 (t),  $OCH_2CH=CH_2$ ); 3.89-3.83 (2 H, m,  $CH_2O$ ); 3.76-3.69 (2 H, m,  $CH_2O$ ); 3.67-3.60 (m, 2 H,  $CH_2O$ ).  $\delta_C$ (75 MHz,  $CDCl_3$ , 25 °C):  $\delta$  = 163.1, 159.3 (C=O), 156.4, 155.4, 130.4, 129.9 (arom. quat.), 134.3 ( $CH=CH_2$ ), 121.5, 121.1, 115.4, 114.6 (*ArCH*), 117.2 ( $CH=CH_2$ ), 72.0, 70.6, 69.5, 69.2, 67.5, 67.4 ( $CH_2O$ ). IR:  $\nu_{max}/cm^{-1}$  3676, 3274, 3130, 3071, 2871, 1669, 1602, 1536, 1509, 1455, 1412, 1351, 1290, 1234, 1176, 1127, 1090, 1062, 994, 923, 872, 827, 726, 643, 633. GC-MS (method 1)  $t_R$  9.19 min,  $m/z$  (%) 163 ( $[M-OCH_2CH_2OAllyl]^+$ , 1.0) 108 (8.7), 87 (8.8), 85 (9.2), 80 (8.1), 65 (9.7), 53 (7.0), 45 (9.5), 44 (5.4), 43 (21), 41 (100), 39 (10).  $m/z$  (ESI+): 266.1392 (M +  $H^+$ ).  $C_{14}H_{20}O_4N$  requires 266.1392.

***N*-4-(2-(2-(allyloxy)ethoxy)ethoxy)phenyl) isocyanide.** A solution of *N*-4-(2-(2-(allyloxy)ethoxy)ethoxy)phenyl)formamide (374 mg, 1.41 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (7 mL) was treated with Et<sub>3</sub>N (590 μL, 4.23 mmol) and cooled at 0 °C. Then diphosgene (103 μL, 0.85 mmol) was added dropwise. After stirring for 1 h at 0 °C, the cold mixture was poured in saturated aqueous NaHCO<sub>3</sub> and extracted with Et<sub>2</sub>O. The organic extracts were washed with saturated brine and evaporated. The crude was purified by chromatography (PE / AcOEt 8:2) to give pure *N*-4-(2-(2-(allyloxy)ethoxy)ethoxy)phenyl) isocyanide as yellow oil (307 mg, 88%). *R*<sub>f</sub> = 0.34 (PE / AcOEt 8:2). δ<sub>H</sub>(300 MHz, CDCl<sub>3</sub>, 25 °C): 7.30 (2 H, d, *J* 9.0, ArCH); 6.89 (2 H, d, *J* 9.0, ArCH); 5.92 (1 H, ddt, *J* 10.4, 17.2 (d), 5.7 (q), CH=CH<sub>2</sub>); 5.28 (1 H, dq, *J* 17.2 (d), 1.6 (q), CH=CHH); 5.19 (1 H, dq, *J* 10.5 (d), 1.2 (q), CH=CHH); 4.17-4.11 (2 H, m, CH<sub>2</sub>O); 4.03 (2 H, dt, *J* 5.7 (d), 1.3 (t), OCH<sub>2</sub>CH=CH<sub>2</sub>); 3.89-3.85 (2 H, m, CH<sub>2</sub>O); 3.75-3.69 (2 H, m, CH<sub>2</sub>O); 3.66-3.60 (2 H, m, CH<sub>2</sub>O). δ<sub>C</sub>(75 MHz, CDCl<sub>3</sub>, 25 °C): 162.5 (NC), 159.1, 119.3 (broad) (arom. quat.), 134.6 (CH=CH<sub>2</sub>), 127.6, 155.2 (ArCH), 117.1 (CH=CH<sub>2</sub>), 72.2, 70.8, 69.5, 69.3, 67.7 (CH<sub>2</sub>O). IR: ν<sub>max</sub>/cm<sup>-1</sup> 3676, 3078, 2871, 2122, 1741, 1646, 1605, 1585, 1504, 1453, 1423, 1394, 1352, 1298, 1252, 1194, 1164, 1127, 1108, 1058, 995, 923, 883, 832, 724, 681, 641. GC-MS (method 1) *t*<sub>R</sub> 7.50 min, *m/z* (%) 159 (1.1), 102 (6.9), 85 (7.1), 73 (6.0), 71 (5.2), 45 (11), 43 (18), 41 (100), 39 (13). *m/z* (ESI<sup>+</sup>): 248.1288 (M + H<sup>+</sup>). C<sub>14</sub>H<sub>18</sub>O<sub>3</sub>N requires 248.1287.

***(R,S)*-N-(4-Acetoxyphenyl)-N-(1-(4-acetoxyphenyl)-2-(tert-butylamino)-2-oxoethyl)propionamide 2a.** A solution of aldehyde **4**<sup>25</sup> (228 mg, 2.0 mmol) in dry methanol (6.7 mL) was treated with amine **3**<sup>23, 24</sup> (302 mg, 2.0 mmol), propionic acid (150 μL, 2.0 mmol), and *tert*-butyl isocyanide (225 μL, 2.0 mmol). The solution was stirred at r.t. for 5 h. Then the solvent was evaporated and the crude purified by chromatography (PE / AcOEt 1:1) to give pure **2a** as a slightly brown solid (665 mg, 73%). M.p. = 144.6-146.8 °C. *R*<sub>f</sub> = 0.29 (PE / AcOEt 40:60). δ<sub>H</sub>(300 MHz, CDCl<sub>3</sub>, 25 °C): 7.13 (2 H, d, *J* 8.5, ArCH from aldehyde); 6.92 (2 H, d, *J* 8.5, ArCH from aldehyde); 7.05-6.85 (2 H, broad m, ArCH from amine) (NOTE: the other 2 ArCH from amine give a very broad signal from 7.50 to 7.00), 5.99 (1H, 1 H, CH), 5.72 (1H, s, NH), 2.25 (6H, s, CH<sub>3</sub>CO), 2.12-2.00 (2H, m, CH<sub>2</sub>CH<sub>3</sub>), 1.34 (9H, s, (CH<sub>3</sub>)<sub>3</sub>C), 1.04 (3 H, t, *J* = 7.4, CH<sub>2</sub>CH<sub>3</sub>). δ<sub>C</sub>(75 MHz, CDCl<sub>3</sub>, 25 °C): 174.3 (C=O), 169.0 (C=O), 168.8 (C=O), 168.7 (C=O), 150.5, 150.0, 137.2, 132.3 (quat.), 131.5, 131.3, 121.8, 121.4 (ArCH), 64.3 (CH), 51.5 (C(CH<sub>3</sub>)<sub>3</sub>), 28.5 (C(CH<sub>3</sub>)<sub>3</sub>), 28.3 (CH<sub>2</sub>CH<sub>3</sub>), 21.0 (CH<sub>3</sub>CO), 9.3 (CH<sub>2</sub>CH<sub>3</sub>). IR: ν<sub>max</sub>/cm<sup>-1</sup> 3339, 3234, 3078, 2976, 1759, 1681, 1636, 1551, 1504, 1459, 1418, 1390, 1366, 1305, 1270, 1251, 1209, 1186, 1166, 1160, 1105, 1099, 1044, 1012, 958, 941, 910, 859, 848, 813, 784, 775, 744, 735, 722, 656, 632. *m/z* (ESI<sup>+</sup>): 455.2180 (M + H<sup>+</sup>). C<sub>25</sub>H<sub>31</sub>O<sub>6</sub>N<sub>2</sub> requires 455.2182.

***(R,S)*-N-(4-Hydroxyphenyl)-N-(2-(tert-butylamino)-1-(4-hydroxyphenyl)-2-oxoethyl)propionamide 1a.** A solution of compound **2a** (137 mg, 0.30 mmol) in tetrahydrofuran (2.25 mL) was treated, at r.t., with 1 M aqueous LiOH (0.78 μL, 0.78 mmol). The solution became yellow. After 20 h, the reaction not being yet complete, other 0.39 μL of LiOH solution were added. After other 20 h, the reaction was worked out with a 1

M NaH<sub>2</sub>PO<sub>4</sub> solution and extracted with AcOEt. After evaporation of the organic phase, two consecutive chromatographies (first PE / AcOEt 3:7; then CH<sub>2</sub>Cl<sub>2</sub> / MeOH 93:7) afforded pure **1a** as a white solid (85 mg, 76%). M.p. = 203.4-204.1 °C. *R*<sub>f</sub> = 0.24 (PE / AcOEt 30:70). δ<sub>H</sub>(300 MHz, DMSO-*d*<sub>6</sub>, 50 °C): 9.22 (1 H, br s, OH), 9.16 (1 H, br s, OH); 7.28 (1 H, s, NH); 6.81 (2 H, d, *J* 8.5, ArCH from aldehyde); 6.49 (2 H, d, *J* 8.5, ArCH from aldehyde); 6.70-6.45 (2 H, broad m, ArCH from amine) (NOTE: the other 2 ArCH from amine give a very broad signal from 7.30 to 6.80), 5.87 (1H, s, CH), 2.01-1.81 (2H, m, CH<sub>2</sub>CH<sub>3</sub>), 1.23 (9H, s, (CH<sub>3</sub>)<sub>3</sub>C), 0.89 (3 H, t, *J* = 7.4, CH<sub>2</sub>CH<sub>3</sub>). δ<sub>C</sub>(75 MHz, DMSO-*d*<sub>6</sub>, 50 °C): 172.6 (C=O), 169.5 (C=O), 156.1, 155.8, 131.5, 131.1 (quat.), 130.9, 126.1, 114.5, 114.2 (ArCH), 63.0 (CH), 49.9 (C(CH<sub>3</sub>)<sub>3</sub>), 28.3 (C(CH<sub>3</sub>)<sub>3</sub>), 27.3 (CH<sub>2</sub>CH<sub>3</sub>), 9.2 (CH<sub>2</sub>CH<sub>3</sub>). IR: ν<sub>max</sub>/cm<sup>-1</sup> 3274, 3234, 2969, 1661, 1614, 1593, 1511, 1452, 1393, 1365, 1258, 1221, 1174, 1096, 1044, 1023, 960, 845, 816, 780, 740, 633. *m/z* (ESI<sup>+</sup>): 371.1976 (M + H<sup>+</sup>). C<sub>21</sub>H<sub>27</sub>O<sub>4</sub>N<sub>2</sub> requires 371.1971. HPLC (see supplementary information) showed a purity of 99.5%.

***(R,S)*-4-((tert-Butyldimethylsilyloxy)-N-(4-((tert-butylidimethylsilyloxy)phenyl)-N-(2-((tert-butylidimethylsilyloxy)phenyl)amino)-2-oxo-1-phenylethyl)benzamide 11.** A solution of amine **9** (224 mg, 1.00 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (10.0 mL) was treated at r.t. with benzaldehyde (102 μL, 1.00 mmol) and anhydrous MgSO<sub>4</sub> (100 mg) and stirred overnight. After filtration of MgSO<sub>4</sub> and evaporation, the residue was taken up in MeOH (5.0 mL), added with freshly activated powdered 3 Å molecular sieves (50 mg), and finally treated with acid **10** (252 mg, 1.00 mmol) and isocyanide **8** (234 mg, 1.00 mmol). After 17 h, the mixture was filtered, evaporated to dryness and chromatographed (PE / AcOEt 60:40) to give pure **11** as a yellow-brown solid (200 mg, 25%). M.p. = 156.7-157.3 °C. *R*<sub>f</sub> = 0.58 (PE / AcOEt 6:4). δ<sub>H</sub>(300 MHz, CDCl<sub>3</sub>, 25 °C): δ 8.19 (1 H, s, NH), 7.35 (2 H, d, *J* 9.0, ArH), 7.32-7.18 (7 H, m), 6.85 (2 H, broad d, *J* = 7.7 Hz, ArH), 6.74 (2 H, d, *J* = 8.8 Hz, ArH), 6.56 (2 H, d, *J* = 8.7 Hz, ArH), 6.50 (2 H, d, *J* = 8.9 Hz, ArH), 6.34 (1 H, s, CH), 0.96, 0.92, 0.91 (3 x 9 H, 3 s, (CH<sub>3</sub>)<sub>3</sub>C); 0.16, 0.11, 0.08 (3 x 6 H, 3 s, (CH<sub>3</sub>)<sub>3</sub>Si). δ<sub>C</sub>(75 MHz, CDCl<sub>3</sub>, 25 °C, TMS): 171.1, 168.1 (C=O), 156.9, 154.6, 152.3, 135.0, 134.5, 131.50, 131.2 (quat.), 131.4 (x2), 130.8 (x2), 130.2 (x2), 128.6, 128.5 (x2), 121.7 (x2), 120.2 (x2), 120.0 (x2), 119.1 (x2) (ArCH), 67.4 (CH), 25.69, 25.64, 25.56 (C(CH<sub>3</sub>)<sub>3</sub>), 18.38 (C(CH<sub>3</sub>)<sub>3</sub>), 18.24, 18.19, 18.14 (quat. C *t*-Bu), -4.48 (x2), -4.54 (CH<sub>3</sub>Si). IR: ν<sub>max</sub>/cm<sup>-1</sup> 3260, 3201, 3075, 2957, 2930, 2896, 2858, 1689, 1617, 1604, 1551, 1505, 1472, 1462, 1410, 1389, 1362, 1341, 1253, 1201, 1166, 1103, 1080, 1052, 1006, 966, 908, 831, 803, 777, 764, 735, 716, 698, 666, 638, 623. *m/z* (ESI<sup>+</sup>): 797.4187 (M + H<sup>+</sup>). C<sub>45</sub>H<sub>65</sub>O<sub>5</sub>N<sub>2</sub>Si<sub>3</sub> requires 797.4201.

***(R,S)*-4-Hydroxy-N-(4-hydroxyphenyl)-N-(2-((4-hydroxyphenyl)amino)-2-oxo-1-phenylethyl)benzamide 1b.** A solution of compound **2a** (117 mg, 0.15 mmol) in tetrahydrofuran (1.2 mL) was treated, at r.t., with 1 M aqueous LiOH (0.59 μL, 0.59 mmol). The solution became orange. After 22 h the solution was evaporated, taken up with MeOH, and treated with previously washed Amberlyst 15 acid resin until pH = 7. The resin was filtered off and the solution



evaporated to dryness and chromatographed (PE / AcOEt 4:6 + 2% EtOH) to give pure **1b** as a white solid (54 mg, 81%). M.p. = 177.7–178.2 °C.  $R_f$  = 0.34 (PE / AcOEt 40:60 + 2% EtOH).  $\delta_H$ (300 MHz, DMSO-*d*<sub>6</sub>, 25 °C):  $\delta$  9.98, 9.68, 9.24 (3 x 1 H, 3 s, OH), 7.40 (2 H, d,  $J$  = 9.0, ArH), 7.24–7.10 (5 H, m, ArH+NH), 7.07 (2 H, d,  $J$  = 9.3, ArH), 6.69 (2 H, d,  $J$  = 9.0, ArH), 6.51 (2 H, d,  $J$  = 8.7, ArH), 6.33 (2 H, broad d,  $J$  = 8.4, ArH), 6.26 (1H, s, CH).  $\delta_C$ (75 MHz, DMSO-*d*<sub>6</sub>, 25 °C): 169.8, 168.2 (C=O), 158.1, 155.4, 153.1, 135.2, 132.0, 131.9, 130.8 (quat.), 130.2 (x4), 127.8 (x2), 127.7, 127.0 (x2), 120.6 (x2), 115.0 (x2), 114.2 (x2), 114.0 (x2) (ArCH) 64.9 (CH). IR:  $\nu_{max}/cm^{-1}$  3275, 3234, 1665, 1607, 1509, 1440, 1365, 1223, 1167, 1103, 1081, 831, 761, 729, 698, 626.  $m/z$  (ESI+): 455.1604 (M + H<sup>+</sup>). C<sub>27</sub>H<sub>23</sub>O<sub>5</sub>N<sub>2</sub> requires 455.1607. HPLC (see supplementary information) showed a purity of 96%.

**General procedure for the preparation of polyphenols 1c,e,h,i through allylated derivatives 14c,e,h,i (Method A).** A solution of the appropriate aldehyde (1 equiv) in dry EtOH and trifluoroethanol in 1:1 ratio (0.26 M) was treated with the amine (1.1 equiv) and molecular sieves (3 Å, 50 mg/mmole). After 5 h, the acid (1.1 equiv), and the isocyanide (1.1 equiv) were added. The reaction mixture was stirred at r.t. for 2–4 days, then filtered with celite on a sintered funnel, washed with AcOEt, concentrated and purified by chromatography. A 0.1 M solution of the Ugi product in MeCN under nitrogen atmosphere, was treated with PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (0.025 equiv for allyl group) and ammonium formate (2.2 equiv for allyl group) at 80 °C for 2 h in a sealed flask. Then, the crude was diluted with AcOEt, washed with saturated aqueous NaHCO<sub>3</sub>. After evaporation, the residue was eluted from a column of silica gel with the suitable eluent.

**(*R,S*)-(E)-N-Benzyl-N-(2-(*tert*-butylamino)-1-(4-hydroxyphenyl)-2-oxoethyl)-3-(4-hydroxy-3-methoxyphenyl)acrylamide 1c.** Following the general procedure A, a mixture of aldehyde **12** (113 mg, 0.70 mmol), benzylamine (80 µL, 0.77 mmol), acid **13** (180 mg, 0.77 mmol), *t*-butyl isocyanide (87 µL, 1.19 mmol) and 3 Å molecular sieves (32 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt 50:50) compound **14c** was obtained pure as white foam (275 mg, 69%). Then a mixture of **14c** (255 mg, 0.45 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (16 mg, 0.023 mmol) and ammonium formate (124 mg, 1.98 mmol) was stirred for 2 h at 80 °C. After work-up and purification (chromatography with PE / AcOEt 3:4, followed by treatment with active coal) compound **1c** was obtained pure as white solid (177 mg, 80%). M.p. = 125 °C with decomposition.  $R_f$  = 0.19 (PE / AcOEt 50:50).  $\delta_H$ (300 MHz, DMSO-*d*<sub>6</sub>, 90 °C):  $\delta$  9.06 (2 H, s, OH), 7.43 (1 H, broad s, NH), 7.41 (1 H, d,  $J$  = 15.0, ArCH=CH), 7.20–6.87 (9 H, m, ArH, ArCH=CH), 6.80–6.60 (1 H, broad signal, ArCH=CH), 6.76 (1 H, d,  $J$  8.1, *H* meta to OMe), 6.67 (2 H, d,  $J$  8.7, *H* ortho to OH), 6.00 (1 H, broad s, CHN), 4.87 (1 H, d,  $J$  16.7, CHHPh), 4.55 (1 H, d,  $J$  16.7, CHHPh), 3.78 (3 H, s, OCH<sub>3</sub>), 1.26 (9 H, s, C(CH<sub>3</sub>)<sub>3</sub>).  $\delta_C$ (75 MHz, DMSO-*d*<sub>6</sub>, 25 °C) (note: at this temperature, 2 conformers are visible and thus most signals are doubled): 169.6, 169.2, 167.1, 166.9 (C=O), 156.9, 156.8, 148.4, 147.7, 147.6, 140.1, 139.4, 130.0 (quat.), 141.7 (ArCH=CH), 130.4, 127.8, 127.4, 126.8, 126.4, 126.2, 126.1,

122.3, 121.9, 116.3, 114.9, 110.6 (ArCH), 115.4 (ArCH=CH), 62.8, 60.2 (CHN), 55.5, 55.4 (OCH<sub>3</sub>), 50.2 (C(CH<sub>3</sub>)<sub>3</sub>), 48.0 (CH<sub>2</sub>Ph), 28.3, 28.1 (C(CH<sub>3</sub>)<sub>3</sub>). IR:  $\nu_{max}/cm^{-1}$  3283, 2967, 1641, 1589, 1511, 1452, 1429, 1364, 1265, 1203, 1171, 1123, 1080, 1030, 976, 947, 891, 865, 837, 814, 726, 696.  $m/z$  (ESI+): 489.2396 (M + H<sup>+</sup>). C<sub>29</sub>H<sub>33</sub>O<sub>5</sub>N<sub>2</sub> requires 489.2389. HPLC (see supplementary information) showed a purity of 99%.

**(*R,S*)-(E)-N-Benzyl-N-(2-(*tert*-butylamino)-1-phenyl-2-oxoethyl)-3-(4-hydroxy-3-methoxyphenyl)acrylamide 1e.**

Following the general procedure A, a mixture of benzaldehyde (67 mg, 0.63 mmol), benzylamine (76 µL, 0.69 mmol), (*E*)-3-(3,4-bis(allyloxy)phenyl)acrylic acid (179 mg, 0.69 mmol), *t*-butyl isocyanide (78 µL, 0.69 mmol) and 3 Å molecular sieves (28 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt 75:25) compound **14e** was obtained pure as white foam (208 mg, 72%). Then a mixture of **14e** (200 mg, 0.37 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (12 mg, 0.017 mmol) and ammonium formate (95 mg, 1.50 mmol) was stirred for 2 h at 80 °C. After work-up and purification (two consecutive chromatographies with PE / AcOEt from 60:40 to 50:50 and a filtration with celite on a sintered funnel) compound **1e** was obtained pure as pale yellow solid (101 mg, 59%). M.p. = 120.0 – 121.0 °C (MeOH).  $R_f$  = 0.40 (PE / AcOEt 50:50).  $\delta_H$ (300 MHz, DMSO-*d*<sub>6</sub>, 80 °C) (note: the OH protons give a very broad signal around 9 ppm): 7.61 (1 H, broad s, 1 H, NH), 7.38 (1 H, d,  $J$  = 15.2, ArCH=CH), 7.30–7.03 (10 H, m, ArH), 6.87 (1 H, s, *H* ortho to OH and CH=CH), 6.79, 6.72 (2 H, AB syst.,  $J$  8.3, *H* para to OH and meta to CH=CH), 6.65 (1 H, broad d,  $J$  15.2, ArCH=CH), 6.12 (1 H, broad s, CHN), 4.91, 4.65 (2 H, AB syst.,  $J$  17.0, CH<sub>2</sub>Ph), 1.24 (9 H, s, C(CH<sub>3</sub>)<sub>3</sub>).  $\delta_C$ (75 MHz, DMSO-*d*<sub>6</sub>, 25 °C) (note: at this temperature, 2 conformers are visible and thus most signals are doubled, although one conformer is prevailing, only the peaks of major conformer are reported): 168.9, 167.1 (C=O), 147.6, 145.3, 139.5, 136.9, 128.2 (quat.), 142.5 (ArCH=CH), 128.9 (x2), 128.1 (x2), 127.8 (x2), 127.4, 126.2, 125.8 (x2), 120.7, 115.4, 114.1 (ArCH), 115.1 (ArCH=CH), 60.6 (CHN), 54.8 (OCH<sub>3</sub>), 50.3 (C(CH<sub>3</sub>)<sub>3</sub>), 48.1 (CH<sub>2</sub>Ph), 28.2 (C(CH<sub>3</sub>)<sub>3</sub>). IR (ATR):  $\nu$  = 3276, 3064, 2969, 1640, 1578, 1513, 1451, 1413, 1363, 1278, 1192, 1112, 1080, 1032, 975, 948, 893, 845, 810, 753, 730, 696, 617 cm<sup>-1</sup>.  $m/z$  (ESI-): 457.2143 (M - H<sup>+</sup>). C<sub>28</sub>H<sub>29</sub>O<sub>4</sub>N<sub>2</sub> requires 457.2127. HPLC (see supplementary information) showed a purity of 100%.

**(*R,S*)-N-(4-hydroxyphenyl)-N-(1-(4-hydroxyphenyl)-2-((4-hydroxyphenyl)amino)-2-oxoethyl)-propanamide 1h.**

Following the general procedure A, a mixture of aldehyde **12** (170 mg, 1.05 mmol), 4-allyloxylaniline (173 mg, 1.16 mmol), propionic acid (87 µL, 1.16 mmol), 4-allyloxyphenyl isocyanide (185 mg, 1.16 mmol) and 3 Å molecular sieves (50 mg) was stirred for 5 days at r.t. After usual work-up, the crude was diluted with AcOEt, washed with HCl 1 N to remove the excess of amine. Then, the crude was purified (PE / Et<sub>2</sub>O 1:2) obtaining compound **14h** as white foam (121 mg, 22%). Then a mixture of **14h** (102 mg, 0.20 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (7 mg, 0.01 mmol) and ammonium formate (83 mg, 1.32 mmol) was stirred for 2 h at 80 °C. After work-up and purification (from CH<sub>2</sub>Cl<sub>2</sub> / AcOEt 3:4 to AcOEt with 1% MeOH) compound **1h** was obtained as red solid (60 mg, 74%). A final treatment with

active coal gave **1h** as a pale yellow solid. M.p. = 158.0 – 160.0 °C (MeOH).  $R_f$  = 0.35 (CH<sub>2</sub>Cl<sub>2</sub> / AcOEt 3:4).  $\delta_H$ (300 MHz, DMSO-*d*<sub>6</sub>, 90 °C) (Note: the 3 phenolic OH exchange with H<sub>2</sub>O contained in the solvent giving a broad signal around 4.90 ppm): 9.38 (1 H, s, NH), 9.30 (1 H, s, OH), 7.32 (2 H, d, J 8.8, *H* ortho to CH), 7.0-6.80 (2 H, broad signal, *H* ortho to N), 6.86 (2 H, d, J 8.4, *H* ortho to NH), 6.67 (2 H, d, J 8.8, *H* meta to CH), 6.50 (4 H, d, J 8.4, *H* meta to N and to NH), 6.03 (2 H, s, CH), 1.97 (2 H, q, J 7.4, CH<sub>2</sub>CH<sub>3</sub>), 0.92 (3 H, t, J 7.5, CH<sub>2</sub>CH<sub>3</sub>).  $\delta_C$ (75 MHz, DMSO-*d*<sub>6</sub>, 25 °C): 173.1, 168.8 (C=O), 157.5, 157.0, 153.3, 130.7, 130.2, 124.5 (quat.), 131.7 (x2), 131.2 (x2), 120.15 (x2), 115.0 (x2), 114.8 (x4) (ArCH), 63.6 (CHN), 27.6 (CH<sub>2</sub>), 9.4 (CH<sub>3</sub>). IR:  $\nu_{max}/cm^{-1}$  3268, 3202, 3005, 1650, 1602, 1584, 1533, 1471, 1379, 1360, 1260, 1205, 1173, 1099, 1043, 1001, 965, 845, 815, 631.  $m/z$  (ESI+): 407.1605 (M + H<sup>+</sup>). C<sub>23</sub>H<sub>23</sub>O<sub>5</sub>N<sub>2</sub> requires 407.1607. HPLC (see supplementary information) showed a purity of 98%.

**(R,S)-(E)-N-(2-(tert-butylamino)-1-(4-hydroxyphenyl)-2-oxoethyl)-3-(4-hydroxy-3-methoxyphenyl)-N-(4-hydroxyphenyl)acrylamide 1i.** Following the general procedure A, a mixture of aldehyde **12** (94 mg, 0.58 mmol), 4-allyloxyaniline (95 mg, 0.64 mmol), acid **13** (150 mg, 0.64 mmol), *t*-butyl isocyanide (72  $\mu$ L, 0.64 mmol) and 3 Å molecular sieves (29 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt 3:2) compound **14i** was obtained pure as yellow foam (117 mg, 33%). Then a mixture of **14i** (88 mg, 0.15 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (8 mg, 0.011 mmol) and ammonium formate (63 mg, 1.00 mmol) was stirred for 2 h at 80 °C. After work-up and purification (the crude was triturated with PE/AcOEt) compound **1i** was obtained pure as white solid (50 mg, 68%). M.p. = 150 °C with decomposition.  $R_f$  = 0.10 (PE / AcOEt 50:50).  $\delta_H$ (300 MHz, DMSO-*d*<sub>6</sub>, 25 °C): 9.42 (2 H, s, OH), 9.30 (1 H, s, OH), 7.56 (1 H, s, NH), 7.38 (1 H, d, J 15.3, ArCH=CH), 6.88 (1 H, s, *H* ortho to OMe), 6.83 (2 H, d, J 8.4, *H* meta to OH), 6.76-6.67 (2 H, m, *H* meta and para to OMe), 6.70-6.40 (2 H, very broad signal, *H* meta to OH), 6.50 (4 H, d, J 8.4, *H* ortho to OH), 5.98 (1 H, s, CHN), 5.95 (1 H, d, J 15.3, ArCH=CH), 3.70 (3 H, s, OCH<sub>3</sub>), 1.24 (9 H, s, (CH<sub>3</sub>)<sub>3</sub>C).  $\delta_H$ (75 MHz, DMSO-*d*<sub>6</sub>, 25 °C): 169.7, 165.4 (C=O), 156.3, 156.0, 148.3, 147.5, 132.0, 130.7, 126.2 (quat.), 140.6 (ArCH=CH), 132.0 (x2), 131.2 (x2), 126.1, 120.2, 114.5 (x4), 112.2 (ArCH), 116.6 (ArCH=CH), 63.3 (CHN), 55.5 (OCH<sub>3</sub>), 50.1 (C(CH<sub>3</sub>)<sub>3</sub>), 28.4 (C(CH<sub>3</sub>)<sub>3</sub>). IR:  $\nu_{max}/cm^{-1}$  3269, 2966, 2930, 1662, 1640, 1592, 1511, 1450, 1388, 1366, 1258, 1216, 1161, 1121, 1030, 1009, 976, 936, 885, 840, 814, 791, 742, 723, 693, 645, 610.  $m/z$  (ESI+): 491.2187 (M + H<sup>+</sup>). C<sub>28</sub>H<sub>31</sub>O<sub>6</sub>N<sub>2</sub> requires 491.2182. HPLC (see supplementary information) showed a purity of 98.5%.

**General procedure for the preparation of acetylated polyphenols 2d,f,g,j,l,m,n,o,p,q and 15 (Method B).** A solution of the appropriate aldehyde (1 equiv) in dry EtOH and trifluoroethanol in 1:1 ratio (0.26 M) was treated with the amine (1.1 equiv) and molecular sieves (3 Å, 50 mg/mmole). After 5 h, the acid (1.1 equiv), and the isocyanide (1.1 equiv) were added. The reaction mixture was stirred at r.t. for 2-4 days, then filtered with celite on a sintered funnel, washed with AcOEt and concentrated. The residue was treated with saturated aqueous NaHCO<sub>3</sub> and extracted with AcOEt. The

organic extracts were washed with saturated brine and evaporated. The crude was purified by chromatography. A 0.1 M solution of the Ugi product in MeCN under nitrogen atmosphere, was treated with PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (0.025 equiv for allyl group) and ammonium formate (2.2 equiv for allyl group) at 80 °C in a sealed flask (2-5 h). Then, the crude was diluted with AcOEt, washed with saturated aqueous NaHCO<sub>3</sub>. After evaporation, the residue was directly dissolved in 1:1 pyridine / acetic anhydride (0.1 M) and stirred for 18 h at r.t. Then, the mixture was poured in 2 N HCl (final pH = 2) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic extracts were washed with saturated brine and evaporated, then the crude was purified by chromatography.

**(R,S)-(E)-3-(4-Acetoxy-3-methoxyphenyl)-N-(1-(4-acetoxyphenyl)-2-(tert-butylamino)-2-oxoethyl)-N-butylacrylamide 2d.** Following the general procedure B, a mixture of aldehyde **12** (176 mg, 1.08 mmol), *n*-butylamine (120  $\mu$ L, 1.19 mmol), acid **13** (280 mg, 1.19 mmol), *t*-butyl isocyanide (135  $\mu$ L, 1.19 mmol) and 3 Å molecular sieves (50 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt 75:25 + 1% EtOH) compound **14d** was obtained pure as yellow oil (330 mg, 57%, 91% based on the recovery of unreacted aldehyde). Then a mixture of **14d** (270 mg, 0.50 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (18 mg, 0.0025 mmol) and ammonium formate (140 mg, 2.22 mmol) was stirred for 5 h at 80 °C. After work-up, the crude was treated with 1:1 pyridine / acetic anhydride (3.7 mL) and stirred for 18 h at r.t. After work-up and purification (PE / AcOEt 7:3 + 3% EtOH) compound **2d** was obtained pure as white foam (226 mg, 76%).  $R_f$  = 0.28 (PE / AcOEt 7:3 + 3% EtOH).  $\delta_H$ (300 MHz, CDCl<sub>3</sub>, 25 °C):  $\delta$  7.72 (1 H, d, J 15.3, ArCH=CH), 7.47 (2 H, d, J 8.4), 7.15-7.01 (5 H, m), 6.78 (1 H, d, J 15.3, ArCH=CH), 6.04 (1 H, s, CH), 5.87 (1 H, s, NH), 3.85 (3 H, s, OCH<sub>3</sub>), 3.57-3.32 (mc = 3.46) (2 H, m, CH<sub>2</sub>N), 2.32, 2.31 (2 x 3 H, 2 s, CH<sub>3</sub>CO), 1.55-1.39 (1 H, m, CHH), 1.37 (9 H, s, (CH<sub>3</sub>)<sub>3</sub>C), 1.22-0.95 (3 H, m, CH<sub>2</sub> and CHH), 0.79 (3 H, t, J 7.2, CH<sub>3</sub>CH<sub>2</sub>).  $\delta_H$ (75 MHz, CDCl<sub>3</sub>, 25 °C): 169.3, 168.9 (x2), 166.9 (C=O), 151.3, 150.6, 140.9, 134.3, 133.4 (quat.), 142.8 (ArCH=CH), 130.5 (x2), 123.2, 121.9 (x2), 120.4, 111.6 (ArCH), 118.0 (ArCH=CH), 61.5 (CH), 58.8 (OCH<sub>3</sub>), 51.7 (C(CH<sub>3</sub>)<sub>3</sub>), 46.1 (NCH<sub>2</sub>), 32.6 (NCH<sub>2</sub>CH<sub>2</sub>), 28.6 (C(CH<sub>3</sub>)<sub>3</sub>), 21.1, 20.7 (CH<sub>3</sub>CO), 20.0 (CH<sub>2</sub>CH<sub>3</sub>), 13.5 (CH<sub>2</sub>CH<sub>3</sub>). IR:  $\nu_{max}/cm^{-1}$  3676, 3295, 3056, 2965, 2930, 2877, 2856, 1768, 1742, 1684, 1640, 1608, 1582, 1542, 1508, 1486, 1473, 1454, 1431, 1407, 1392, 1364, 1349, 1311, 1302, 1288, 1268, 1259, 1246, 1208, 1194, 1166, 1123, 1066, 1045, 1032, 1012, 979, 951, 937, 918, 906, 889, 865, 842, 815, 801, 792, 753, 737, 731, 709, 688, 633, 620.  $m/z$  (ESI+): 539.2758 (M + H<sup>+</sup>). C<sub>30</sub>H<sub>39</sub>O<sub>7</sub>N<sub>2</sub> requires 539.2757.

**(R,S)-(E)-N-(1-(4-Acetoxyphenyl)-2-(tert-butylamino)-2-oxoethyl)-3-(3,4-diacetoxyphenyl)-N-butylacrylamide 2f.** Following the general procedure B, a mixture of aldehyde **12** (596 mg, 3.70 mmol), benzylamine (445  $\mu$ L, 4.07 mmol), (*E*)-3-(3,4-bis(allyloxy)phenyl)acrylic acid (1.06 g, 4.07 mmol), *t*-butyl isocyanide (460  $\mu$ L, 4.07 mmol) and 3 Å molecular sieves (200 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt 2:1) compound **14f** was obtained pure as white foam (1.59 g, 73%). Then a mixture of **14f** (710 mg, 1.20 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (70 mg, 0.1 mmol) and ammonium

formate (500 mg, 7.90 mmol) was stirred for 2 h at 80 °C. After work-up, the crude was treated with 1:1 pyridine / acetic anhydride (8 mL) and stirred for 18 h at r.t. After work-up and purification (the crude was triturated with AcOEt and the mother liquor was purified by chromatography PE / Et<sub>2</sub>O 1:2 + 2% EtOH) pure compound **2f** was obtained as white solid (602 mg, 83%). M.p. = 189.0 – 190.0 °C (CH<sub>2</sub>Cl<sub>2</sub>). R<sub>f</sub> = 0.70 (CH<sub>2</sub>Cl<sub>2</sub> / MeOH 15:1). δ<sub>H</sub>(300 MHz, CDCl<sub>3</sub>, 25 °C): δ 7.70 (1 H, d, J 15.4, ArCH=CH), 7.38 (2 H, d, J 8.4), 7.25-7.10 (6 H, m), 7.02-6.92 (4 H, m), 6.66 (1 H, d, J 15.4, ArCH=CH), 6.06 (1 H, s, CH), 5.67 (1 H, s, NH), 4.90 (1 H, d, J 17.5, CHPh), 4.66 (1 H, d, J 17.5, CHPh), 2.27 (3 H, s, CH<sub>3</sub>CO), 2.26 (6 H, s, CH<sub>3</sub>CO), 1.35 (9 H, s, (CH<sub>3</sub>)<sub>3</sub>C). δ<sub>C</sub>(75 MHz, CDCl<sub>3</sub>, 25 °C): 169.1, 168.6, 168.0 (x2), 167.6 (C=O), 150.6, 143.0, 142.2, 137.9, 134.0, 132.6 (quat.), 141.9 (ArCH=CH), 130.8 (x2), 128.5 (x2), 127.0, 126.1 (x3), 123.7, 122.6, 121.7 (x2) (ArCH), 119.2 (ArCH=CH), 62.0 (CH), 51.7 (C(CH<sub>3</sub>)), 49.6 (NCH<sub>2</sub>), 28.6 (C(CH<sub>3</sub>)<sub>3</sub>), 21.1, 20.6, 20.5 (CH<sub>3</sub>CO). IR: ν<sub>max</sub>/cm<sup>-1</sup> 3295, 3071, 2939, 1758, 1694, 1646, 1602, 1546, 1505, 1451, 1408, 1368, 1302, 1258, 1185, 1162, 1122, 1013, 979, 953, 909, 832, 793, 730, 692, 638. m/z (ESI<sup>-</sup>): 599.2381 (M - H<sup>+</sup>). C<sub>34</sub>H<sub>35</sub>O<sub>8</sub>N<sub>2</sub> requires 599.2393.

**(R,S)-N-(4-Acetoxyphenyl)-N-(1-(4-Acetoxyphenyl)-2-((4-acetoxyphenyl)amino)-2-oxoethyl)benzamide 2g.** Following the general procedure B, a mixture of aldehyde **12** (122 mg, 0.75 mmol), 4-allyloxyaniline (114 μL, 0.82 mmol), benzoic acid (100 mg, 0.82 mmol), 4-allyloxyphenyl isocyanide (130 mg, 0.82 mmol) and 3 Å molecular sieves (70 mg) was stirred for 3 days at r.t. After usual work-up, the crude was diluted with AcOEt, washed with HCl 1 N to remove the excess of amine. Then, the crude was purified (PE / AcOEt 3:2) obtaining compound **14g** as brown foam (119 mg, 28%). Then a mixture of **14g** (94 mg, 0.16 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (9 mg, 0.01 mmol) and ammonium formate (67 mg, 1.06 mmol) was stirred for 4 h at 80 °C. After work-up, the crude was treated with 1:1 pyridine / acetic anhydride (3 mL) and stirred for 18 h at r.t. After work-up and purification (PE / AcOEt 1:1) pure compound **2g** was obtained as white foam (55 mg, 59%). R<sub>f</sub> = 0.72 (PE / AcOEt 3:7). δ<sub>H</sub>(300 MHz, CDCl<sub>3</sub>, 25 °C): 8.43 (1 H, s, NH), 7.48 (2 H, d, J 8.8), 7.28 (4 H, d, J 8.1), 7.25-7.10 (3 H, m), 7.02-6.93 (6 H, m), 6.78 (2 H, d, J 8.5), 6.42 (1 H, s, CH), 2.26 (6 H, s, CH<sub>3</sub>CO), 2.19 (3 H, s, CH<sub>3</sub>CO). δ<sub>H</sub>(75 MHz, CDCl<sub>3</sub>, 25 °C): δ 171.6, 169.5, 169.1, 168.7, 167.9 (C=O), 150.9, 149.5, 146.9, 138.0, 135.4, 135.3, 129.9 (quat.), 131.51 (x2), 131.43 (x2), 131.37, 128.6 (x2), 127.8 (x2), 121.9 (x4), 121.5 (x2), 120.9 (x2) (ArCH), 66.2 (CH), 21.1 (CH<sub>3</sub>CO). IR: ν<sub>max</sub>/cm<sup>-1</sup> 3275, 3054, 2980, 1743, 1691, 1615, 1594, 1533, 1551, 1488, 1463, 1401, 1333, 1301, 1198, 1133, 1101, 1045, 1007, 965, 841, 761, 732, 679, 630, 603. m/z (ESI<sup>+</sup>): 581.1914 (M + H<sup>+</sup>). C<sub>33</sub>H<sub>29</sub>O<sub>8</sub>N<sub>2</sub> requires 581.19244.

**(R,S)-4-Acetoxy-N-(1-(4-Acetoxyphenyl)-2-((4-acetoxyphenyl)amino)-2-oxoethyl)-N-phenylbenzamide 2j.** Following the general procedure B, a mixture of aldehyde **12** (122 mg, 0.75 mmol), aniline (74 μL, 0.82 mmol), 4-(allyloxy)benzoic acid (146 mg, 0.82 mmol), 4-allyloxyphenyl isocyanide (130 mg, 0.82 mmol) and 3 Å molecular sieves (70 mg) was stirred for 3 days at r.t. After usual work-up, the crude was diluted with AcOEt, washed with HCl 1 N to remove the excess of amine. Then, the crude was purified (PE / AcOEt 7:3)

obtaining compound **14j** as yellow oil (74 mg, 17%). Then a mixture of **14j** (74 mg, 0.13 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (7 mg, 0.01 mmol) and ammonium formate (54 mg, 0.86 mmol) was stirred for 3 h at 80 °C. After work-up, the crude was treated with 1:1 pyridine / acetic anhydride (3 mL) and stirred for 18 h at r.t. After work-up and purification (PE / AcOEt from 3:2 to 1:1) pure compound **2j** was obtained as white foam (48 mg, 64%). R<sub>f</sub> = 0.50 (PE / AcOEt 3:7). δ<sub>H</sub>(300 MHz, CDCl<sub>3</sub>, 25 °C): 8.05 (1 H, s, NH), 7.55 (2 H, d, J 8.9), 7.36 (2 H, d, J 8.7), 7.35 (2 H, d, J 8.4), 7.12-6.96 (9 H, m), 6.89 (2 H, d, J 8.7), 6.28 (1 H, s, CH), 2.28 (6 H, s, CH<sub>3</sub>CO), 2.22 (3 H, s, CH<sub>3</sub>CO). δ<sub>C</sub>(75 MHz, CDCl<sub>3</sub>, 25 °C): 170.5, 169.5, 169.1, 168.7, 167.8 (C=O), 151.5, 150.8, 146.8, 140.6, 135.5, 132.9, 131.4 (quat.), 131.3 (x2), 130.4 (x2), 130.2 (x2), 128.6 (x2), 127.6, 121.9 (x2), 121.7 (x2), 120.9 (x2), 120.8 (x2) (ArCH), 66.6 (CH), 21.1 (CH<sub>3</sub>CO). IR: ν<sub>max</sub>/cm<sup>-1</sup> 3282, 3070, 2988, 1756, 1697, 1621, 1595, 1546, 1505, 1494, 1453, 1409, 1367, 1310, 1187, 1163, 1106, 1075, 1046, 1014, 966, 909, 846, 757, 737, 700, 675, 634, 611. m/z (ESI<sup>+</sup>): 581.1929 (M + H<sup>+</sup>). C<sub>33</sub>H<sub>29</sub>O<sub>8</sub>N<sub>2</sub> requires 581.1924.

**(R,S)-4-Acetoxy-N-(4-acetoxyphenyl)-N-(1-(4-(pivaloyloxy)phenyl)-2-((4-acetoxyphenyl)amino)-2-oxoethyl)benzamide 2k.** Following the general procedure B, a mixture of 4-pivaloyloxybenzaldehyde (203 mg, 1.00 mmol), 4-allyloxyaniline (153 μL, 1.11 mmol), 4-allyloxybenzoic acid (200 mg, 1.11 mmol), and 2,6-dimethylphenyl isocyanide (144 mg, 1.11 mmol) and 3 Å molecular sieves (50 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt 7:3) compound **15** was obtained pure as off-white foam (198 mg, 31%). Then a mixture of **15** (168 mg, 0.26 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (9 mg, 0.013 mmol) and ammonium formate (72 mg, 1.14 mmol) was stirred for 5 h at 80 °C. After work-up, the crude was treated with 1:1 pyridine / acetic anhydride (2.6 mL) and stirred for 18 h at r.t. After work-up and purification (PE / Et<sub>2</sub>O 1:3 + 1% EtOH) compound **2d** was obtained pure as white foam (117 mg, 69%). R<sub>f</sub> = 0.15 (PE / Et<sub>2</sub>O 1:3 + 1% EtOH). δ<sub>H</sub>(300 MHz, CDCl<sub>3</sub>, 25 °C): δ 7.42 (2 H, d, J 8.2), 7.42 (2 H, d, J 8.4), 7.26 (1 H, s, NH), 7.20-6.98 (7 H, m), 6.90 (2 H, d, J 8.3), 6.82 (2 H, d, J 8.5), 6.28 (1 H, s, CH), 2.23 (12 H, s, CH<sub>3</sub>CO and CH<sub>3</sub>Ar), 1.35 (9 H, s, C(CH<sub>3</sub>)<sub>3</sub>). δ<sub>H</sub>(75 MHz, CDCl<sub>3</sub>, 25 °C): 176.8, 170.3, 168.7 (x2), 167.9 (C=O), 151.5 (x2), 149.6, 138.4, 135.5 (x2), 133.4, 133.0, 131.6 (quat.), 131.4 (x4), 130.0 (x2), 128.2 (x2), 127.4, 121.9 (x2), 121.6 (x2), 120.9 (x2) (ArCH), 66.3 (CH), 39.1 (C(CH<sub>3</sub>)<sub>3</sub>), 27.1 (ArCH<sub>3</sub>), 21.1, 18.6 (CH<sub>3</sub>CO). IR: ν<sub>max</sub>/cm<sup>-1</sup> 3269, 3046, 2970, 2874, 1752, 1643, 1603, 1503, 1479, 1418, 1367, 1279, 1263, 1189, 1164, 1107, 1015, 945, 910, 850, 803, 765, 703, 678, 626. m/z (ESI<sup>+</sup>): 651.2713 (M + H<sup>+</sup>). C<sub>38</sub>H<sub>39</sub>O<sub>8</sub>N<sub>2</sub> requires 651.2706.

**(R,S)-(E)-N-(3-Acetoxybenzyl)-N-(1-(4-acetoxyphenyl)-2-((4-acetoxyphenyl)amino)-2-oxoethyl)-3-(4-acetoxy-3-methoxyphenyl)acrylamide 2l.** Following the general procedure B, a mixture of aldehyde **12** (404 mg, 2.49 mmol), 3-allyloxybenzylamine (406 mg, 2.49 mmol), acid **13** (530 mg, 2.26 mmol), 4-allyloxyphenyl isocyanide (397 mg, 2.49 mmol) and 3 Å molecular sieves (250 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt from 2:1 to 1:1) compound **14l** was obtained as yellow foam (1.19 g, 75%). Then a mixture of **14l** (200 mg, 0.29 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (16

mg, 0.02 mmol) and ammonium formate (161 mg, 2.56 mmol) was stirred for 4 h at 80 °C. After work-up, the crude was treated with 1:1 pyridine / acetic anhydride (4 mL) and stirred for 18 h at r.t. After work-up and purification (PE / AcOEt 3:4) compound **2l** was obtained pure as white foam (118 mg, 58%).  $R_f = 0.80$  (PE / AcOEt 1:6).  $\delta_H$ (300 MHz, CDCl<sub>3</sub>, 25 °C): 8.50 (1 H, s, NH), 7.67 (1 H, d, J 15.3, ArCH=CH), 7.48 (2 H, d, J 8.7), 7.42 (2 H, d, J 8.4), 7.20 (1 H, t, J 7.9), 7.03-6.80 (10 H, m), 6.67 (1 H, d, J 15.3, ArCH=CH), 6.30 (1 H, s, CH), 4.93 and 4.69 (2 H, AB syst., J 17.9, CH<sub>2</sub>Ar), 3.76 (3 H, s, OCH<sub>3</sub>), 2.30, 2.26, 2.24 (4 x 3 H, 4 s, CH<sub>3</sub>CO).  $\delta_C$ (75 MHz, CDCl<sub>3</sub>, 25 °C): 169.5, 169.2, 169.1, 168.8, 168.3, 167.9 (C=O), 151.2, 150.9 (x2), 146.9, 141.1, 139.5, 135.4, 133.7, 131.5, 144.2 (ArCH=CH), 131.0 (x2), 129.6, 123.6, 123.1, 122.0 (x2), 121.9 (x2), 120.9 (x3), 120.4, 119.6, 111.5 (ArCH), 117.6 (ArCH=CH), 63.0 (CH), 55.9 (OCH<sub>3</sub>), 49.6 (ArCH<sub>2</sub>), 21.10 (x2), 21.08, 20.6 (CH<sub>3</sub>CO). IR:  $\nu_{max}/cm^{-1}$  3285, 3072, 2940, 1758, 1694, 1647, 1602, 1545, 1505, 1452, 1408, 1368, 1302, 1258, 1185, 1161, 1122, 1013, 978, 955, 908, 831, 792, 730, 691, 637.  $m/z$  (ESI+): 709.2405 (M + H<sup>+</sup>). C<sub>39</sub>H<sub>37</sub>O<sub>11</sub>N<sub>2</sub> requires 709.2397.

**(R,S)-(E)-N-(2-(4-Acetoxyphenyl)ethyl)-N-(1-(4-acetoxyphenyl)-2-(methylamino)-2-oxoethyl)-3-(4-acetoxy-3-methoxyphenyl)acrylamide 2m.** Following the general procedure B, a mixture of aldehyde **12** (135 mg, 0.83 mmol), 4-allyloxyphenethylamine (148 mg, 0.83 mmol), acid **13** (176 mg, 0.75 mmol), methyl isocyanide (74  $\mu$ L, 1.23 mmol) and 3 Å molecular sieves (50 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt 1:2) compound **14m** was obtained as yellow foam (263 mg, 59%). Then a mixture of **14m** (237 mg, 0.40 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (23 mg, 0.03 mmol) and ammonium formate (170 mg, 2.70 mmol) was stirred for 4 h at 80 °C. After work-up, the crude was treated with 1:1 pyridine / acetic anhydride (9 mL) and stirred for 18 h at r.t. After work-up and purification (PE / AcOEt from 1:3 to 1:4) compound **2m** was obtained pure as white foam (188 mg, 78%).  $R_f = 0.32$  (PE / AcOEt 1:6).  $\delta_H$ (300 MHz, CDCl<sub>3</sub>, 50 °C): 7.68 (1 H, d, J 15.3, ArCH=CH), 7.56-7.45 (2 H, m), 7.13 (2 H, d, J 8.4), 7.11-6.89 (7 H, m), 6.74 (1 H, d, J 15.3, ArCH=CH), 6.21 (1 H, broad d, J 4.8, NH), 6.12 (1 H, s, CH), 3.84 (3 H, s, OCH<sub>3</sub>), 3.80-3.55 (2 H, broad m, CH<sub>2</sub>N), 2.87-2.73 (1 H, m, ArCHH), 2.83 (3 H, d, J 4.8, CH<sub>3</sub>NH), 2.45-2.17 (1 H, m, ArCH<sub>2</sub>), 2.30, 2.29, 2.24 (3 x 3 H, 3 s, CH<sub>3</sub>CO).  $\delta_C$ (75 MHz, CDCl<sub>3</sub>, 25 °C): 170.0, 169.2, 169.0, 168.6, 167.2 (C=O), 151.5, 151.0, 149.5, 141.3, 135.8, 134.1, 132.9 (quat.), 143.1 (ArCH=CH), 130.7 (x2), 129.5 (x2), 123.3, 122.1 (x2), 121.7 (x2), 120.6, 111.8 (ArCH), 117.8 (ArCH=CH), 61.6 (CH), 56.0 (OCH<sub>3</sub>), 48.0 (NCH<sub>2</sub>), 36.4 (ArCH<sub>2</sub>), 26.4 (CH<sub>3</sub>NH), 21.0 (x2), 20.5 (CH<sub>3</sub>CO). IR:  $\nu_{max}/cm^{-1}$  3311, 2941, 2249, 1759, 1674, 1647, 1601, 1506, 1466, 1450, 1416, 1369, 1262, 1190, 1153, 1121, 1032, 1013, 979, 908, 829, 726, 646, 623.  $m/z$  (ESI+): 603.2352 (M + H<sup>+</sup>). C<sub>33</sub>H<sub>35</sub>O<sub>9</sub>N<sub>2</sub> requires 603.2343.

**(R,S)-(E)-3-(4-Acetoxy-3-methoxyphenyl)-N-(1-(4-acetoxy-3-methoxyphenyl)-2-(tert-butylamino)-2-oxoethyl)-N-benzylacrylamide 2n.** Following the general procedure B, a mixture of 4-allyloxy-3-methoxybenzaldehyde (140 mg, 0.70 mmol), benzylamine (84  $\mu$ L, 0.77 mmol), acid **13** (180 mg, 0.77 mmol), *t*-butyl isocyanide (87  $\mu$ L, 0.77 mmol) and 3 Å

molecular sieves (35 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt 6:4) compound **14n** was obtained pure as pale yellow foam (249 mg, 63%). Then a mixture of **14n** (249 mg, 0.42 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (15 mg, 0.021 mmol) and ammonium formate (117 mg, 1.85 mmol) was stirred for 5 h at 80 °C. After work-up, the crude was treated with 1:1 pyridine / acetic anhydride (4.2 mL) and stirred for 18 h at r.t. After work-up and purification (PE / AcOEt 6:4 + 3% EtOH) compound **2n** was obtained pure as white foam (144 mg, 36%).  $R_f = 0.26$  (PE / AcOEt 6:4 + 3% EtOH).  $\delta_H$ (300 MHz, CDCl<sub>3</sub>, 25 °C): 7.71 (1 H, d, J 15.3, ArCH=CH), 7.45-7.10 (4 H, m), 7.10-6.90 (6 H, m), 6.83 (1 H, s), 6.65 (1 H, d, J 15.3, ArCH=CH), 6.12 (1 H, s, CH), 5.76 (1 H, s, NH), 4.92, 4.68 (2 H, AB syst., J 17.9, CH<sub>2</sub>Ph), 3.74, 3.67 (2 x 3 H, 2 s, OCH<sub>3</sub>), 2.29, 2.28 (2 x 3 H, 2 s, CH<sub>3</sub>CO), 1.37 (9 H, s, C(CH<sub>3</sub>)<sub>3</sub>).  $\delta_C$ (75 MHz, CDCl<sub>3</sub>, 25 °C):  $\delta$  168.8 (x2), 168.0 (C=O), 151.1, 151.0, 140.9, 139.8, 138.4, 134.1, 133.9 (quat.), 143.0 (ArCH=CH), 128.5 (x2), 127.0, 126.2 (x2), 123.0, 122.8, 122.0, 120.8, 114.1, 111.3 (ArCH), 118.6 (ArCH=CH), 62.2 (CH), 55.8 (OCH<sub>3</sub>), 51.8 (C(CH<sub>3</sub>)<sub>3</sub>), 49.6 (NCH<sub>2</sub>), 28.6 (C(CH<sub>3</sub>)<sub>3</sub>), 20.6 (CH<sub>3</sub>CO). IR:  $\nu_{max}/cm^{-1}$  3279, 3064, 2939, 1761, 1735, 1688, 1646, 1601, 1547, 1507, 1454, 1414, 1368, 1298, 1213, 1191, 1156, 1121, 1031, 1012, 976, 953, 909, 829, 723, 696, 635, 603.  $m/z$  (ESI+): 603.2697 (M + H<sup>+</sup>). C<sub>34</sub>H<sub>39</sub>O<sub>8</sub>N<sub>2</sub> requires 603.2706.

**(R,S)-(E)-3-(4-Acetoxy-3-methoxyphenyl)-N-(1-(4-acetoxyphenyl)-2-(methylamino)-2-oxoethyl)-N-benzylacrylamide 2o.** Following the general procedure B, a mixture of 4-allyloxybenzaldehyde (150 mg, 0.93 mmol), benzylamine (111  $\mu$ L, 1.02 mmol), acid **13** (238 mg, 1.02 mmol), methyl isocyanide (61  $\mu$ L, 1.02 mmol) and 3 Å molecular sieves (47 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt 4:6) compound **14o** was obtained pure as pale yellow foam (246 mg, 51%). Then a mixture of **14o** (229 mg, 0.43 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (15 mg, 0.022 mmol) and ammonium formate (120 mg, 1.91 mmol) was stirred for 5 h at 80 °C. After work-up, the crude was treated with 1:1 pyridine / acetic anhydride (4.3 mL) and stirred for 18 h at r.t. After work-up and purification (PE / AcOEt 4:6 + 3% EtOH) compound **2o** was obtained pure as white foam (168 mg, 74%).  $R_f = 0.40$  (PE / AcOEt 4:6 + 3% EtOH).  $\delta_H$ (300 MHz, CDCl<sub>3</sub>, 25 °C): 7.70 (1 H, d, J 15.3, ArCH=CH), 7.43 (2 H, d, J 8.4), 7.27-7.13 (3 H, m), 7.09 (2 H, broad d, J 7.2), 7.05-6.93 (4 H, m), 6.84 (1 H, s), 6.63 (1 H, d, J 15.3, ArCH=CH), 5.99 (1 H, s, CH), 5.94 (1 H, s, NH), 4.89, 4.66 (2 H, AB syst., J 17.8, CH<sub>2</sub>Ph), 3.75 (3 H, s, OCH<sub>3</sub>), 2.84 (3 H, d, J 4.8, CH<sub>3</sub>NH), 2.29, 2.28 (2 x 3 H, 2 s, CH<sub>3</sub>CO).  $\delta_C$ (75 MHz, CDCl<sub>3</sub>, 25 °C): 169.9, 169.2, 168.8, 168.0 (C=O), 151.2, 150.7, 141.0, 137.7, 134.0, 132.5 (quat.), 143.3 (ArCH=CH), 130.9 (x2), 128.6 (x2), 127.2, 126.2 (x2), 123.1, 121.9 (x2), 120.8, 111.3 (ArCH), 118.3 (ArCH=CH), 62.5 (CH), 55.8 (OCH<sub>3</sub>), 50.2 (NCH<sub>2</sub>), 26.5 (NCH<sub>3</sub>), 21.1, 20.6 (CH<sub>3</sub>CO). IR:  $\nu_{max}/cm^{-1}$  3302, 3065, 2940, 1760, 1674, 1647, 1601, 1506, 1453, 1407, 1368, 1300, 1257, 1189, 1155, 1121, 1080, 1030, 1012, 976, 957, 907, 844, 829, 724, 697, 676, 635.  $m/z$  (ESI+): 531.2137 (M + H<sup>+</sup>). C<sub>30</sub>H<sub>31</sub>O<sub>7</sub>N<sub>2</sub> requires 531.2131.



**(*R,S*)-(E)-*N*-2-((4-(2-(2-acetoxyethoxy)ethoxy)phenyl)amino)-1-(4-acetoxyphenyl)-2-oxoethyl)-3-(4-acetoxy-3-methoxyphenyl)-*N*-benzylacrylamide 2p.**

Following the general procedure B, a mixture of 4-allyloxybenzaldehyde (146 mg, 0.90 mmol), benzylamine (106  $\mu$ L, 0.99 mmol), acid **13** (231 mg, 0.99 mmol), *N*-(4-(2-(2-(allyloxy)ethoxy)ethoxy)phenyl) isocyanide (244 mg, 0.99 mmol) and 3 Å molecular sieves (45 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt 1:1) compound **14p** was obtained pure as pale yellow foam (449 mg, 71%). Then a mixture of **14p** (449 mg, 0.61 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (32 mg, 0.046 mmol) and ammonium formate (254 mg, 4.03 mmol) was stirred for 5 h at 80 °C. After work-up, the crude was treated with 1:1 pyridine / acetic anhydride (6.1 mL) and stirred for 18 h at r.t. After work-up and purification (PE / AcOEt 1:1 + 3% EtOH) compound **2p** was obtained pure as white foam (271 mg, 60%). *R<sub>f</sub>* = 0.40 (PE / AcOEt 4:6 + 3% EtOH).  $\delta_{\text{H}}$ (300 MHz, CDCl<sub>3</sub>, 25 °C):  $\delta$  7.90 (1 H, s, NH), 7.71 (1 H, d, J 15.2, ArCH=CH), 7.49 (2 H, d, J 8.2), 7.39 (2 H, d, J 8.7), 7.30-7.07 (5 H, m), 7.03 (2 H, d, J 8.3), 6.96 (2 H, s), 6.90-6.81 (3 H, m), 6.67 (1 H, d, J 15.3, ArCH=CH), 6.19 (1 H, s, CH), 4.92, 4.72 (2 H, AB syst., J 17.6, CH<sub>2</sub>Ph), 4.25 (2 H, dd, J 4.5, 5.0, OCH<sub>2</sub>), 4.10 (2 H, broad t, J 4.5, CH<sub>2</sub>O), 3.83 (2 H, broad t, J 4.5, CH<sub>2</sub>O), 3.76 (2 H, dd, J 4.5, 5.0, OCH<sub>2</sub>), 3.75 (3 H, s, OCH<sub>3</sub>), 2.29, 2.28, 2.07 (3 x 3 H, 3 s, CH<sub>3</sub>CO).  $\delta_{\text{C}}$ (75 MHz, CDCl<sub>3</sub>, 25 °C):  $\delta$  171.1, 169.1, 168.3, 167.6 (C=O), 155.6, 151.2, 150.8, 141.0, 137.7, 133.9, 135.5, 132.0 (quat.), 143.6 (ArCH=CH), 130.8 (x2), 128.7 (x2), 127.3, 126.3 (x2), 123.1, 122.0 (x2), 121.8 (x2), 120.8, 114.9 (x2), 111.4 (ArCH), 118.1 (ArCH=CH), 69.7, 69.3, 67.7, 63.6 (CH<sub>2</sub>O), 63.0 (CH), 55.8 (OCH<sub>3</sub>), 50.2 (NCH<sub>2</sub>), 21.1, 21.0, 20.6 (CH<sub>3</sub>CO). IR:  $\nu_{\text{max}}$ /cm<sup>-1</sup> 3278, 3065, 2940, 1760, 1735, 1688, 1647, 1601, 1546, 1507, 1454, 1414, 1368, 1299, 1191, 1157, 1121, 1031, 1012, 975, 953, 909, 829, 723, 697, 636, 604. *m/z* (ESI+): 739.2875 (M + H<sup>+</sup>). C<sub>41</sub>H<sub>43</sub>O<sub>11</sub>N<sub>2</sub> requires 739.2867.

**(*R,S*)-(E)-*N*-(3-acetoxybenzyl)-3-(4-acetoxy-3-methoxyphenyl)-*N*-(1-(4-acetoxyphenyl)-2-(methylamino)-2-oxoethyl)-acrylamide 2q.**

Following the general procedure B, a mixture of aldehyde **12** (135 mg, 0.83 mmol), 3-allyloxybenzylamine (135 mg, 0.83 mmol), acid **13** (176 mg, 0.75 mmol), methyl isocyanide (49  $\mu$ L, 0.83 mmol) and 3 Å molecular sieves (50 mg) was stirred for 3 days at r.t. After work-up and purification (PE / AcOEt 3:7) compound **14q** was obtained as white foam (313 mg, 72%). Then a mixture of **14q** (97 mg, 0.17 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (10 mg, 0.014 mmol) and ammonium formate (71 mg, 1.12 mmol) was stirred for 3 h at 80 °C. After work-up, the crude was treated with 1:1 pyridine / acetic anhydride (2.5 mL) and stirred for 18 h at r.t. After work-up and purification (PE / AcOEt 1:6) compound **2q** was obtained pure as white foam (73 mg, 75%). *R<sub>f</sub>* = 0.49 (PE / AcOEt 1:6).  $\delta_{\text{H}}$ (300 MHz, CDCl<sub>3</sub>, 25 °C): 7.72 (1 H, d, J 15.3, ArCH=CH), 7.40 (2 H, d, J 8.1), 7.21 (1 H, t, J 7.8), 7.03-6.82 (8 H, m), 6.63 (1 H, d, J 15.3, ArCH=CH), 5.97 (1 H, s, CH), 5.94 (1 H, broad s, NH), 4.89, 4.65 (2 H, AB syst., J 17.9, CH<sub>2</sub>Ar), 3.77 (3 H, s, OCH<sub>3</sub>), 2.83 (3 H, d, J 4.6, CH<sub>3</sub>N), 2.29, 2.27, 2.26 (3 x 3 H, 3 s, CH<sub>3</sub>CO).  $\delta_{\text{C}}$ (75 MHz, CDCl<sub>3</sub>, 25 °C): 169.9, 169.21, 169.16, 168.8, 168.0 (C=O), 151.2, 150.9, 150.8, 141.0, 139.6, 133.9, 132.2 (quat.), 143.8 (ArCH=CH), 131.0 (x2),

129.6, 123.6, 123.1, 121.9 (x2), 120.9, 120.4, 119.6, 111.4 (ArCH), 117.9 (ArCH=CH), 62.5 (CH), 55.9 (OCH<sub>3</sub>), 49.7 (NCH<sub>2</sub>), 26.4 (CH<sub>3</sub>N), 21.1 (x2), 20.6 (CH<sub>3</sub>CO). IR:  $\nu_{\text{max}}$ /cm<sup>-1</sup> 3302, 3072, 2967, 2940, 1760, 1676, 1648, 1602, 1506, 1438, 1414, 1368, 1302, 1258, 1190, 1156, 1120, 1014, 975, 909, 829, 794, 751, 722, 694, 635. *m/z* (ESI+): 589.2200 (M + H<sup>+</sup>). C<sub>32</sub>H<sub>33</sub>O<sub>9</sub>N<sub>2</sub> requires 589.2186.

**General procedure for the preparation of polyphenols 1d,f,g,h,j,k,l,m,n,o,p,q from the corresponding acetylated derivatives 2.** The peracylated Ugi product was treated with 0.2 M MeONa in MeOH (freshly prepared by adding Na to dry MeOH) under N<sub>2</sub> atmosphere. After stirring for 1 h (18 h for compound **15**) at r.t., the mixture was treated with previously washed Amberlyst 15 acid resin until pH = 4. The resin was filtered off and the solution evaporated to dryness (with exception of compound **1n**). The resulting polyphenols were not fully characterized, but only examined at <sup>1</sup>H NMR and HPLC in order to establish their degree of purity.

**(*R,S*)-(E)-*N*-Butyl-*N*-(2-(tert-butylamino)-1-(4-hydroxyphenyl)-2-oxoethyl)-3-(4-hydroxy-3-methoxyphenyl)acrylamide 1d.**

$\delta_{\text{H}}$ (300 MHz, DMSO-*d*<sub>6</sub>, 90 °C) (Note: the 2 phenolic OH exchange with H<sub>2</sub>O contained in the solvent giving a broad signal around 9 ppm): 7.42 (1 H, d, J 15.2, ArCH=CH), 7.35 (1 H, s, NH), 7.17 (1 H, d, J 1.8), 7.14 (2 H, d, J 8.7), 7.06 (1 H, dd, J 8.2, 1.8), 6.82 (1 H, d, J 15.2, ArCH=CH), 6.81 (1 H, d, J 8.2), 6.79 (2 H, d, J 8.7), 5.87 (1 H, s, CH), 3.83 (3 H, s, OCH<sub>3</sub>), 3.50-3.25 (mc = 3.38) (2 H, m, CH<sub>2</sub>N), 1.50-1.30 (1 H, m, CHH), 1.29 (9 H, s, (CH<sub>3</sub>)<sub>3</sub>C), 1.20-0.95 (3 H, m, CH<sub>2</sub> and CHH), 0.74 (3 H, t, J 7.2, CH<sub>3</sub>CH<sub>2</sub>). HPLC (see supplementary information) showed a purity of 92%.

**(*R,S*)-(E)-*N*-Benzyl-*N*-(2-(tert-butylamino)-1-(4-hydroxyphenyl)-2-oxoethyl)-3-(3,4-dihydroxyphenyl)acrylamide 1f.**

$\delta_{\text{H}}$ (300 MHz, DMSO-*d*<sub>6</sub>, 70 °C) (Note: the 2 phenolic OH exchange with H<sub>2</sub>O contained in the solvent giving a very broad signal around 9 ppm): 7.53 (1 H, s, NH), 7.35 (1 H, d, J 15.2, ArCH=CH), 7.20-7.00 (6 H, m, ArCH), 6.95-6.75 (2 H, m, ArCH), 6.73-6.62 (4 H, m, ArCH=CH and ArCH), 6.01 (1 H, s, CH), 4.85 (1 H, d, J 16.9, CHHPh), 4.55 (1 H, d, J 16.9, CHHPh), 1.24 (9 H, s, (CH<sub>3</sub>)<sub>3</sub>C). HPLC (see supplementary information) showed a purity of 97%.

**(*R,S*)-*N*-(4-Hydroxyphenyl)-*N*-(1-(4-hydroxyphenyl)-2-((4-hydroxyphenyl)amino)-2-oxoethyl)benzamide 1g.**

$\delta_{\text{H}}$ (300 MHz, DMSO-*d*<sub>6</sub>, 70 °C) (Note: the 3 phenolic OH exchange with H<sub>2</sub>O contained in the solvent giving a very broad signal not visible in the spectrum): 9.69 (1 H, s, NH), 7.39 (2 H, d, J 8.8), 7.25-7.12 (5 H, m), 6.96 (2 H, d, J 8.5), 6.81 (2 H, broad d, J 7.2), 6.70 (2 H, d, J 9.0), 6.57 (2 H, d, J 8.7), 6.33 (2 H, d, J 8.9), 6.21 (1 H, s, CH). HPLC (see supplementary information) showed a purity of 99%.

**(*R,S*)-4-Hydroxy-*N*-(1-(4-hydroxyphenyl)-2-((4-hydroxyphenyl)amino)-2-oxoethyl)-*N*-phenylbenzamide 1j.**

$\delta_{\text{H}}$ (300 MHz, DMSO-*d*<sub>6</sub>, 30 °C) (Note: the 3 phenolic OH exchange with H<sub>2</sub>O contained in the solvent giving a very broad signal not visible in the spectrum):  $\delta$  9.91 (1 H, s, NH), 7.40 (2 H, d, J 8.9), 7.04 (2 H, d, J 8.7), 6.99 (5 H, broad s), 6.93 (2 H, d, J 8.4), 6.69 (2 H, d, J 8.9), 6.53 (2 H, d, J 8.4), 6.49 (2 H,

d, J 8.7), 6.22 (1 H, s, CH). HPLC (see supplementary information) showed a purity of 100%.

**(R,S)-4-Hydroxy-N-(4-hydroxyphenyl)-N-(1-(4-hydroxyphenyl)-2-((4-hydroxyphenyl)amino)-2-oxoethyl)benzamide 1k.**  $\delta_{\text{H}}$ (300 MHz, DMSO-*d*6, 70 °C):  $\delta$  9.39, 9.14, 9.03 (3 x 1 H, 3 broad s, OH), 9.21 (1 H, s, NH), 7.08 (2 H, d, J 8.7), 7.03 (2 H, d, J 8.4), 7.03 (3 H, s), 6.77 (2 H, d, J 8.2), 6.59 (2 H, d, J 8.5), 6.52 (2 H, d, J 8.6), 6.36 (2 H, d, J 8.8), 6.23 (1 H, s, CH), 2.12 (6 H, s, CH<sub>3</sub>Ar), 1.35 (9 H, s, C(CH<sub>3</sub>)<sub>3</sub>). HPLC (see supplementary information) showed a purity of 100%.

**(R,S)-(E)-N-(3-Hydroxybenzyl)-N-(1-(4-hydroxyphenyl)-2-((4-hydroxyphenyl)amino)-2-oxoethyl)-3-(4-hydroxy-3-methoxyphenyl)acrylamide 1l.**  $\delta_{\text{H}}$ (300 MHz, DMSO-*d*6, 90 °C) (Note: the 4 phenolic OH exchange with H<sub>2</sub>O contained in the solvent giving a very broad signal at around 9 ppm): 9.63 (1 H, s, NH), 7.41 (1 H, d, J 15.3, ArCH=CH), 7.33 (2 H, d, J 8.7), 7.17 (2 H, d, J 8.5), 7.00-6.85 (3 H, m), 6.75 (2 H, d, J 8.1), 6.73-6.63 (4 H, m), 6.58 (1 H, s), 6.55-6.48 (2 H, m), 6.17 (1 H, s, CH), 4.79 and 4.53 (2 H, AB syst., J 17.1, CH<sub>2</sub>Ar), 3.76 (3 H, s, OCH<sub>3</sub>). HPLC (see supplementary information) showed a purity of 92%.

**(R,S)-(E)-N-(2-(4-hydroxyphenyl)ethyl)-N-(1-(4-hydroxyphenyl)-2-(methylamino)-2-oxoethyl)-3-(4-hydroxy-3-methoxyphenyl)acrylamide 1m.**  $\delta_{\text{H}}$ (300 MHz, DMSO-*d*6, 90 °C) (Note: the 4 phenolic OH exchange with H<sub>2</sub>O contained in the solvent giving a very broad signal at around 6 ppm): 7.73 (1 H, broad s, NH), 7.44 (1 H, d, J 15.3, ArCH=CH), 7.21-7.14 (3 H, m), 7.05 (1 H, dd, J 8.2, 1.9), 6.87-6.74 (6 H, m), 6.63 (2 H, d, J 8.4), 6.01 (1 H, s, CH), 3.85 (3 H, s, OCH<sub>3</sub>), 3.64-3.40 (2 H, m, CH<sub>2</sub>N), 2.66 (3 H, d, J 4.5, CH<sub>3</sub>NH), 2.63-2.48 (1 H, m, ArCHH), 2.18-2.03 (1 H, m, ArCH<sub>2</sub>). HPLC (see supplementary information) showed a purity of 92%.

**(R,S)-(E)-N-Benzyl-3-(4-hydroxy-3-methoxyphenyl)-N-(2-(tert-butylamino)-1-(4-hydroxy-3-methoxyphenyl)-2-oxoethyl)acrylamide 1n.** After the treatment with the resin and evaporation, in this case the residue was further purified by chromatography (PE / AcOEt 6:4).  $\delta_{\text{H}}$ (300 MHz, DMSO-*d*6, 90 °C): 9.00 (1 H, s, OH), 8.57 (1 H, s, OH), 7.45 (1 H, broad s, NH), 7.42 (1 H, d, J 15.3, ArCH=CH), 7.21-6.99 (6 H, m), 6.94 (1 H, d, J 8.0), 6.84-6.66 (4 H, m), 6.70 (1 H, d, J 15.3, ArCH=CH), 5.97 (1 H, s, CH), 4.87, 4.55 (2 H, AB syst., J 16.7, PhCH<sub>2</sub>), 3.78 (3 H, s, OCH<sub>3</sub>), 3.63 (3 H, s, OCH<sub>3</sub>), 1.27 (9 H, s, (CH<sub>3</sub>)<sub>3</sub>C). HPLC (see supplementary information) showed a purity of 99%.

**(R,S)-(E)-N-Benzyl-3-(4-hydroxy-3-methoxyphenyl)-N-(1-(4-hydroxyphenyl)-2-(methylamino)-2-oxoethyl)-acrylamide 1o.**  $\delta_{\text{H}}$ (300 MHz, DMSO-*d*6, 90 °C): 9.09, 9.00 (2 x 1 H, 2 broad s, OH), 7.73 (1 H, s, NH), 7.40 (1 H, d, J 15.3, ArCH=CH<sub>2</sub>), 7.21-7.02 (6 H, m), 6.96 (1 H, s), 6.90 (1 H, d, J 8.1), 6.75 (2 H, d, J 8.4), 6.67 (2 H, d, J 8.7), 6.66 (1 H, broad signal, ArCH=CH), 6.04 (1 H, s, CH), 4.85, 4.62 (2 H, AB syst., J 17.1, CH<sub>2</sub>Ph), 3.77 (3 H, s, OCH<sub>3</sub>), 2.63 (3 H, d, J 4.6, CH<sub>3</sub>NH). HPLC (see supplementary information) showed a purity of 96%.

**(R,S)-(E)-N-Benzyl-N-(2-((4-(2-(2-hydroxyethoxy)ethoxy)phenyl)amino)-1-(4-hydroxyphenyl)-2-oxoethyl)-3-(4-hydroxy-3-methoxyphenyl)acrylamide 1p.**  $\delta_{\text{H}}$ (300 MHz, DMSO-*d*6, 90 °C): 9.80 (1 H, s, NH), 9.15, 9.01 (2 x 1 H, 2 broad s, OH), 7.46 (2 H, d, J 9.0), 7.43 (1 H, d, J 15.0, ArCH=CH), 7.22-7.06 (4 H, m), 6.94 (1 H, broad d, J 15.0,

ArCH=CH), 6.88 (2 H, d, J 9.0), 6.75 (1 H, d, J 8.1), 6.69 (2 H, d, J 8.4), 6.21 (1 H, s, CH), 4.90, 4.63 (2 H, AB syst., J 17.1, CH<sub>2</sub>Ph), 4.08 (2 H, t, J 5.0, OCH<sub>2</sub>), 3.75 (2 H, t, J 4.8, CH<sub>2</sub>O), 3.75 (3 H, s, OCH<sub>3</sub>), 3.58-3.48 (4 H, m, CH<sub>2</sub>O). HPLC (see supplementary information) showed a purity of 97.5%.

**(R,S)-(E)-N-(3-hydroxybenzyl)-3-(4-hydroxy-3-methoxyphenyl)-N-(1-(4-hydroxyphenyl)-2-(methylamino)-2-oxoethyl)acrylamide 1q.**  $\delta_{\text{H}}$ (300 MHz, DMSO-*d*6, 90 °C) (Note: the 4 phenolic OH exchange with H<sub>2</sub>O contained in the solvent giving a very broad signal at around 9 ppm): 7.71 (1 H, s, NH), 7.39 (1 H, d, J 15.3, ArCH=CH), 7.11 (2 H, d, J 8.1), 7.00-6.87 (3 H, m), 6.75 (1 H, d, J 8.1), 6.68 (2 H, d, J 8.7), 6.67-6.46 (4 H, m), 6.02 (1 H, s, CH), 4.76, 4.53 (2 H, AB syst., J 17.1, CH<sub>2</sub>Ar), 3.77 (3 H, s, OCH<sub>3</sub>), 2.63 (3 H, d, J 4.5, CH<sub>3</sub>N). HPLC (see supplementary information) showed a purity of 99%.

### Biophysical and Biochemical tests

#### UV Spectroscopy

Synthetic polyphenols stock solutions were obtained by dissolving the compounds in 100% dimethyl sulfoxide (DMSO; Sigma) at given concentration (1.25 – 50 mM). Work solutions were prepared diluting the appropriate stock solution in PBS (150 mM, pH 7.4) at 12.5 – 500  $\mu$ M, in such a manner that each tube contained 1% of stock solution in DMSO. Solubility and turbidity of polyphenols in function of the concentration was determined by spectrophotometric measures using Shimadzu UV-2700 Spectrophotometer and reading the absorbance respectively at characteristic wavelength of each polyphenols and at  $\lambda = 405$  nm.

#### A $\beta$ sample preparation

One milliliter of DMSO was added to 1 mg of lyophilized synthetic peptide (A $\beta$ 1-42, A $\beta$ pE3-42 AnaSpec), reaching a final concentration of 1 mg/mL. Aliquots of 75  $\mu$ L were lyophilized and stored at -20°C until used. For all experiments, stock peptides were reconstituted as reported.<sup>54</sup> The concentration of the peptide in the stock solution was estimated using a molar extinction coefficient at 214 nm, by Shimadzu UV-2700 Spectrophotometer.<sup>55</sup>

For the preparation of the working samples, stock solution of each peptide was divided in two or more aliquots. One was diluted to 5  $\mu$ M in PBS containing 1% (v/v) DMSO to have a reference sample, the others were diluted in PBS containing the appropriate quantity of polyphenols stock solution in DMSO in such a manner that each samples contains 1% of DMSO. Final pH was measured and eventually corrected at 7.4 with few  $\mu$ L of 1M HCl.

#### Thioflavine T Fluorescence Spectroscopy

A $\beta$  peptides (5  $\mu$ M) were incubated at 37 °C in presence /absence of polyphenols as previous described and analyzed in parallel. ThT fluorescence was followed in time during aggregation. For this purpose, 47.5  $\mu$ L of A $\beta$  with and without test compounds, were mixed with 2.5  $\mu$ L ThT (400  $\mu$ M) in a 3 mm path length fluorescence cuvette. ThT fluorescence was measured by Luminescence Spectrometer Perkin Elmer LS50B at excitation and emission wavelengths of 440 nm (slit width=5 nm) and 482 nm (slit width = 10 nm), respectively.

#### Transmission electron microscopy (TEM)

## ARTICLE

## Organic and Biomolecular Chemistry

A $\beta$  peptides (10  $\mu$ M) were separately co-incubated with single polyphenol at molar ratio 1:5 (A $\beta$ :polyphenol) in sterile microtubes. To evaluate the morphology and the sizes of the species in the different samples, 5  $\mu$ L of each one were adsorbed for 5 min onto carbon coated 300-mesh copper grids. The aggregates species were negatively stained for 1 min with 5  $\mu$ L of 1% Uranyl Acetate. All air-dried specimens were examined with a Zeiss LEO 900 electron microscope (Zeiss, Stuttgart, Germany) operating at 80 kV. Images flattening and analysis was performed by ImageJ software.

## Conflicts of interest

There are no conflicts to declare



## Acknowledgements

This study is supported by Fondazione Cariplo, under "Integrated Biotechnology and Bioeconomy" programme. We thank Valeria Rocca for HPLC analyses, Giuliana Ottonello for HRMS, Giulia Baruzzo, Gianluca Pusceddu, and Riccardo Loddo for their experimental contribution to this work.


## References

1. A. J. Doig and P. Derreumaux, *Curr. Opin. Struct. Biol.*, 2015, **30**, 50-56.
2. Q. Nie, X. G. Du and M. Y. Geng, *Acta Pharm. Sinica*, 2011, **32**, 545-551.
3. F. Belluti, A. Rampa, S. Gobbi and A. Bisi, *Expert Opin. Ther. Pat.*, 2013, **23**, 581-596.
4. C. Ran, X. Xu, S. B. Raymond, B. J. Ferrara, K. Neal, B. J. Bacskai, Z. Medarova and A. Moore, *J. Amer. Chem. Soc.*, 2009, **131**, 15257-15261.
5. Y. Porat, A. Abramowitz and E. Gazit, *Chem. Biol. Drug Des.*, 2006, **67**, 27-37.
6. M. Hirohata, K. Hasegawa, S. Tsutsumi-Yasuhara, Y. Ohhashi, T. Ookoshi, K. Ono, M. Yamada and H. Naiki, *Biochemistry*, 2007, **46**, 1888-1899.
7. V. L. N. Ngoungoure, J. Schluesener, P. F. Moundipa and H. Schluesener, *Mol. Nutr. Food Res.*, 2015, **59**, 8-20.
8. N. Ferreira, I. Cardoso, M. R. Domingues, R. Vitorino, M. Bastos, G. Bai, M. J. Saraiva and M. R. Almeida, *FEBS Lett.*, 2009, **583**, 3569-3576.
9. M. Yamada, K. Ono, T. Hamaguchi and M. Noguchi-Shinohara, in *Natural Compounds as Therapeutic Agents for Amyloidogenic Diseases*, ed. N. Vassallo, Springer International Publishing, Cham, 2015, pp. 79-94.
10. K. M. Pate, M. Rogers, J. W. Reed, N. van der Munnik, S. Z. Vance and M. A. Moss, *Cns Neurosci. Therap.*, 2017, **23**, 135-144.
11. R. Randino, M. Grimaldi, M. Persico, A. De Santis, E. Cini, W. Cabri, A. Riva, G. D'Errico, C. Fattorusso, A. M. D'Ursi and M. Rodriguez, *Sci Rep*, 2016, **6**.
12. T. T. An, S. Feng and C. M. Zeng, *Redox Biol.*, 2017, **11**, 315-321.
13. J. M. Ringman, S. A. Frautschy, E. Teng, A. N. Begum, J. Bardens, M. Beigi, K. H. Gylys, V. Badmaev, D. D. Heath, L. G. Apostolova, V. Porter, Z. Vanek, G. A. Marshall, G. Hellemann, C. Sugar, D. L. Masterman, T. J. Montine, J. L. Cummings and G. M. Cole, *Alzheimer's Res. Ther.*, 2012, **4**, 43.
14. S. Feng, X.-H. Song and C.-M. Zeng, *FEBS Lett.*, 2012, **586**, 3951-3955.
15. S. Montanari, M. Bartolini, P. Neviani, F. Belluti, S. Gobbi, L. Pruccoli, A. Tarozzi, F. Falchi, V. Andrisano, P. Misztka, A. Cavalli, S. Filipek, A. Bisi and A. Rampa, *ChemMedChem*, 2016, **11**, 1296-1308.
16. L. A. Wessjohann, G. Kaluderovic, R. A. W. Neves Filho, M. C. Morejon, G. Lemanski and T. Ziegler, in *Science of Synthesis: Multicomponent Reactions, Vol. 1*, ed. T. J. J. Müller, 2013, pp. 415-497.
17. L. Banfi, R. Riva and A. Basso, *Synlett*, 2010, 23-41.
18. M. Spallarossa, L. Banfi, A. Basso, L. Moni and R. Riva, *Adv. Synth. Cat.*, 2016, **358**, 2940-2948.
19. L. Moni, C. F. Gers-Panther, M. Anselmo, T. J. J. Muller and R. Riva, *Chem. Eur. J.*, 2016, **22**, 2020-2031.
20. S. Caputo, A. Basso, L. Moni, R. Riva, V. Rocca and L. Banfi, *Beilstein J. Org. Chem.*, 2016, **12**, 139-143.
21. E. Tassano, A. Alama, A. Basso, G. Dondo, A. Galatini, R. Riva and L. Banfi, *Eur. J. Org. Chem.*, 2015, 6710-6726.
22. D. Goyal, S. Shuaib, S. Mann and B. Goyal, *Acs Combi. Sci.*, 2017, **19**, 55-80.
23. E. Elhalem, B. N. Bailey, R. Docampo, I. Ujváry, S. H. Szajnman and J. B. Rodriguez, *J. Med. Chem.*, 2002, **45**, 3984-3999.
24. S. W. Youn and E. M. Lee, *Org. Lett.*, 2016, **18**, 5728-5731.
25. B. Schmidt, N. Elizarov, R. Berger and F. Holter, *Org. Biomol. Chem.*, 2013, **11**, 3674-3691.
26. C. D'Arrigo, M. Tabaton and A. Perico, *Biopolymers*, 2009, **91**, 861-873.
27. C. Dammers, M. Schwarten, A. K. Buell and D. Willbold, *Chem. Sci.*, 2017, **8**, 4996-5004.
28. S. Schilling, T. Hoffmann, S. Manhart, M. Hoffmann and H. U. Demuth, *FEBS Lett.*, 2004, **563**, 191-196.
29. S. Schilling, T. Lauber, M. Schaupp, S. Manhart, E. Scheel, G. Bohm and H. U. Demuth, *Biochemistry*, 2006, **45**, 12393-12399.
30. D. Galante, F. S. Ruggeri, G. Dietler, F. Pellistri, E. Gatta, A. Corsaro, T. Florio, A. Perico and C. D'Arrigo, *Int. J. Biochem. Cell Biol.*, 2016, **79**, 261-270.
31. Y. Liu, S. H. Wang, S. Z. Dong, P. Chang and Z. F. Jiang, *Rsc Adv.*, 2015, **5**, 62402-62413.
32. F. Ding, J. M. Borreguero, S. V. Buldyrey, H. E. Stanley and N. V. Dokholyan, *Proteins*, 2003, **53**, 220-228.
33. A. Sgarbossa, D. Giacomazza and M. di Carlo, *Nutrients*, 2015, **7**, 5764-5782.
34. S. Sinha, D. H. J. Lopes and G. Bitan, *Acs Chem. Neurosci.*, 2012, **3**, 473-481.
35. E. Gazit, *Faseb J.*, 2002, **16**, 77-83.
36. C. Soto, E. M. Sigurdsson, L. Morelli, R. A. Kumar, E. M. Castano and B. Frangione, *Nat. Med.*, 1998, **4**, 822-826.


37. M. Convertino, R. Pellarin, M. Catto, A. Carotti and A. Caflich, *Protein Sci.*, 2009, **18**, 792-800.
38. M. Benchekroun, A. Romero, J. Egea, R. Leon, P. Michalska, I. Buendia, M. L. Jimeno, D. Jun, J. Janockova, V. Sepsova, O. Soukup, O. M. Bautista-Aguilera, B. Refouvelet, O. Ouari, J. Marco-Contelles and L. Ismaili, *J. Med. Chem.*, 2016, **59**, 9967-9973.
39. H. Josien, S. B. Ko, D. Bom and D. P. Curran, *Chem. Eur. J.*, 1998, **4**, 67-83.
40. M. Suzuki, J. F. K. Kotyk, S. I. Khan and Y. Rubin, *J. Am. Chem. Soc.*, 2016, **138**, 5939-5956.
41. W. Gu and R. B. Silverman, *Org. Lett.*, 2003, **5**, 415-418.
42. S. Imai and H. Togo, *Tetrahedron*, 2016, **72**, 6948-6954.
43. A. J. Harnoy, G. Slor, E. Tirosh and R. J. Amir, *Org. Biomol. Chem.*, 2016, **14**, 5813-5819.
44. J. Hoffmann and U. Kazmaier, *Angew. Chem., Int. Ed.*, 2014, **53**, 11356-11360.
45. R. Jaiswal, M. H. Dickman and N. Kuhnert, *Org. Biomol. Chem.*, 2012, **10**, 5266-5277.
46. L. Jiménez-González, S. García-Muñoz, M. Álvarez-Corral, M. Muñoz-Dorado and I. Rodríguez-García, *Chem. Eur. J.*, 2007, **13**, 557-568.
47. M. Kim, W. B. Euler and W. Rosen, *J. Org. Chem.*, 1997, **62**, 3766-3769.
48. J. Liu, K. Wu, T. Shen, Y. Liang, M. Zou, Y. Zhu, X. Li, X. Li and N. Jiao, *Chem. Eur. J.*, 2017, **23**, 563-567.
49. A. Massink, J. Louvel, I. Adlere, C. van Veen, B. J. H. Huisman, G. S. Dijksteel, D. Guo, E. B. Lenselink, B. J. Buckley, H. Matthews, M. Ranson, M. Kelso and A. P. Ijzerman, *J. Med. Chem.*, 2016, **59**, 4769-4777.
50. T. Mäkelä, J. Matikainen, K. Wähälä and T. Hase, *Tetrahedron*, 2000, **56**, 1873-1882.
51. C. D. Selassie, C. Hansch, T. A. Khwaja, C. B. Dias and S. Pentecost, *J. Med. Chem.*, 1984, **27**, 347-357.
52. G. H. Lee, E. B. Choi, E. Lee and C. S. Pak, *J. Org. Chem.*, 1994, **59**, 1428-1443.
53. T. Ventrice, E. M. Campi, W. R. Jackson and A. F. Patti, *Tetrahedron*, 2001, **57**, 7557-7574.
54. D. Galante, A. Corsaro, T. Florio, S. Vella, A. Pagano, F. Sbrana, M. Vassalli, A. Perico and C. D'Arrigo, *Int. J. Biochem. Cell Biol.*, 2012, **44**, 2085-2093.
55. L. W. Hung, G. D. Ciccotosto, E. Giannakis, D. J. Tew, K. Perez, C. L. Masters, R. Cappai, J. D. Wade and K. J. Barnham, *J. Neurosci.*, 2008, **28**, 11950-11958.




108x139mm (600 x 600 DPI)



202x83mm (200 x 200 DPI)



151x107mm (200 x 200 DPI)



145x94mm (200 x 200 DPI)





## SUPPORTING INFORMATION

**MULTICOMPONENT, FRAGMENT-BASED, SYNTHESIS OF NEW NATURAL-BASED POLYPHENOLS AND THEIR INHIBITING ACTIVITY ON BETA-AMYLOID OLIGOMERIZATION**

Chiara Lambruschini, Denise Galante, Lisa Moni, Francesco Ferraro, Giulio Gancia, Renata Riva, Alessia Traverso, Luca Banfi\* and Cristina D'Arrigo\*


***Table of contents***


Additional biophysical assays on polyhenols .....	S2
Copies of <sup>1</sup> H and <sup>13</sup> C NMR spectra of new compounds.....	S7
Copies of HPLC chromatograms of polyphenols <b>1a-q</b> .....	S66


## Additional biophysical assays on polyphenols


### Solubility tests

The first biophysical analysis performed on the new polyphenols was the solubility in aqueous solution because the working condition is Phosphate Buffer Solution (PBS) at pH 7.4 to mimic the physiological environment. Almost all compounds showed low solubility in a range from 25 to 500  $\mu\text{M}$  in aqueous solution except **1g**, **1h**, **1j** and **1q**. To overcome this obstacle, we dissolved all samples in 100% dimethyl sulfoxide (DMSO). In this solvent new polyphenols result all fully soluble. To continue working in aqueous solution, DMSO samples were diluted in PBS at the desired concentration by keeping the 1% DMSO in the final solution. Together with the solubility analysis, the turbidity trend depending on the concentration of the samples was also investigated, in order to verify that the new polyphenols did not aggregate or form micelles as the concentration increases, precipitating in water solution. In the following Figures, the blue solid line represents the absorbance of the samples at their characteristic wavelength over the concentration increase. The dashed red line shows the turbidity of the polyphenols at 405 nm over the concentration increase. For compounds **1a** and **1b** only the absorbance experiments were carried out.













## Aggregation kinetic tests on some polyphenols





Copies of  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of new compounds

























































































































































































## Copies of HPLC chromatograms of polyphenols 1a-q

Data File C:\HPCHEM\1\DATA\LISSEQ\AT14\_000.D

Sample Name: AT14

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
AT14 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml/m  
in, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300, 33  
0nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

=====  
Injection Date : 5/16/2017 6:24:27 PM           Seq. Line : 13  
Sample Name : AT14                                Location : Vial 11  
Acq. Operator : AeVeO                            Inj : 1  
Acq. Instrument : stanza306new                 Inj Volume : 5 µl  
Acq. Method : C:\HPCHEM\1\METHODS\POLIFEN.M  
Last changed : 5/16/2017 5:48:58 PM by AeVeO  
   (modified after loading)  
Analysis Method : C:\HPCHEM\1\METHODS\POLIFEN.M  
Last changed : 5/17/2017 9:14:31 AM by AeVeO  
   (modified after loading)  
=====



## Area Percent Report

Sorted By : Signal  
Multiplier : 1.0000  
Dilution : 1.0000  
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=220,4 Ref=450,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	5.441	BB	0.1305	2993.46826	347.44159	99.6945
2	12.148	PB	0.0737	9.17285	1.88715	0.3055

Totals :                                       3002.64111   349.32874

stanza306new 5/17/2017 9:35:44 AM AeVeO


Page 1 of 2

Data File C:\HPCHEM\1\DATA\LISSEQ\AT80\_000.D

Sample Name: AT80

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
AT80 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml/m  
in, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300, 33  
0nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

=====  
Injection Date : 5/16/2017 6:52:28 PM           Seq. Line : 14  
Sample Name : AT80                                Location : Vial 12  
Acq. Operator : AeVeO                            Inj : 1  
Acq. Instrument : stanza306new                 Inj Volume : 5 µl  
Acq. Method : C:\HPCHEM\1\METHODS\POLIFEN.M  
Last changed : 5/16/2017 5:48:58 PM by AeVeO  
   (modified after loading)  
Analysis Method : C:\HPCHEM\1\METHODS\POLIFEN.M  
Last changed : 5/17/2017 9:14:31 AM by AeVeO  
   (modified after loading)  
=====



## Area Percent Report

Sorted By : Signal  
Multiplier : 1.0000  
Dilution : 1.0000  
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 B, Sig=254,16 Ref=500,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	3.575	BB	0.1084	87.62709	11.97564	2.6084
2	5.730	PV	0.1116	41.26830	5.25543	1.2285
3	5.993	VB	0.1290	3230.46387	384.63614	96.1631

stanza306new 5/17/2017 9:43:45 AM AeVeO

Page 1 of 2




Data File C:\HPCHEM\1\DATA\LISA\FEF17\_02.D

Sample Name: FEF17 fr9-18

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
FEF17 fr9-18 (conc.:100ugr/ml MeOH), flusso 0,34ml/min,  
Vinj=5ul, Temp. 25°C Term.ON, Dad 326nm, grad. A=CH3CN  
- B=H2O, 0min B=90%, 20min B=0%

=====  
Injection Date : 5/10/2017 3:07:04 PM  
Sample Name : FEF17 fr9-18 Location : Vial 1  
Acq. Operator : AeVeO  
Acq. Instrument : stanza306new Inj Volume : 5 µl  
Acq. Method : C:\HPCHEM\1\METHODS\GRADACN.M  
Last changed : 5/10/2017 3:05:02 PM by AeVeO  
(modified after loading)  
Analysis Method : C:\HPCHEM\1\METHODS\GRADACN.M  
Last changed : 5/10/2017 3:57:22 PM by AeVeO  
(modified after loading)



=====  
Area Percent Report  
=====

Sorted By : Signal  
Multiplier : 1.0000  
Dilution : 1.0000  
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 B, Sig=326,16 Ref=550,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	15.757	PV	0.0572	15.45907	4.08185	0.6156
2	15.892	VB	0.0872	2495.76001	427.02893	99.3844

Totals : 2511.21908 431.11079

stanza306new 5/10/2017 3:57:38 PM AeVeO


Page 1 of 2

Data File C:\HPCHEM\1\DATA\LISSEQ\GB028\_00.D

Sample Name: GB-028

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
GB-028 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml  
/min, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300,  
330nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

=====  
Injection Date : 5/16/2017 2:40:19 PM Seq. Line : 5  
Sample Name : GB-028 Location : Vial 5  
Acq. Operator : AeVeO Inj : 1  
Acq. Instrument : stanza306new Inj Volume : 5 µl  
Acq. Method : C:\HPCHEM\1\METHODS\POLIFEN.M  
Last changed : 5/16/2017 12:31:23 PM by AeVeO  
Analysis Method : C:\HPCHEM\1\METHODS\POLIFEN.M  
Last changed : 5/17/2017 8:26:04 AM by AeVeO  
(modified after loading)



=====  
Area Percent Report  
=====

Sorted By : Signal  
Multiplier : 1.0000  
Dilution : 1.0000  
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 D, Sig=330,16 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.851	BB	0.1590	39.08525	2.94181	2.5704
2	7.852	BV	0.0899	81.44060	13.59060	5.3559
3	8.147	VB	0.0883	1400.04663	239.05367	92.0737

Totals : 1520.57248 255.58608




Data File C:\HPCHEM\1\DATA\LISSEQ\FEF07\_00.D

Sample Name: FEF07

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
FEF07 (conc.:100 ugr/ml MeOH), sequenza, flusso 0,34ml  
/min, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300,  
330nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

```
=====
Injection Date : 5/16/2017 5:28:27 PM      Seq. Line : 11
Sample Name    : FEF07                      Location  : Vial 10
Acq. Operator  : AeVeO                      Inj       : 1
Acq. Instrument: stanza306new               Inj Volume: 5 µl
Acq. Method    : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/16/2017 12:31:23 PM by AeVeO
Analysis Method: C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/17/2017 9:14:31 AM by AeVeO
              (modified after loading)
=====
```



=====  
Area Percent Report  
=====

```
Sorted By      :      Signal
Multiplier     :      1.0000
Dilution       :      1.0000
Use Multiplier & Dilution Factor with ISTDs
```

Signal 1: DAD1 D, Sig=330,16 Ref=360,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	9.198	PB	0.0937	1152.51221	180.00525	100.0000

Totals : 1152.51221 180.00525

Results obtained with enhanced integrator!

stanza306new 5/17/2017 9:25:26 AM AeVeO


Page 1 of 2

Data File C:\HPCHEM\1\DATA\LISSEQ\FEF51\_00.D

Sample Name: FEF51

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
FEF51 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml/  
min, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300, 3  
30nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

```
=====
Injection Date : 5/16/2017 4:32:17 PM      Seq. Line : 9
Sample Name    : FEF51                      Location  : Vial 8
Acq. Operator  : AeVeO                      Inj       : 1
Acq. Instrument: stanza306new               Inj Volume: 5 µl
Acq. Method    : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/16/2017 12:31:23 PM by AeVeO
Analysis Method: C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/17/2017 9:14:31 AM by AeVeO
              (modified after loading)
=====
```



=====  
Area Percent Report  
=====

```
Sorted By      :      Signal
Multiplier     :      1.0000
Dilution       :      1.0000
Use Multiplier & Dilution Factor with ISTDs
```

Signal 1: DAD1 D, Sig=330,16 Ref=360,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	7.439	MF	0.1247	836.88159	111.86374	96.8411
2	7.864	FM	0.2011	27.29815	2.26276	3.1589

Totals : 864.17974 114.12650

Data File C:\HPCHEM\1\DATA\LISSEQ\FEF57\_00.D


Sample Name: FEF57

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
FEF57 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml/m  
in, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300, 33  
0nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

```

=====
Injection Date : 5/16/2017 2:12:17 PM      Seq. Line : 4
Sample Name   : FEF57                      Location  : Vial 4
Acq. Operator : AeVeO                      Inj      : 1
Acq. Instrument : stanza306new              Inj Volume : 5 µl
Acq. Method   : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed  : 5/16/2017 12:31:23 PM by AeVeO
Analysis Method : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed  : 5/17/2017 9:01:18 AM by AeVeO
              (modified after loading)
=====

```



=====  
Area Percent Report  
=====

```

Sorted By      : Signal
Multiplier     : 1.0000
Dilution      : 1.0000
Use Multiplier & Dilution Factor with ISTDs

```

Signal 1: DAD1 B, Sig=254,16 Ref=500,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	5.776	BB	0.1403	2679.42847	291.30542	99.3218
2	9.893	PB	0.0748	18.29558	3.69103	0.6782

Totals : 2697.72405 294.99645

Data File C:\HPCHEM\1\DATA\LISSEQ\FEF21\_02.D


Sample Name: FEF21

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
FEF21 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml/m  
in, Vinj=5ul, Temp. 25°C Term.ON, Dad 232nm, grad. A=CH  
3CN - B=H2O, 0min B=90%, 15min B=0%

```

=====
Injection Date : 5/17/2017 10:18:55 AM      Seq. Line : 4
Sample Name   : FEF21                      Location  : Vial 2
Acq. Operator : AeVeO                      Inj      : 1
Acq. Instrument : stanza306new              Inj Volume : 5 µl
Acq. Method   : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed  : 5/17/2017 9:46:06 AM by AeVeO
              (modified after loading)
Analysis Method : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed  : 5/17/2017 11:27:06 AM by AeVeO
              (modified after loading)
=====

```



=====  
Area Percent Report  
=====

```

Sorted By      : Signal
Multiplier     : 1.0000
Dilution      : 1.0000
Use Multiplier & Dilution Factor with ISTDs

```

Signal 1: DAD1 C, Sig=254,16 Ref=500,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	11.033	VB	0.0774	970.13867	190.45187	97.9454
2	13.679	MM	0.1411	20.35086	2.40455	2.0546

Totals : 990.48953 192.85643

Data File C:\HPCHEM\1\DATA\LISSEQ\FEF20\_00.D


Sample Name: FEF20

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
FEF20 triturato (conc.:100ugr/ml MeOH), sequenza, flusso  
o 0,34ml/min, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 25  
4, 300, 330nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min  
B=0%

```

=====
Injection Date   : 5/16/2017 7:20:29 PM      Seq. Line : 15
Sample Name     : FEF20                      Location  : Vial 13
Acq. Operator   : AeVeO                      Inj       : 1
Acq. Instrument : stanza306new                Inj Volume: 5 µl
Acq. Method     : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed    : 5/16/2017 5:48:58 PM by AeVeO
                  (modified after loading)
Analysis Method : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed    : 5/17/2017 9:46:06 AM by AeVeO
                  (modified after loading)
=====

```



=====  
Area Percent Report  
=====

```

Sorted By      : Signal
Multiplier     : 1.0000
Dilution       : 1.0000
Use Multiplier & Dilution Factor with ISTDs

```

Signal 1: DAD1 D, Sig=330,16 Ref=500,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.730	VB	0.1041	2904.12012	422.79178	98.5112
2	9.256	VB	0.0918	28.66213	4.53398	0.9723
3	9.545	BB	0.0826	15.22732	2.71001	0.5165

Data File C:\HPCHEM\1\DATA\LISSEQ\FEF59\_00.D


Sample Name: FEF59

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
FEF59 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml/  
min, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300, 3  
30nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

```

=====
Injection Date   : 5/16/2017 8:16:25 PM      Seq. Line : 17
Sample Name     : FEF59                      Location  : Vial 15
Acq. Operator   : AeVeO                      Inj       : 1
Acq. Instrument : stanza306new                Inj Volume: 5 µl
Acq. Method     : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed    : 5/16/2017 5:48:58 PM by AeVeO
                  (modified after loading)
Analysis Method : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed    : 5/17/2017 9:46:06 AM by AeVeO
                  (modified after loading)
=====

```



=====  
Area Percent Report  
=====

```

Sorted By      : Signal
Multiplier     : 1.0000
Dilution       : 1.0000
Use Multiplier & Dilution Factor with ISTDs

```

Signal 1: DAD1 B, Sig=254,16 Ref=500,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.743	PB	0.1396	3149.39502	344.53494	100.0000
Totals :				3149.39502	344.53494	

Data File C:\HPCHEM\1\DATA\LISSEQ\RL19\_000.D


Sample Name: RL19

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
RL19 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml/m  
in, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300, 33  
0nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

```

=====
Injection Date : 5/16/2017 7:48:27 PM      Seq. Line : 16
Sample Name   : RL19                      Location : Vial 14
Acq. Operator : AeVeO                    Inj      : 1
Acq. Instrument : stanza306new           Inj Volume : 5 µl
Acq. Method   : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed  : 5/16/2017 5:48:58 PM by AeVeO
                (modified after loading)
Analysis Method : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed  : 5/17/2017 9:46:06 AM by AeVeO
                (modified after loading)
=====

```



=====  
Area Percent Report  
=====

```

Sorted By      : Signal
Multiplier     : 1.0000
Dilution       : 1.0000
Use Multiplier & Dilution Factor with ISTDs

```

Signal 1: DAD1 B, Sig=254,16 Ref=500,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.651	BB	0.1050	1787.80994	257.55042	100.0000

Totals :                    1787.80994   257.55042

Data File C:\HPCHEM\1\DATA\LISSEQ\FEF97\_00.D


Sample Name: FEF97

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
FEF97 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml/  
min, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300, 3  
30nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

```

=====
Injection Date : 5/16/2017 5:00:18 PM      Seq. Line : 10
Sample Name   : FEF97                      Location : Vial 9
Acq. Operator : AeVeO                    Inj      : 1
Acq. Instrument : stanza306new           Inj Volume : 5 µl
Acq. Method   : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed  : 5/16/2017 12:31:23 PM by AeVeO
Analysis Method : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed  : 5/17/2017 9:14:31 AM by AeVeO
                (modified after loading)
=====

```



=====  
Area Percent Report  
=====

```

Sorted By      : Signal
Multiplier     : 1.0000
Dilution       : 1.0000
Use Multiplier & Dilution Factor with ISTDs

```

Signal 1: DAD1 D, Sig=330,16 Ref=360,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	3.818	BB	0.1166	15.16598	1.57098	1.2582
2	5.251	FV	0.1110	13.12027	1.40461	1.0885
3	5.594	VB	0.1427	1112.28748	118.24020	92.2797
4	6.613	BB	0.1066	64.77035	9.26406	5.3736



Data File C:\HPCHEM\1\DATA\LISSEQ\FEF77\_00.D


Sample Name: FEF77

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
FEF77 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml/  
min, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300, 3  
30nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

```

=====
Injection Date : 5/16/2017 12:48:16 PM      Seq. Line : 1
Sample Name    : FEF77                      Location  : Vial 1
Acq. Operator  : AeVeO                      Inj      : 1
Acq. Instrument: stanza306new               Inj Volume: 5 µl
Acq. Method    : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/16/2017 12:31:23 PM by AeVeO
Analysis Method: C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/17/2017 9:01:18 AM by AeVeO
              (modified after loading)
=====

```



=====  
Area Percent Report  
=====

```

Sorted By      :      Signal
Multiplier     :      1.0000
Dilution       :      1.0000
Use Multiplier & Dilution Factor with ISTDs

```

Signal 1: DAD1 D, Sig=330,16 Ref=360,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.473	BB	0.1222	1101.45911	136.36723	91.8820
2	7.714	BB	0.0909	90.25157	14.85730	7.5286
3	8.071	BV	0.0836	7.06468	1.21925	0.5893

Totals :                    1198.77535   152.44378

Data File C:\HPCHEM\1\DATA\LISSEQ\GB017\_00.D


Sample Name: GB-017

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
GB-017 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml/  
min, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300,  
330nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

```

=====
Injection Date : 5/16/2017 8:44:25 PM      Seq. Line : 18
Sample Name    : GB-017                    Location  : Vial 16
Acq. Operator  : AeVeO                      Inj      : 1
Acq. Instrument: stanza306new               Inj Volume: 5 µl
Acq. Method    : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/16/2017 5:48:58 PM by AeVeO
              (modified after loading)
Analysis Method: C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/17/2017 9:46:06 AM by AeVeO
              (modified after loading)
=====

```



=====  
Area Percent Report  
=====

```

Sorted By      :      Signal
Multiplier     :      1.0000
Dilution       :      1.0000
Use Multiplier & Dilution Factor with ISTDs

```

Signal 1: DAD1 D, Sig=330,16 Ref=500,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	8.098	PV	0.0883	17.60988	2.96553	0.7288
2	8.298	VB	0.0874	2398.80054	408.96167	99.2712

Totals :                    2416.41042   411.92720

Data File C:\HPCHEM\1\DATA\LISSEQ\GB025\_00.D


Sample Name: GB-025

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
GB-025 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml  
/min, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300,  
330nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

```

=====
Injection Date : 5/16/2017 4:04:15 PM      Seq. Line : 8
Sample Name    : GB-025                    Location  : Vial 7
Acq. Operator  : AeVeO                     Inj       : 1
Acq. Instrument: stanza306new              Inj Volume: 5 µl
Acq. Method    : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/16/2017 12:31:23 PM by AeVeO
Analysis Method: C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/17/2017 9:14:31 AM by AeVeO
(modified after loading)
=====

```



=====  
Area Percent Report  
=====

```

Sorted By      : Signal
Multiplier     : 1.0000
Dilution       : 1.0000
Use Multiplier & Dilution Factor with ISTDs

```

Signal 1: DAD1 D, Sig=330,16 Ref=360,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	5.968	BB	0.1242	1476.92041	179.12212	95.7654
2	8.067	VB	0.0910	53.76893	8.96376	3.4864
3	9.152	VV	0.0785	11.53749	2.26333	0.7481

Totals : 1542.22683 190.34921

Data File C:\HPCHEM\1\DATA\LISSEQ\GB018\_00.D


Sample Name: GB-018

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
GB-018 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml  
/min, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300,  
330nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

```

=====
Injection Date : 5/16/2017 3:36:15 PM      Seq. Line : 7
Sample Name    : GB-018                    Location  : Vial 6
Acq. Operator  : AeVeO                     Inj       : 1
Acq. Instrument: stanza306new              Inj Volume: 5 µl
Acq. Method    : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/16/2017 12:31:23 PM by AeVeO
Analysis Method: C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/17/2017 9:14:31 AM by AeVeO
(modified after loading)
=====

```



=====  
Area Percent Report  
=====

```

Sorted By      : Signal
Multiplier     : 1.0000
Dilution       : 1.0000
Use Multiplier & Dilution Factor with ISTDs

```

Signal 1: DAD1 D, Sig=330,16 Ref=360,100  
Signal has been modified after loading from rawdata file!

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	7.076	PB	0.0922	1076.16138	176.38182	97.5291
2	8.063	PB	0.0850	14.37284	2.53977	1.3026
3	8.574	VB	0.0861	12.89170	2.21003	1.1683

Totals : 1103.42592 181.13162

Data File C:\HPCHEM\1\DATA\LISSEQ\FEF93\_00.D


Sample Name: FEF93

Colonna Phenyl C6 150x3mm 3um con precolonna, Campione:  
FEF93 (conc.:100ugr/ml MeOH), sequenza, flusso 0,34ml/m  
in, Vinj=5ul, Temp. 25°C Term.ON, Dad 220, 254, 300, 33  
0nm, grad. A=CH3CN - B=H2O, 0min B=60%, 10min B=0%

```

=====
Injection Date : 5/16/2017 1:44:17 PM      Seq. Line : 3
Sample Name    : FEF93                      Location  : Vial 3
Acq. Operator  : AeVeO                      Inj       : 1
Acq. Instrument: stanza306new                Inj Volume: 5 µl
Acq. Method    : C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/16/2017 12:31:23 PM by AeVeO
Analysis Method: C:\HPCHEM\1\METHODS\POLIFEN.M
Last changed   : 5/17/2017 8:26:04 AM by AeVeO
                (modified after loading)
=====

```



```

=====
Area Percent Report
=====

```

```

Sorted By      :      Signal
Multiplier     :      1.0000
Dilution       :      1.0000
Use Multiplier & Dilution Factor with ISTDs

```

Signal 1: DAD1 D, Sig=330,16 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.214	BB	0.1171	1499.56262	192.03836	99.3795
2	6.610	PB	0.1039	9.36271	1.36681	0.6205

Totals : 1508.92533 193.40517

Results obtained with enhanced integrator!