



Contents lists available at ScienceDirect

Cytokine

journal homepage: [www.journals.elsevier.com/cytokine](http://www.journals.elsevier.com/cytokine)

## Cigarette smoke affects IL-17A, IL-17F and IL-17 receptor expression in the lung tissue: Ex vivo and in vitro studies

Angela Marina Montalbano<sup>a</sup>, Loredana Riccobono<sup>a</sup>, Liboria Siena<sup>a</sup>, Giuseppina Chiappara<sup>a</sup>, Caterina Di Sano<sup>a</sup>, Giulia Anzalone<sup>a</sup>, Rosalia Gagliardo<sup>a</sup>, Fabio L.M. Ricciardolo<sup>b</sup>, Valentina Sorbello<sup>b</sup>, Loredana Pipitone<sup>c</sup>, Patrizio Vitulo<sup>c</sup>, Mirella Profita<sup>a,\*</sup>

<sup>a</sup> Unit: "Ex vivo/In vitro Models to Study the Immunopathology and Pharmacology of Airway Diseases", Institute of Biomedicine and Molecular Immunology (IBIM), Italian National Research Council (CNR), Palermo, Italy

<sup>b</sup> Department of Clinical and Biological Sciences, University of Torino, Torino, Italy

<sup>c</sup> Istituto Mediterraneo per i Trapianti e Terapie ad Alta Specializzazione (ISMETT), Palermo, Italy

### ARTICLE INFO

#### Article history:

Received 27 April 2015

Received in revised form 16 June 2015

Accepted 13 July 2015

Available online xxxxx

#### Keywords:

Th17 immunity

Epithelial cells

Airway inflammation

COPD

### ABSTRACT

Cigarette smoke is a risk factor for Chronic Obstructive Pulmonary Disease (COPD). Th-17 cytokines are involved in the pathogenesis of COPD. We aimed to evaluate the role of cigarette smoke on the expression of IL-17A, IL-17F and IL-17R in airways of COPD patients. Epithelial and subepithelial immunoreactivity for IL-17A, IL-17F and IL-17R was assessed in surgical specimens from COPD patients ( $n = 15$ ) and from healthy subjects (HC) ( $n = 10$ ) by immunohistochemistry. In vitro, human epithelial cell line 16HBE and A549 as well as PBMC from normal donors were stimulated with cigarette smoke extract (CSE) (0%, 2.5%, 5%, 10%) to evaluate the IL-17A, IL-17F and IL-17R expression by flow cytometry. Furthermore, rhIL-17A and CSE stimulation was evaluated on proliferation and apoptosis in 16HBE and in A549. In central and distal airways immunoreactivity for IL-17A, IL-17F and IL-17R significantly increased in the epithelium and IL-17A in the subepithelium from COPD than in HC. In distal airway, immunoreactivity for IL-17F increased in the subepithelium of COPD than in HC. IL-17A immunoreactivity positively correlate with IL-17R and total pack years in the epithelium from central and distal airways of COPD patients. In vitro, CSE stimulation significantly increased IL-17F and IL-17R in 16HBE (2.5%) and A549 (5%) while IL-17A and IL-17F in PBMC (10%). IL-17A and CSE stimulation, rather than CSE or rhIL-17A alone, significantly increased proliferation in 16HBE and apoptosis in A549. Cigarette smoke increases Th17 immunity in lung tissue of COPD patients, promoting the mechanism of proliferation and apoptosis in airway epithelial cells.

© 2015 Elsevier Ltd. All rights reserved.

### 1. Introduction

Chronic obstructive pulmonary disease (COPD) is a complex disease characterized by chronic innate and adaptive inflammatory immune responses. Cigarette smoke is the major risk factor for the development of COPD, cigarette smokers constituting more than 90% of all COPD patients in developed countries [1]. The pathological hallmarks of COPD are destruction of the lung parenchyma, which characterizes pulmonary emphysema, inflammation of the peripheral airways, respiratory bronchiolitis and inflammation of the central airways [1–3]. However, there are evidences of increased inflammation in both central and peripheral airways,

involving a range of cell types, including neutrophils, macrophages, lymphocytes and epithelial cells. To what extent central airways may mirror events, occurring in distal lung is uncertain.

Adaptive immune processes are implicated in the pathogenesis of COPD. It has been hypothesized that susceptibility to COPD may arise by a shift from the non-specific innate response present in every smoker toward an adaptive immune response with features typical of autoimmune processes [4,5]. Th17-cells are effector T-cell subsets characterized by the production of IL-17A, IL-17F and IL-22, implicated in the pathogenesis of several inflammatory and autoimmune diseases such as multiple sclerosis, rheumatoid arthritis and psoriasis [6,7]. IL-17A and IL-17F targeted the cells on IL-17 receptor (IL-17R). The IL-17R is expressed in blood cells as well as in structural cells such as airway epithelial cells [8–11]. Th17 immunity and the related receptor are involved in both the innate and the adaptive aspects of airway immunity,

\* Corresponding author at: Institute of Biomedicine and Molecular Immunology "A. Monroy", Italian National Research Council, Palermo, Italy.

E-mail address: [profita@ibim.cnr.it](mailto:profita@ibim.cnr.it) (M. Profita).

which represent the crucial crosstalk between the immune system and structural cells [12]. It has been reported that IL-17A levels are increased in submucosal biopsy specimens from the large airways of patients with COPD compared with control subjects, but there is no difference for IL-17F [13] or in contrast that both IL-17A and IL-17F positive cells were higher in bronchial biopsies from COPD patients than in healthy control [14]. Furthermore, it was observed that the levels of IL-17A are increased in the submucosa of lung tissue from COPD than in S and HC while there were no differences in the subepithelium [15]. However, no findings clarify the role of IL-17 immunity in central or distal airway of COPD suggesting the necessity to compare the levels IL-17A, IL-17F and the related receptor (IL-17R) in the two lung districts.

The aim of the present study is to evaluate the role of Cigarette Smoke on IL-17A, IL-17F and IL-17R using *ex vivo/in vitro* studies. *Ex vivo*, we investigated the immunoreactivity of IL-17A, IL-17F and IL-17R in the epithelium and in the subepithelium of the central and distal airways from surgical specimens of smoking COPD patients. *In vitro*, we evaluated the effect of cigarette smoke extract (CSE) on the expression of IL-17A, IL-17F and IL-17R in human bronchial epithelial cell line (16HBE), in alveolar cell line (A549) and in human peripheral blood mononuclear cells (PBMC) from normal donors. Finally, we studied whether CSE, via the increased expression of IL-17R, strengthen the IL-17A mediated activities on the proliferation and on the apoptosis of epithelial cell line from central and distal airways.

## 2. Materials and methods

### 2.1. Patient population

Patients underwent surgery for lung cancer and were recruited at ISMETT-Palermo, Italy. The study was approved by the ISMETT Ethic Committee (#217806-30/06/2008) and was in agreement with Helsinki Declaration. Written informed consent was obtained from each patient. The study was carried out using samples of three groups of subjects: 10 healthy asymptomatic non-smoking subjects with normal lung function (HC); 15 patients with chronic obstructive pulmonary disease (COPD). The diagnosis of COPD and the assessment of its severity were defined and classified according to the criteria reported by the Global Initiative for Obstructive Lung Disease (GOLD) guidelines for COPD management (GOLD stage  $\geq$  I) [16]. COPD subjects with exacerbations within 1-month prior to the study were excluded. Patients with COPD had a smoking history of 10  $\geq$  pack years or more.

COPD patients were treated with bronchodilators and were classified based on preoperative lung function: FEV1 less than 80% of reference, FEV1/FVC less than 70%, and bronchodilatation effect less than 12%. The patients were not under corticosteroid therapy (neither inhaled nor systemic) and not under antibiotics and did not have exacerbations during the month preceding the study. Subjects had negative skin tests for common allergen extracts and had no past history of asthma or allergic rhinitis.

### 2.2. Immunohistochemistry of paraffine sections

Tissue specimens from tumor-free central bronchi and peripheral lung tissue were selected, fixed with 10% neutral buffer formalin and embedded in paraffin wax. Sequential sections (3  $\mu$ m thick) were placed on poly-L-lysine coated slides, deparaffinized in xylene, rehydrated in a descending ethanol series and stained with haematoxylin and eosin (HE). Immunoreactivity for Rabbit Polyclonal IL-17A Antibody (H-132): sc7927 (Santa Cruz Biotechnology, Santa Cruz, CA), Human IL-17F Polyclonal Goat IgG (R&D System) and Monoclonal Anti-Human IL-17R Antibody

(R&D System) was evaluated in Central (internal perimeter > 6 mm) and Distal (internal perimeter,  $\leq$  6 mm) airways [17]. LSAB2 Dako kit (Code Nu K0674) (Dako, Glostrup, Denmark) and Fuchsin Substrate-Chromogen System Dako [18] were used for the staining for IL-17A and IL-17R Antibodies, meanwhile LSAB Dako kit Universal (Code Nu K0689) (Dako) and Fuchsin Substrate-Chromogen System Dako were used for staining of IL-17F antibody. Rabbit, mouse and goat negative control immunoglobulins (Dako) were used for negative controls. Two independent investigators, using image analysis (Leica microscope, Wetzlar, Germany) 400 $\times$  magnification, evaluated sample immunoreactivity blindly.

The length of the basement membrane was evaluated using a Leica Application Suite V3.3 (LAS) software (Leica) for Image Analysis. Results were expressed as the number of positive epithelial cells/mm basement membrane as previously described [19]. Finally, immunostained cells were quantified in the subepithelium of central and distal airway, and the results were expressed as the number of positive cell/mm<sup>2</sup> [15,13].

### 2.3. Epithelial cell cultures

The SV40 large T antigen-transformed 16HBE cell line (16HBE), from normal bronchial epithelial cells [20] was used for this study. The 16HBE cell line retains the differentiated morphology and function of normal airway epithelial cells. The cells represent a clonal diploid (2n<sup>1/4</sup>/6) cell line isolated from human lungs previously used to study the functional properties of bronchial epithelial cells in inflammation and repair processes [21]. 16HBE cells were cultured as adherent monolayers in Eagle's minimum essential medium (MEM) supplemented with 10% heat-inactivated (56 °C, 30 min) foetal bovine serum (FBS), 1% MEM (non-essential aminoacids, Euroclone), 2 mM L-glutamine and gentamicin 250  $\mu$ g/ml at 37 °C in a humidified 5% CO<sub>2</sub> atmosphere [22]. Type II alveolar epithelial cell-derived A549, cell line were purchased from American Type Culture Collection (ATCC; Rockville, MD). Cells were cultured in complete culture medium (RPMI 1640 containing 10% FCS and 200 IU/mL penicillin/100  $\mu$ g/mL streptomycin).

### 2.4. Preparation of cigarette smoke extracts (CSE)

Commercial cigarettes (Marlboro; Philip Morris USA, Richmond, VA) were used in this study. CSE was prepared as described previously described [22]. Cigarette smoke extract was used to stimulate cultured 16-HBE and A549 cell line and Peripheral Blood Mononuclear Cells (PBMC) obtained from healthy control.

### 2.5. Isolation of human peripheral blood mononuclear cells (PBMC)

Blood samples were collected in EDTA vacutainer tubes (BD Biosciences) and used for plasma selection and PBMC isolation from Healthy donors. The cells were isolated using a density gradient centrifugation (Ficoll-paque™ PLUS; Amersham Biosciences SE-751 84, Uppsala, Sweden). After two washes, the cells were suspended in RPMI 1640 cell culture medium (Invitrogen Life Technologies, Italy) supplemented with 10% heat-inactivated fetal bovine serum (FBS), 2 mM L-glutamine, 20 mM HEPES, 100 U/ml penicillin, 100  $\mu$ g/ml streptomycin, 5  $\times$  10<sup>-5</sup> M 2-ME and 85  $\mu$ g/ml gentamicin. Purity of PBMC was assessed by May-Grünwald-Giemsa staining and was 93  $\pm$  3% accordingly with the purity declared in the data sheet of the Ficoll-Paque™ PLUS density gradient. The viability of the PBMC was tested using trypan blue exclusion and was 93  $\pm$  5% as previously described [23].

**Table 1**  
Demographic characteristics of patients.

	Control	COPD	Overall <i>p</i> value
Subject number	<i>n</i> = 10	<i>n</i> = 15	
Sex, male/female	8/2	12/3	
Age, yr	69.5 ± 5.8	76.3 ± 13.2	N.S.
FEV1, % predicted	106 ± 19.4	68.8 ± 17.5	<i>p</i> < 0.001
FEV1/FVC, %	80.3 ± 2.9	60.9 ± 3.5	<i>p</i> < 0.001
Smoking, pack/yr	0	50.9 ± 36.2	–

Data are shown as mean ± S.D.

Abbreviations: FEV1 = forced expiratory volume in 1 s.

FVC = forced vital capacity.

Statistical analysis for multiple comparisons was performed by Mann–Whitney U-test.

## 2.6. PBMC stimulation


PBMC ( $1 \times 10^6$  cells/ml) were cultured in the presence of CSE (2.5%, 5%, 10%) for 48 h in 24-well cell culture plates in complete medium RPMI 1640 (Invitrogen Life Technologies, Italy) plus 10% heat-inactivated FBS. Cultured PBMC were assessed for intracellular cytokines IL-17A, IL-17F, IL-17R expression, by immunocytochemistry, immunofluorescence and flowcytometric analysis. After the stimulation, the viability of the cells was tested using trypan blue exclusion.

## 2.7. 16HBE and A549 cells stimulation


16HBE and A549 were plated (200,000 cells/well) in standard six-well plates, in a suitable medium at 10% FBS and grown to confluence (70–80%). After, the mediums were replaced and the 16HBE and A549 cells were stimulated in the presence or absence of different concentration of CSE (2.5%, 5%, 10%) for 24 h to evaluate IL-17A, IL-17F, IL-17R expression by immunocytochemistry, immunofluorescence, and flowcytometric analysis. Furthermore, 16HBE and A549 were stimulated in the presence or absence of CSE (2.5% and 5% respectively) with or without hrIL-17A (20 ng/ml) to study cell apoptosis and cell proliferation after 24 h.

## 2.8. Flowcytometric analysis for IL-17A, IL-17F, and IL-17 Receptor

Monensin 1 mM was added to 16HBE, A549 cells and PBMC (800,000 cells/well) stimulated with CSE for 12 h before the collection. The cells were washed in PBS and collected in FACS tubes. After wards, cells were washed in staining buffer (PBS containing 1% FCS and 0.1% Na azide) and then incubated with PBS containing 4% paraformaldehyde for 20 min. Fixation was followed by two washes in permeabilization buffer (PBS containing 1% FCS, 0.3% saponin, and 0.1% Na azide) and fixed permeabilized cells were stained with the primary antibody, 1 h, 4 °C. The 16HBE, A549 cells and PBMC were analyzed for rabbit-polyclonal antibody anti IL-17



**Fig. 1.** Immunoreactivity for IL-17A, IL-17F and IL-17R in surgical specimens from central airways of HC (*n* = 10) and COPD (*n* = 15) subjects. Cells were stained with an anti-IL-17A, anti-IL-17F and anti-IL-17R antibodies. Negative control was performed using rabbit immunoglobulins negative control (see Section 2 for details). (A) Counts for the number of positive epithelial cells/mm basement membrane (left column); (B) Counts for the number positive subepithelial cells/mm<sup>2</sup> (right column) in central airways (right column). Results were expressed as box plot representing median and 25–75 percentiles. Statistical analysis was performed by Kruskal Wallis test followed by Fisher's PLSD correction for multiple comparison. Significance was accepted at *p* < 0.05.



**Fig. 2.** Immunoreactivity for IL17A, IL17F and IL-17R in surgical specimens from distal airways of HC subjects ( $n = 10$ ) and COPD patients ( $n = 15$ ). Cells were stained with an anti-IL-17A, anti IL-17F and anti IL-17R antibodies. Negative control were performed using rabbit immunoglobulins (see Section 2 for details). (A) Counts of the number of positive epithelial cells/mm basement membrane (left column); (B) Counts of the number positive subepithelial cells/mm<sup>2</sup> (right column) in central airways (right column). Results were expressed as box plot representing median and 25–75 percentiles. Statistical analysis was performed by Kruskal Wallis test followed by Fisher's PLSD correction for multiple comparison. Significance was accepted at  $p < 0.05$ .

(H-132), Santa Cruz Biotechnology, Santa Cruz, Calif.), anti-human IL-17F Antibody (R&D System), and Monoclonal anti-Human IL-17Receptor Antibody (R&D Systems, Inc. Minneapolis), expression by FACS analysis. Non-immune IgG at the same titre as the primary antibody was used as a negative control. Cells were washed in cold PBS and incubated with FITC-conjugated polyclonal swine anti-Rabbit Ig (DAKO Glostrup, Denmark), Polyclonal Rabbit anti mouse/FITC IgG (DAKO Glostrup, Denmark) and Polyclonal Rabbit Anti-goat IgG/FITC (DAKO Glostrup, Denmark) in the dark 1 h at 4 °C before flow cytometric analysis. After washing, the cells were analysed by FACS Calibur (Becton Dickinson, Mountain View, CA, USA) flow cytometer. Fluorescence-positive cells were quantified. Percentages of positive cells for IL-17A, IL-17F and IL-17R of positive cells were determined from forward (FS) and sideways (SS) scatter patterns, after gating on the cells, excluding debris. Non-specific binding and background fluorescence were quantified by analyzing negative control.


### 2.9. Immunocytochemistry

Cytospins were made from cultured PBMC stimulated with and without CSE 10% and fixed and permeabilized as previously described. After washing in PBS, slides were incubated with blocking solutions (0.5% BSA in PBS) 1 h RT, and then slides were incubated with the primary antibodies IL17 (1:50 dilution), IL17F

(1:10 dilution), and IL17R (1:10 dilution) overnight 4 °C. The reaction was revealed by LSAB and LSAB2 KIT phosphatase method (DAKO Glostrup Denmark) in 16-HBE, A549 and PBMC according to the manufacturer's instructions. Non-immune IgG at the same titre as the primary antibody were used as a negative controls. Cytospins were examined under light microscopy with a final magnification of 400×.

### 2.10. Immunofluorescence Co-Localization of IL-17A/IL-17R and IL-17F/IL-17R on airway epithelium

Double immunofluorescence tests were performed to assess the co-localization of rabbit polyclonal antibody IL-17 (H-132), Santa Cruz Biotechnology or the goat polyclonal antibody IL17 F (R&D System), with mouse monoclonal IL-17 R Antibody (R&D System). Frozen sections of Bronchial ring (6 μm) and Parenchymal sections (9 μm) were subjected to indirect immunofluorescence as follow described. Briefly, the samples were fixed in 4% paraformaldehyde for 15 min at room temperature, followed by 3 washes in PBS, 3 min each. The samples were then permeabilized with Saponin 0.05% in PBS-plus 3% BSA for 5 min at room temperature and after 3 washes in PBS, were fixed in cold Aceton at -20 °C for 7 min. After washes in PBS, samples were blocked in 3% BSA in PBS for 1hour at room temperature. Primary antibody incubation was performed in 3% BSA in PBS



**Fig. 3.** Representative immunostaining of IL17A, IL17F and IL17R in central and in distal airways. (A) Representative images for IL17A, IL-17F and IL-17R immunostaining (red stain) of epithelium and subepithelium in central (left panel) and distal (right panel) airways of HC, and COPD subjects (original magnification 400 $\times$ ). Arrow show immunoreactive cells in the epithelium (higher magnification in black box) and in the subepithelium (higher magnification in white box). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

overnight. The following primary antibodies were used: anti IL-17A (rabbit, 1:50), anti IL-17R (mouse, 1:50) and anti IL-17F (goat, 1:50) all overnight at 4 °C. The day after, slides were wash in PBS and the secondary antibodies incubation was in 3% BSA in PBS at room temperature for 1 h at room temperature, followed by three PBS washes. FITC-conjugated sheep anti-rabbit IgG (Sigma Aldrich, F7512), R-Phycoerythrin-conjugated goat anti-mouse IgG (Sigma P9287) and Alexa 488 conjugated rabbit anti-goat IgG secondary antibodies (Thermo Scientific Cat. N. SA5-10078) were used all 1:200 1 h in the dark at room temperature. The samples were then incubated with Hoechst (Sigma-Aldrich, Inc, Milan, Italy) 1:1000 in PBS for 10 min at room temperature, followed by PBS wash. The slides were mounted with Vectashield (Vector Laboratories, Burlingame, CA), and images were analyzed by using a laser scanning microscope ZEISS at a final magnification of 400 $\times$ . All immunofluorescence tests were performed with negative controls where no primary antibody was added.

#### 2.11. Immunofluorescence of 16HBE, A549 and PBMC

16HBE and A549 cells were seed in a 6 wells plates within we have insert a sterile cover slide. Cells growth until a 70–80% of confluence in presence or absence of CSE (2.5% and 5% respectively) for 24 h and the cover slide was recovered. Cultured PBMC were stimulated with CSE (10%) for 24 h and at the end of stimulation, cytopspins were performed to obtained the slides. The cells were fixed in Paraformaldehyde 4% 15 min. RT, washed in PBS and treated with Permeabilization Buffer (0.05% Saponin in PBS-plus 3% BSA). Then, the cells were incubated with blocking solution (BSA 0,5%in PBS) 1 h RT. Immunofluorescent staining was performed with the rabbit polyclonal antibody IL-17A (H-132), Santa Cruz Biotechnology, the goat-polyclonal antibody IL17F (R&D System), and IL-17Receptor Antibody (R&D System)


all at 1:25 dilution, overnight at 4 °C. The primary antibodies were diluted in PBS plus 3% BSA. Non immune IgG was used as a negative control. Secondary antibodies, anti-Rabbit IgG (whole molecule)-FITC F7512 Sigma Aldrich, Anti Mouse IgG (whole Molecule) R-Phycoeritrin conjugate P9287 Sigma and Alexa Fluor 488 Donkey anti-goat IgG (A11055) was performed in the dark for 1 h at room temperature. After wash in PBS, the slides were counterstained with DAPI and after wash in PBS and mounting with Mounting Medium Vectashield. Slides were cover slipped in Vectashield (Vector Laboratories, Burlingame, CA). Images were analyzed by using a laser-scanning microscope ZEISS at a final magnification of 400 $\times$ .

#### 2.12. Cell apoptosis by flow cytometry

The cells were stained with a solution containing a mixture of Annexin V-FITC in binding buffer 1X. After incubation (15 min in total darkness, RT) was add Propidium Iodide just before analysis. The number of viable, apoptotic and necrotic cells were determined using the FACS Calibur flow cytometer (Becton Dickinson, San Jose, CA). Results were presented as a percentage of counted cells.

#### 2.13. Cell proliferation assay

Cell proliferation was measured using carboxyfluorescein succinimidyl ester (CFSE) labeling assay. CFSE is used as fluorescently label live cells and is equally partitioned to daughter cells during division [24,25]. Briefly, the cells were incubated with CFSE (Molecular Probes, Inc. Eugene, OR) (at a final concentration of 5  $\mu$ M) at 37 °C for 10 min. Labeling was blocked by the addition of an equal volume of heat inactivated FCS. Tubes were placed in ice for 5 min and then washed. The cells were plated at  $8 \times 10^5$  - cells/well in six-well plates and incubated at 37 °C with 5% CO<sub>2</sub>. Cell proliferation was assessed by flow-cytometry.



**Fig. 4.** Spearman's rank correlations in central and distal airways of COPD ( $n = 15$ ) patients. (A) Correlation between the Total Pks year and immunoreactivity for IL-17A in the epithelium of central and distal airways; (B) Correlation between the immunoreactivity for IL-17A and immunoreactivity for IL-17R in the epithelium of central and distal airways. Data were reported as individual values. Statistical analysis was performed by Spearman's rank test. Significance was set at  $p < 0.05$ .

#### 2.14. Statistical analysis

Statistical comparisons in order to test differences between the three groups (HC, COPD) were made by use of the Kruskal–Wallis test followed by Fisher's PLSD correction for multiple comparisons. Data were expressed as median and inter quartile range. ANOVA test was used for the analysis of the data obtained from in vitro experimental conditions expressed as mean  $\pm$  S.D. Correlations were calculated according to Spearman test. All statistical analyses were performed using StatView<sup>®</sup> 5 software (SAS institute Inc.). A  $p$  value of less than 0.05 was considered to indicate statistical significance in these analyses.

### 3. Results

#### 3.1. Demographic characteristics of the subjects

The demographic characteristics and the functional evaluations of the studied groups are shown in Table 1. All recruited patient groups were similar with regard to age.

#### 3.2. IL-17A, IL17F and IL-17R expression in central airways and distal airways


We detected statistical higher levels of immunoreactivity for IL-17A, IL17F and IL17R expressed as positive cells/mm in the epithelium from the central airways of COPD patients compared

with HC subjects ( $p < 0.0001$ ,  $p = 0.0001$  and  $p = 0.0005$ , respectively) (Fig. 1A). The immunoreactivity for IL-17A and IL-17F expressed as positive cells/mm<sup>2</sup> showed statistical higher levels in the subepithelium from central airways of COPD subjects than in HC subjects ( $p < 0.0009$  and  $p = 0.0005$ , respectively). No significant differences were observed for the IL-17R immunoreactivity in the cells of the subepithelial region from central airways of COPD patients than in HC subjects (Fig. 1B).

Immunoreactivity for IL-17A ( $p = 0.0011$ ), IL-17F ( $p < 0.0005$ ) and IL-17R ( $p = 0.007$ ) were statistical significant higher in the epithelium from distal airways of COPD than in HC subjects (Fig. 2A). Additionally, immunoreactivity for IL-17A was statistical significant higher in the subepithelium from distal airways of COPD patients compared with HC subjects ( $p < 0.0001$ ) (Fig. 2B). No significant differences were observed for immunoreactivity of IL-17F or IL-17R in subepithelial region from central airways of COPD patients than in HC subjects. A representative images of the immunoreactivity for IL-17A, IL-17F and IL-17R in the epithelial and sub-epithelial regions from central and distal airways of HC and COPD are represented Fig. 3.

#### 3.3. Correlations

Immunoreactivity for IL-17A positively correlated with the total packs year in epithelial cells (cells/mm) from central and distal airways of COPD patients ( $p = 0.006$ ,  $Rho = -0.7$  and  $p = 0.04$ ,  $Rho = 0.6$  respectively) (Fig. 4A). Furthermore immunoreactivity for IL-17A positively correlated with epithelial cells



**Fig. 5.** Double immunofluorescence for IL-17A or IL-17F and IL-17R in epithelial cells from surgical specimens. Epithelial cells from HC subjects and COPD patients showed single staining for IL-17A (green) (A) or IL-17R (red) (B), or a double positivity (orange) (C) in central airways. Some of the latter are indicated by arrows. Epithelial cells from HC subjects and COPD patients showed single staining for IL-17A (green) (D) or IL-17R (red) (E), or double positivity (orange) (F) in distal airways. Some of the latter are indicated by arrows. Epithelial cells from HC subjects and COPD patients showed single staining for IL-17F (green) (A) or IL-17R (red) (B), or a double positivity (orange) (C) in central airways. Some of the latter are indicated by arrows. Epithelial cells from HC subjects and COPD patients showed single staining for IL-17A (green) (D) or IL-17R (red) (E), or double positivity (orange) (F) in distal airways. Some of the latter are indicated by arrows. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

immunoreactivity for IL-17R (cells/ml) in epithelial cells (cells/mm) from central and distal airways of COPD patients ( $p = 0.05$ ,  $Rho = -0.5$  and  $p = 0.02$ ,  $Rho = 0.6$  respectively) (Fig. 4B).

#### 3.4. Double immunofluorescence for IL-17A and IL-17R in epithelial cells from surgical specimens

Double immunofluorescence experiments, performed on tissue sections from surgical specimens, showed that epithelial cells from central and distal airways had higher levels of IL-17A and IL-17R co-localization in COPD patients than in HC subjects (Fig. 5). Furthermore, epithelial cells from central and distal airways of COPD had low levels of IL-17F and IL-17R co-localization in COPD patients. These results support the positive correlation obtained between the immunoreactivity of epithelium for IL-17A and IL-17R obtained with the detection of positive cells by immunohistochemistry.


#### 3.5. In vitro, cigarette smoke extract induces IL-17A, IL-17F and IL-17R expression in 16HBE, A549 and in PBMC

In vitro, we studied the effect of CSE on IL17A, IL17F and IL17R expression in epithelial cell line 16HBE and A549 as well as in


PBMC from normal donors. Using flow cytometry, immunocytochemistry and immunofluorescence analyses, we showed that CSE used at the percentage of 2.5% statistically significant increase the expression of IL-17F ( $p < 0.021$ ) and IL-17R ( $p < 0.024$ ) in 16HBE cell line compared with untreated cells (Fig. 6). Using the same technique, we showed that CSE stimulation at the percentage of 5% statistically significant increase the expression of IL-17F ( $p < 0.015$ ) and IL-17R ( $p < 0.002$ ) in A549 cell line (Fig. 7). Furthermore, we did not observed IL-17A expression in both 16HBE and A549 untreated or stimulated with CSE. Finally, we observed that CSE at the percentage of 10% statistical significant increase the IL-17A ( $p < 0.036$ ), IL-17F ( $p < 0.044$ ) while did not affect the expression of IL-17R (data not shown) in PBMC compared with untreated cells (Fig. 8).

#### 3.6. Effect of CSE and IL-17A on cell proliferation and apoptosis of 16HBE and A549

CSE 2.5% or rhIL-17A did not affect cell proliferation in stimulated 16HBE cell line compared to untreated cells. The combined use of CSE and rhIL-17A significant increase cell proliferation compared with 16HBE cell line stimulated with CSE, rhIL-17A or untreated cells ( $p < 0.014$ ;  $p < 0.017$  and  $p < 0.003$  respectively)



**Fig. 6.** Effect of CSE 2.5% on the expression of IL-17F and IL-17R in 16-HBE cells. Cells were incubated with and without CSE for 24 h and tested for IL-17F and IL-17R (cytokines or proteins) expression by (A) Flowcytometric analysis. Bars represent mean  $\pm$  SD of % of positive gated cells of 6 separate experiments. (B) Representative flow-cytometry analysis for IL17F and IL17R in 16HBE cell line; (C) Representative immunocytochemical and immunofluorescent analysis (original magnification 400 $\times$ ). Statistical analysis was performed by ANOVA test with Fisher's correction for multiple comparisons. Significance was set at  $p < 0.05$ .




**Fig. 7.** Effect of CSE 5% on the expression of IL-17F and IL-17R in A549 cell line. Cells were incubated with and without CSE for 24 h and tested for IL-17F, and IL-17R (cytokines or proteins) expression by (A) Flowcytometric analysis. Bars represent mean  $\pm$  SD of % of positive gated cells of 6 separate experiments. (B) Representative flow-cytometry analysis for IL17F and IL17R in A549 cell line; (C) Representative immunocytochemical and immunofluorescent analysis (original magnification 400 $\times$ ). Statistical analysis was performed by ANOVA test with Fisher's correction for multiple comparisons. Significance was set at  $p < 0.05$ .

(Fig. 9A and B). CSE 5% or rhIL-17A increased cell apoptosis in A549 cell line compared to untreated cells although without a statistical significant difference. Furthermore, we showed that the combined use of CSE 5% and rhIL-17A statistical significant increase cell apoptosis in A549 cell line than in untreated cells or than in CSE or in rhIL-17A treated cells ( $p < 0.003$ ,  $p < 0.012$  and  $p < 0.011$ ; respectively) (Fig. 9C and D).

#### 4. Discussion

This study compare for the first time the levels of IL-17A, IL-17F and IL-17R in the central and distal airways from smoking COPD patients. Our results showed that the immunoreactivity of IL-17A, IL-17F and IL-17R was increased in airway epithelial cells in central and distal airways from COPD patients. Particularly, we showed






**Fig. 8.** Effect of CSE 10% on the expression of IL-17A and IL-17F in PBMC from normal donors. Cells were incubated with and without CSE for 24 h and tested for IL17A and IL17F (cytokines or proteins) expression by (A) Flowcytometric analysis. Bars represent mean  $\pm$  SD of % of positive gated cells of 6 separate experiments. (B). Representative flow-cytometry analysis for IL17A and IL17F in PBMC. (C) Representative immunocytochemical and immunofluorescent analysis (original magnification 400 $\times$ ). Statistical analysis was performed by ANOVA test with Fisher's correction for multiple comparisons. Significance was set at  $p < 0.05$ .

that the levels of IL-17A immunoreactivity was higher, in submucosa than in epithelium of central and distal airways from COPD patients. Furthermore, we observed that the low levels of IL-17A immunoreactivity observed in epithelial cells of both districts positively correlated with IL-17R expression and with smoke habit in COPD patients. These findings might suggest that in the presence of cigarette smoke IL-17A, exclusively produced by infiltrating cells, target the epithelium in the IL-17R highly expressed in COPD. By in vitro experiments, we investigated a possible involvement of IL-17A in the processes of tissue renewal and damage of the epithelial cells in bronchial and distal airways from COPD patients. Accordingly, we found that IL-17A is able to promote airway epithelial cell activation generating cell proliferation in 16HBE and cell apoptosis in A549, representing epithelial cell line from bronchial and distal airways respectively. Further studies might be necessary to clarify the role of IL-17A in the induction of inflammatory gene expression during COPD in airway epithelial cells.

The airway epithelium is a dynamic tissue capable of self-renewal and proliferation after injury that undergoes slow but constant renewal [26]. Acting as a physical barrier, the lung epithelium regulates lung fluid balance, modulates metabolism and clearance of inhaled agents, and secretes numerous mediators, several of which recruit and activate inflammatory cells in response to injury [27]. Dysregulation of airway epithelial cell function related to environmental triggers, like cigarette smoke, may contribute to the pathogenesis of major lung diseases such as COPD. Th17 immunity orchestrated mechanisms inherent tissue damage in response to microbial infection and perpetuation of an autoimmune response in airway inflammation. Furthermore, Th17 immunity explain these phenomena underlying T cell-mediated damage to tissue. IL-17A was exclusively expressed in inflammatory cells present in the subepithelial cells even if some immunoreactive cells were present in the epithelium and within smooth muscle bundles in patients with asthma as well as IL-17F [28]. Our results showed that IL-17A, IL-17F and the related receptor IL-17R are increased in the epithelial cells as well as that IL-17A and IL-17F were increased in the sub epithelium of both central and distal airways from smoking COPD. These findings might suggest that cigarette smoke habit are able to increase the expression

of these cytokines and of the IL-17R in both district of COPD patients. The data of our study are in agreement with previous results showing that cigarette smoke exposure, upregulated IL-17A/F in human lung tissue explants from both non-COPD and COPD subjects [29]. It was observed that COPD and Healthy smokers have increased numbers of interleukin 17A+ cells in the bronchial submucosa where T-cells might represent an important source of this cytokine in the presence of smoking habit [13]. Furthermore, it was shown that the up-regulation of Th17 might be associated with cigarette smoke in lung tissue of mice [30]. These data together with our findings support the implications of Th17 immunity in the pathogenesis of COPD in association with the risk factor cigarette smoke. Particularly, we found higher levels of cells immunoreactive for IL-17A in the subepithelium than in epithelium of both central and distal airways from smoking COPD. Furthermore, higher levels of IL-17F were observed in the subepithelium of distal airways from smoking COPD than in the epithelium. The co-localization, observed by immunofluorescence, demonstrated the presence of IL-17A or IL-17F immunoreactivity in the epithelial cells expressing IL-17R in both central and distal airways from smoking COPD patients than in HC subjects. These findings might suggest the concept that both IL-17A and IL-17F are involved in the activation of epithelium in COPD patients having smoking habit. Moreover, the analysis of the correlations showed only a positive relationship between IL-17A expression and smoking habit or between IL-17A and IL-17R expression in both central and distal airway epithelial cells from smoking COPD patients. No positive correlation were found for the cells immunoreactive for IL-17F. These findings leave to suppose a secondary role for IL-17F in the pathogenesis of COPD. On the other hand, although it was observed that IL-17F may also have similar neutrophil-promoting effects in chronic inflammatory and allergic lung disease [31], an animal model of lung inflammation showed that the administration of anti-IL-17F antibodies has no effect on lung neutrophilia (pathogenic mechanism involved in airway inflammation of COPD patients) [32]. Accordingly, with ambiguous role of IL-17F, Hizawa et al. [31] showed that Mutant IL-17F was unable to activate the Raf1-MEK-ERK1/2 pathway, but antagonized wild-type IL-17F activity, suggesting that IL-17F might be able to



**Fig. 9.** Effects of CSE and rhIL-17A on cell proliferation in 16HBE and on apoptosis in A549. (A) 16HBE, cultured in the presence and absence of CSE 2.5% alone or in combination with rhIL-17A 20 ng/ml for 24 h, were used for evaluating proliferation using a CFSE test (see Section 2 for details). Bars represent mean  $\pm$  S. D. fluorescence intensity (MFI) of three separate experiments and were plotted as fold-change compared to untreated cells, which were chosen as the reference sample. (B) A representative flow cytometry of cell proliferation was shown (C) A549, cultured in the presence and absence of CSE 5% alone or in combination with rhIL-17A 20 ng/ml for 24 h were used for evaluating apoptosis using an Annexin V Test (see Section 2 for details). Bars represent mean  $\pm$  S. D. fluorescence intensity (MFI) of three separate experiments and were plotted as fold-change compared to untreated cells, which were chosen as the reference sample. (D) A representative flow cytometry of cell apoptosis was shown. Statistical analysis was performed by ANOVA test with Fisher's correction for multiple comparisons. Significance was set at  $p < 0.05$ .

bind the receptor, but not activate the signaling pathways. There has been speculation whether both IL-17A and IL-17F signal act via the IL-17R although IL-17F has at least an order of magnitude lower affinity for IL-17R than IL-17A despite the similar biological activity, [33]. Given that asthma and COPD are complex diseases involving a number of genetic and environmental factors, the genetic impact of IL-17F H161R with regard to the development of chronic airway inflammation likely varies among individuals with different genetic backgrounds and environmental exposures [34]. All together, these observations might support the concept that IL-17F present in the airways of smoking COPD might not be involved in the activation of airway epithelial cells during inflammation of COPD patients. However further study might be

necessary to clarify the role of IL-17F and IL-17R in the pathogenesis of airway diseases. In this scenario, we suggest that Th17 immunity involved in the activation of airway epithelial cells during the inflammatory actions of cigarette smoke is represented by IL-17A. Particularly we suppose that IL-17A detected in the epithelium (lower levels than subepithelium) from COPD patients, is IL-17A produced in the submucosa of central airways and in the parenchyma of distal airways by infiltrating cells to target epithelial cells in the IL-17R (highly expressed). By exploiting the growing understanding of the epithelium and its interactions with inflammatory cells obtained by our findings might open new pharmacological perspective to treat the epithelial dysregulation associated with Th17 immunity during inflammatory lung conditions.

Chronic cigarette smoke has been shown to induce both Th1 and Th17 cells in parenchymal tissue of mice [35] suggesting that this tissue might represent an important source of IL-17A/F in individuals who smoke. To study the role of cigarette smoke on IL-17A, IL-17F and IL-17R expression in epithelial cells and in inflammatory cells from central and distal airways of the lung, we stimulated epithelial cell line 16HBE, A549 (representing epithelial cell line from bronchial and distal airways, respectively) as well as PBMC from normal donors with different concentration of CSE. We showed that CSE was able to increase the production of IL-17F and IL-17R in 16HBE and in A549, while increase IL-17A and IL-17F in PBMC. This effect was obtained at different concentration of CSE: 2.5% for 16HBE, 5% for A549 and 10% for PBMC. These findings might support the concept that IL-17 immunity and the related activation in the lung, might be regulated by smoking habit (pack years) in COPD patients.

Th17 cells may play an important role in the pathogenesis of emphysema, and Th1 and Th2 cells may have contributory roles in the central rather than distal airways of COPD. Th17 cells are credited for causing and sustaining tissue damage in diverse situations. The Th1 pathway antagonizes the Th17 pathway in an intricate fashion. The evolution of the understanding of Th17 pathway illuminates a shift in immunologists' perspectives regarding the basis of tissue damage [36]. Much of our knowledge from the interactions between environmental and inflammatory stimuli, and the airway epithelium has been derived extensively from in vitro cell culture models. Taken together, in vitro studies have shown that differentiated cell culture is an invaluable model in understanding the physiological properties of the human airway epithelium. Accordingly, with this observation using an in vitro model, we studied the action of hrIL-17A on cell proliferation and apoptosis in 16HBE and in A549, with the aim to understand its different role in central and distal airways during the pathogenesis of COPD. Particularly, we found that in the presence of CSE, promoting higher levels of IL-17R receptor expression (as observed), hrIL-17A is able to increase proliferative mechanism of bronchial epithelial cell line 16HBE. These findings might suggest that epithelium proliferation, at baseline of the mechanism of squamous metaplasia observed in central airways of COPD patients [37], is a considerably disturb of the innate immune functions of the airway epithelium that might involve IL-17A. Furthermore, it was observed that IL-17A is essential to the development of emphysema in mice [38] and that cigarette smoke induces apoptosis in A549 [39]. Accordingly, we observed that CSE or hrIL-17A generate the mechanism of apoptosis in type II alveolar epithelial cell-derived A549 cell line with a synergistic effect when hrIL-17A and CSE were used in combination. These last findings suggest that CSE stimulation might be able to potentiate the action of hrIL-17A in 16HBE and A549 cell line in agreement with the increased expression of IL-17R expression. Duan et al. showed that Th17 immunity is involved in cigarette smoke-induced emphysema observed in the lung of mice [30]. Accordingly, our data might underline the relevant contribution of IL-17A in the development of pulmonary emphysema in COPD patients.

In conclusion, Th17 immunity associated with IL17A and IL-17F cytokines and the related IL-17 receptor showed a relevant role in the epithelium and subepithelium of both, central and distal airway from smoking COPD patients. Using our ex vivo/in vitro studies we showed that smoke habit might contribute to increase IL-17F and IL-17R production in epithelial cells as well as to increase IL-17A and IL-17F production in infiltrating cells present in the subepithelium from large and small airways. Finally, by in vitro study, we identified a potential role of IL-17A rather than IL-17F in the phenomena of epithelial cell proliferation and apoptosis involved in the tissue damage and repair in the central and distal airways of COPD patients.

## Funding source

The study was supported by Italian National Research Council, Unit: "Ex vivo/In vitro models to study the Immunopathology and Pharmacology of airway diseases", Institute of Biomedicine and Molecular Immunology (IBIM), Palermo, Italy.

## Disclosure statements

The authors of the manuscript have not conflict of interest.

## Authors' contributions

The authors MP, AMM and LR conceived the study and designed the experiments. LS, GC, GA, RG, CDS performed the technical procedures. FLMR and VS give us some suggestions about the data analysis. PV and LP give us the support for surgical specimens. MP revised the final draft of the manuscript. All authors read and approved the final version of the MS.

## Acknowledgments

The authors thank the Italian National Research Council and ISMETT for the support.

## References

- [1] P.J. Barnes, Chronic obstructive pulmonary disease, *N. Engl. J. Med.* 343 (2000) 269–280.
- [2] Global initiative for chronic obstructive lung disease, Global strategy for the diagnosis, management and prevention of chronic obstructive pulmonary disease, NHLBI/WHO workshop report. NIH Publication No. 2701:1-100. Bethesda, MD: National Heart, Lung and Blood Institute, April 2001, Last update 2008, <<http://www.goldcopd.com>> (accessed 12.03.2009).
- [3] A. Di Stefano, G. Caramori, F.L.M. Ricciardolo, A. Capelli, I.M. Adcock, C.F. Donner, Cellular and molecular mechanisms in chronic obstructive pulmonary disease: an overview, *Clin. Exp. Allergy* 34 (2004) 1156–1167.
- [4] M.G. Cosio, J. Majo, M.G. Cosio, Inflammation of the airways and lung parenchyma in COPD: role of T cells, *Chest* 121 (2002) 160S–165S.
- [5] S.H. Lee, S. Goswami, A. Grudo, L.Z. Song, V. Bandi, S. Goodnight-White, et al., Antielastin autoimmunity in tobacco smoking-induced emphysema, *Nat. Med.* 13 (2007) 567–569.
- [6] H. Ikeuchi, T. Kuroiwa, N. Hiramatsu, Y. Kaneko, K. Hiromura, K. Ueki, et al., Expression of interleukin-22 in rheumatoid arthritis: potential role as a proinflammatory cytokine, *Arthritis Rheum.* 52 (2005) 1037–1046.
- [7] K. Wolk, E. Witte, K. Warszawska, G. Schulze-Tanzil, K. Witte, S. Philipp, et al., The Th17cytokine IL-22 induces IL-20 production in keratinocytes: a novel immunological cascade with potential relevance in psoriasis, *Eur. J. Immunol.* 39 (2009) 3570–3581.
- [8] C.E. Jones, K. Chan, Interleukin-17 stimulates the expression of interleukin-8, growth-related oncogene-alpha, and granulocyte-colony-stimulating factor by human airway epithelial cells, *Am. J. Respir. Cell Mol. Biol.* 26 (2002) 748–753.
- [9] C.Y. Kao, F. Huang, Y. Chen, P. Thai, S. Wachi, C. Kim, et al., Up-regulation of CC chemokine ligand 20 expression in human airway epithelium by IL-17 through a JAK-in-dependent but MEK/NF-kappaB-dependent signaling pathway, *J. Immunol.* 175 (2005) 6676–6685.
- [10] C. Nembrini, B.J. Marsland, M. Kopf, IL-17-producing T cells in lung immunity and inflammation, *J. Allergy Clin. Immunol.* 123 (2009) 986–994, <http://dx.doi.org/10.1016/j.jaci.2009.03.03> (Review).
- [11] A. Lindén, H. Hoshino, M. Laan, Airway neutrophils and interleukin-17, *Eur. Res. J.* 15 (2000) 973–977.
- [12] L.Y. Hung, S. Velichko, F. Huang, P. Thai, R. Wu, Regulation of airway innate and adaptive immune responses: the IL-17 paradigm, *Critical Rev. Immunol.* 28 (2008) 269–279.
- [13] A. Di Stefano, G. Caramori, I. Gnemmi, M. Contoli, C. Vicari, A. Capelli, et al., T helper type 17-related cytokine expression is increased in the bronchial mucosa of stable chronic obstructive pulmonary disease patients, *Clin. Exp. Immunol.* 2009 (157) (1960) 316–324, <http://dx.doi.org/10.1111/j.1365-2249.2009.03965.x>, PubMed PMID:4272; PubMed Central PMCID: PMC2730858.
- [14] Y. Chang, J. Nadigel, N. Boulais, J. Bourbeau, F. Maltais, D.H. Eidelman, et al., CD8 positive T cells express IL-17 in patients with chronic obstructive pulmonary disease, *Respir. Res.* 10 (2011) 12–43, <http://dx.doi.org/10.1186/1465-9921-12-43>, PubMed PMID: 21477350; PubMed Central PMCID: PMC3082241.
- [15] A. Eustace, L.J. Smyth, L. Mitchell, K. Williamson, J. Plumb, D. Singh, Identification of cells expressing IL-17A and IL-17F in the lungs of patients

- with COPD, *Chest* 139 (2011) 1089–1100. doi: 0.1378/chest.10-0779. Epub 2010 Sep 23. PubMed PMID: 20864612..
- [16] R. Pauwels, Global initiative for chronic obstructive lung diseases (GOLD): time to act, *Eur. Respir. J.* 18 (2001) 901–902.
- [17] M. Saetta, A. Di Stefano, G. Turato, F.M. Facchini, L. Corbino, C.E. Mapp, et al., CD8<sup>+</sup> T-lymphocytes in peripheral airways of smokers with chronic obstructive pulmonary disease, *Am. J. Respir. Crit. Care Med.* 157 (1998) 822–826.
- [18] G. Chiappara, P. Chanez, A. Bruno, E. Pace, F. Pompeo, J. Bousquet, et al., Variable p-CREB expression depicts different asthma phenotypes, *Allergy* 62 (2007) 787–794.
- [19] D. Miotto, M.D. Hollenberg, N.W. Bunnett, A. Papi, F. Braccioni, P. Boschetto, et al., Expression of protease activated receptor-2 (PAR-2) in central airways of smokers and non-smokers, *Thorax* 57 (2002) 146–151.
- [20] A.L. Cozens, M.J. Yezzi, M. Yamaya, D. Steiger, J.A. Wagner, S.S. Garber, et al., A transformed human epithelial cell line that retains tight junctions post crisis, *In Vitro Cell. Dev. Biol.* 28A (1992) 735–744.
- [21] A. Bruno, S. Conus, I. Schimid, H.U. Simon, Apoptotic pathways are inhibited by leptin receptor activation in neutrophils, *J. Immunol.* 174 (2005) 8090–8096.
- [22] A.M. Montalbano, G. Anzalone, G.D. Albano, C. Di Sano, R. Gagliardo, A. Bonanno, L. Riccobono, G. Nicolini, E. Ingrassia, M. Gjomarkaj, M. Profita, Beclomethasone dipropionate and formoterol reduce oxidative/nitrosative stress generated by cigarette smoke extracts and IL-17A in human bronchial epithelial cells, *Eur. J. Pharmacol.* 15 (2013) 418–427. <http://dx.doi.org/10.1016/j.ejphar.2013.08.001> Epub.
- [23] G.D. Albano, C. Di Sano, A. Bonanno, L. Riccobono, R. Gagliardo, P. Chanez, M. Gjomarkaj, A.M. Montalbano, G. Anzalone, S. La Grutta, F.L. Ricciardolo, M. Profita, Th17 immunity in children with allergic asthma and rhinitis: a pharmacological approach, *PLoS One* 8 (4) (2013) e58892, <http://dx.doi.org/10.1371/journal.pone.0058892>. Epub 2013 Apr 3, PubMed PMID: 23573194; PubMed Central PMCID: PMC3616002..
- [24] O. Palomares, B. Rückert, T. Jartti, U.C. Küçüksezer, T. Puhakka, E. Gomez, H.B. Fahrner, A. Speiser, A. Jung, W.W. Kwok, L. Kalogjera, M. Akdis, C.A. Akdis, Induction and maintenance of allergen-specific FOXP3<sup>+</sup> Treg cells in human tonsils as potential first-line organs of oral tolerance, *J. Allergy Clin. Immunol.* 129 (2012) 510–520. 520.e1–9. <http://dx.doi.org/10.1016/j.jaci.2011.09.031>, Epub 2011 Nov 3.
- [25] P. Pochard, B. Vickery, M.C. Berin, A. Grishin, H.A. Sampson, M. Caplan, K. Bottomly, Targeting toll-like receptors on dendritic cells modifies the T(H)2 response to peanut allergens in vitro, *J. Allergy Clin. Immunol.* 126 (2010). 92–97.e5 <http://dx.doi.org/10.1016/j.jaci.2010.04.003>, Epub 2010 Jun 9, PubMed PMID: 20538332; PubMed Central PMCID: PMC2902661.
- [26] R.G. Crystal, S.H. Randell, J.F. Engelhardt, J. Voynow, M.E. Sunday, Airway epithelial cells: current concepts and challenges, *Proc. Am. Thorac. Soc.* 5 (2008) 772–777.
- [27] D.A. Knight, S.T. Holgate, The airway epithelium: structural and functional properties in health and disease, *Respirology* 8 (2003) 432–446.
- [28] W. Al-Ramli, D. Préfontaine, F. Chouiali, J.G. Martin, R. Olivenstein, C. Lemièrre, et al., T(H)17-associated cytokines (IL-17A and IL-17F) in severe asthma, *J. Allergy Clin. Immunol.* 123 (2009) 1185–1187, <http://dx.doi.org/10.1016/j.jaci.2009.02.024>. Epub 2009 Apr 10.
- [29] Y. Chang, L. Al-Alwan, S. Alshakfa, S. Audusseau, A. Mogas, F. Chouiali, P. Nair, C.J. Baglole, Q. Hamid, D.H. Eidelman, Upregulation of IL-17A/F from human lung tissue explants with cigarette smoke exposure: implications for COPD, *Respir. Res.* 27 15(1) (2014) 145. [Epub ahead of print] PubMed PMID: 25427574; PubMed Central PMCID: PMC4268887.
- [30] M.C. Duan, X.N. Zhong, Y. Huang, Z.Y. He, H.J. Tang, Mechanisms and dynamics of Th17 cells in mice with cigarette smoke-induced emphysema, *Zhonghua Yi Xue Za Zhi.* 26 (2011) 1996–2000 (Chinese).
- [31] N. Hizawa, M. Kawaguchi, S.K. Huang, M. Nishimura, Role of interleukin-17F in chronic inflammatory and allergic lung disease, *Clin. Exp. Allergy* 36 (2006) 1109–1114.
- [32] S.C. Liang, A.J. Long, F. Bennett, M.J. Whitters, R. Karim, M. Collins, et al., An IL-17F/A heterodimer protein is produced by mouse Th17 cells and induces airway neutrophil recruitment, *J. Immunol.* 179 (2007) 7791–7799.
- [33] S.G. Hymowitz, E.H. Filvaroff, J.P. Yin, J. Lee, L. Cai, P. Risser, M. Maruoka, W. Mao, J. Foster, R.F. Kelley, G. Pan, A.L. Gurney, A.M. de Vos, M.A. Starovsknik, IL-17s adopt a cystine knot fold: structure and activity of a novel cytokine, IL-17F, and implications for receptor binding, *EMBO J.* 20 (2001) 5332–5341. PubMed PMID: 11574464; PubMed Central PMCID: PMC125646.
- [34] Y. Chang, L. Al-Alwan, S. Audusseau, F. Chouiali, J. Carlevaro-Fita, Y. Iwakura, et al., Genetic deletion of IL-17A reduces cigarette smoke-induced inflammation and alveolar type II cell apoptosis, *Am. J. Physiol. Lung Cell Mol. Physiol.* 306 (2014) L132–L143, <http://dx.doi.org/10.1152/ajplung.00111.2013>. Epub 2013 Oct 4.
- [35] O.J. Harrison, J. Foley, B.J. Bolognese, E. Long 3rd, P.L. Podolin, P.T. Walsh, Airway infiltration of CD4<sup>+</sup> CCR6<sup>+</sup> Th17 type cells associated with chronic cigarette smoke induced airspace enlargement, *Immunol. Lett* 121 (2008) 13–21.
- [36] L. Steinman, A brief history of T(H)17, the first major revision in the T(H)1/T(H)2 hypothesis of T cell-mediated tissue damage, *Nat. Med.* 13 (2007) 139–145. Review. Erratum in: *Nat. Med.* 13(3) (2007) 385. PubMed PMID: 17290272.
- [37] E. Puchelle, J.M. Zahm, J.M. Tournier, C. Coraux, Airway epithelial repair, regeneration, and remodeling after injury in chronic obstructive pulmonary disease, *Proc. Am. Thorac. Soc.* 3 (2006) 726–733. Review.
- [38] Z.X. Jiao, Q.L. Ao, M. Xiong, Cigarette smoke extract inhibits the proliferation of alveolar epithelial cells and induces apoptosis, *Sheng Li Xue Bao. J* 58 (2006) 244–254. PubMed PMID: 16786.
- [39] E. Kurimoto, N. Miyahara, A. Kanehiro, K. Waseda, A. Taniguchi, G. Ikeda, et al., IL-17A is essential to the development of elastase-induced pulmonary inflammation and emphysema in mice, *Respir Res* 20 (14) (2013) 5. <http://dx.doi.org/10.1186/1465-9921-14-5>. PubMed PMID: 23331548; PubMed Central PMCID: PMC3564829.