

The sound of longevity: music and technology for healthy ageing

Received: 22 January 2026

Accepted: 29 April 2026

Published online: 09 May 2026

Cite this article as: Ecarnot F., Brattico E., Camurri A. *et al.* The sound of longevity: music and technology for healthy ageing. *Aging Clin Exp Res* (2026). <https://doi.org/10.1007/s40520-026-03410-7>

Fiona Ecarnot, Elvira Brattico, Antonio Camurri, Andrea Cera, Simone Dalla Bella, Massimo Grassi, Stefan Koelsch, Wanda Morganti, Alois Saller, Cornel Sieber, Fiorella Tonello, Peter Vuust, Stefania Maggi & Alberto Pilotto

We are providing an unedited version of this manuscript to give early access to its findings. Before final publication, the manuscript will undergo further editing. Please note there may be errors present which affect the content, and all legal disclaimers apply.

If this paper is publishing under a Transparent Peer Review model then Peer Review reports will publish with the final article.

ARTICLE IN PRESS

The Sound of Longevity: Music and Technology for Healthy Ageing

Fiona Ecarnot^{1,2}, Elvira Brattico^{3,4}, Antonio Camurri⁵, Andrea Cera⁶,
Simone Dalla Bella⁷⁻¹⁰, Massimo Grassi¹¹, Stefan Koelsch¹², Wanda
Morganti¹³, Alois Saller¹⁴, Cornel Sieber^{15,16}, Fiorella Tonello¹⁷, Peter
Vuust^{3,4}, Stefania Maggi¹⁷, Alberto Pilotto¹⁸.

Affiliations:

1. Department of Cardiology, University Hospital Besançon, Besançon, France
2. SINERGIES, University Marie & Louis Pasteur, Besançon, France
3. Center for Music in the Brain, Department of Clinical Medicine, Aarhus University, Aarhus, & Royal Academy of Music Aarhus/Aalborg, Aarhus, Denmark
4. Department of Education, Psychology, Communication, University of Bari, Italy
5. InfoMus -Casa Paganini Research Centre, University of Genoa, Italy
6. IRCAM / InfoMus-Casa Paganini Research Centre, University of Genoa, Italy
7. International Laboratory for Brain, Music and Sound Research (BRAMS), Montreal, Québec, Canada

8. Department of Psychology, Université de Montréal, Montreal, Québec, Canada
9. Centre for Research on Brain, Language and Music (CRBLM), Montreal, Québec, Canada
10. VIZJA University, Warsaw, Poland
11. Department of General Psychology, University of Padua, Padua, Italy
12. Department of Biological and Medical Psychology, University of Bergen, 5020 Bergen, Norway
13. Geriatrics Unit, Department of Medicine and Geriatrics, E.O. Galliera Hospitals, Genoa, Italy
14. Department of Medicine, Sant'Antonio Hospital, Padua, Italy
15. Institute for Biomedicine of Ageing (IBA), Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany.
16. CMO, County Hospital Winterthur, Winterthur, Switzerland.
17. CNR of Italy, Neuroscience Institute, Viale G. Colombo 3, 35131 Padua, Italy
18. Research, Development and Scientific Coordination Unit, E.O. Galliera Hospitals, Genoa, Italy

Corresponding author:

Fiona Ecarnot, PhD

Department of Cardiology, University Hospital Besancon, 3-8 Boulevard
Fleming, 25000 Besancon, France

ARTICLE IN PRESS

Abstract

A growing body of research is focusing on how music, technology, and neuroscience can converge to promote healthy ageing and counteract pathological decline. In particular, music interventions for older adults have been garnering increasing attention, with numerous reports showing positive effects of music on various health outcomes, including psychological well-being, cognitive function, physiological responses, quality of life, and overall well-being. In this context, the European Interdisciplinary Council on Ageing (EICA) in collaboration with the "Age-It" project (Ageing well in an Ageing Society), specifically, the work package 1 of Spoke 8, run by the University of Bari, and the "RAISE" project (Robotic and Artificial Intelligence for Socio-economic Empowerment) run by the E.O. Galliera Hospitals, Genoa, Italy, convened a 2-day meeting in June 2025 to review the state of the evidence on the relationship between music and health. Speakers were tasked with exploring how music, neuroscience, technology, and environmental design contribute to understanding ageing and to developing interventions that support healthy ageing. This report presents the Executive Summary of the proceedings of this meeting.

Introduction

Ageing is a multidimensional process shaped by biological, psychological, social, and environmental factors. Across recent decades, research has increasingly focused on how music, technology, and neuroscience can converge to promote healthy ageing and counteract pathological decline. Healthy ageing is broadly and heterogeneously defined in the literature, reflecting its multidimensional and context-dependent nature, but is generally understood as a dynamic process shaped by personal characteristics, resources, goals, and the broader social, cultural, economic, and environmental context across multiple dimensions of life (1). It involves both maintaining health and function as much as possible, and adapting effectively over time through supportive interactions between the person and their environment (1).

In this context, music interventions for older adults have been garnering increasing attention in recent years, with a number of published reviews that have shown positive effects of music on various health outcomes, including psychological well-being, cognitive function, physiological responses, quality of life, and overall well-being (2,3) . There is a growing consensus that music interventions may enhance biopsychosocial health outcomes in older adults, although overall confidence in the evidence remains quite low. According to a white paper by Jespersen and colleagues, the current evidence corresponds to Level 2 in the Oxford Centre for Evidence-Based Medicine hierarchy, indicating support from randomized or large observational studies but insufficient systematic

synthesis to draw definitive conclusions ageing(4). Similarly, while mechanistic models of the rehabilitative impact of music on neurological disorders have been put forward in the literature, a consensus is still lacking on the actual mechanisms driving the beneficial effects of music on health, especially in older adults (5-7). Nevertheless, music is a valuable and highly versatile support for the well-being of older adults. In today's world, where life expectancy is rising, technological advancements have transformed how we experience music, and thus the role of both music and technology in the daily life of older individuals deserves greater scientific attention.

To this end, the European Interdisciplinary Council on Ageing (EICA) in collaboration with the "Age-It" project (Ageing well in an Ageing Society), specifically the work package 1 of Spoke 8 run by the University of Bari, and the "RAISE" project (Robotic and Artificial Intelligence for Socio-economic Empowerment) run by the E.O. Galliera Hospitals, Genoa, Italy, convened a 2-day international meeting entitled "The Sound of Longevity: Music and Technology for a Health Ageing"(including a live streaming webinar, available at <https://www.youtube.com/watch?v=Cz2CpZkMrIE>, and a workshop) in June 2025. The scope of the meeting was to review the state of the evidence on the relationship between music, technology, and health, thus, speakers were tasked with exploring how music, technology, neuroscience, and environmental design contribute to understanding ageing and to developing interventions that support healthy ageing (see Figure 1). We present here the Executive Summary of the proceedings of this meeting.

2. A Multidimensional Approach to Ageing: Technology and Music

Overview of multidimensional assessment models

A multidimensional understanding of ageing is essential for designing and evaluating effective interventions. While it is known that biological age alone does not adequately capture the great heterogeneity of ageing trajectories, the Comprehensive Geriatric Assessment (CGA) is recognized as a cornerstone of the Geriatric Medicine and the gold standard for evaluating medical, cognitive, functional, nutritional, and psychosocial domains in older adults (8). A central tool derived from the CGA is the Multidimensional Prognostic Index (MPI). By integrating standardized measurements across domains, the MPI predicts mortality and other adverse outcomes with high accuracy. Importantly, the MPI supports clinical decision-making by identifying risk profiles and informing personalized care plans (9). A great number of studies have confirmed its validity and utility across hospital, primary care, and telemedicine contexts (10).

Several major ongoing projects illustrate how CGA and the MPI can be combined with technology and lifestyle interventions. The MULTIPLAT-AGE project linked multiple technological platforms to create interoperable data systems supporting personalized treatment pathways (11). Examples of such platforms include an adapted physical activity programme supported by information and communication technologies, or at-home cognitive telerehabilitation / remote cognitive stimulation (11). The

Pro-Home project further demonstrated how home-based monitoring, incorporating domotics, robotics, and music training, can help maintain mobility, reduce frailty, and sustain improvements, up to several months after the intervention ends (12).

Role of technology and music in supporting healthy ageing

Music appears as an emerging component in multidomain interventions. Evidence from numerous systematic reviews suggests that music listening, exercise with music, and structured music therapy improve mood, physical activity, socialization, and aspects of cognitive function (13). The mechanisms remain under investigation (5-7), but involve interactions among motor networks, emotional regulation circuits, and neuroendocrine pathways. The integration of music into CGA-based interventions is conceptually aligned with person-centred geriatric care: it targets multiple domains simultaneously, supports adherence, and enhances quality of life.

3. Neural Perspectives on Music Across the Lifespan

The importance of music for humans resides in the phenomenon of group coherence. Indeed, it creates an ambiance conducive to cooperation, and fosters group coordination and promote prosocial behaviour (14,15).

Previous research has shown that interpersonal synchrony, in the form of synchronized drumming, enhances activity in the caudate nucleus in the brain (16). This activates reward-like signals that have the potential to shape future behaviour, specifically by promoting prosocial commitment (17). It has also been shown that temporally aligned movements enhance

social bonding, and this in turn was associated with higher empathy during movement interactions that featured music (18,19).

The question of why humans spontaneously move when they hear music has been the subject of intense research, and there is evidence that most humans can move to the beat, musicians and non-musicians alike, albeit with some exceptions (20,21). This concept is known as "groove", namely the "feel" of the rhythm, or the subtle quality of music that compels people to dance or move along (22). The balance achieved between the degree of syncopation of the music, and the feelings of pleasure and wanting to move, indicate that both motor and pleasure systems (like dopamine) are stimulated when a human listens to music. These findings regarding activation of the associated brain areas have been corroborated by functional magnetic resonance imaging studies performed while musicians and non-musicians listened to and rated experimentally controlled music (23). The study showed that rhythms with a medium level of complexity elicited greater pleasure and a greater sensation of desire to move (23). Cultural differences and musicianship experience may affect these responses (24,25). Interestingly, the disruption to the dopaminergic system resulting from drug addiction can cause individuals with substance use disorders to experience stronger groove with high rhythmic and harmonic complexities (26). High rhythmic complexity means the beat is hard to predict or follow, i.e. due to the use of unusual time signatures, syncopation, overlapping patterns, or sudden accents. Harmonic complexity means the chords and note combinations are richer or less familiar than in simpler songs, i.e. the music may change key

unexpectedly, use dense or dissonant chords, or move through chord progressions that create ambiguity or tension. Furthermore, individuals with Parkinson's disease have also been shown to have a significantly flattened response for both pleasure and the urge to move (27).

Current thinking suggests that the violation and confirmation of temporal predictions is the key driver behind the pleasurable effect of music.

Accordingly, the inverted U-shaped relation between the urge to dance and rhythmic complexity is purported to represent a balance between predictability and uncertainty when listening to the music (28). In this regard, the predictive coding theory proposes that prediction is a fundamental principle of brain processing (29).

Auditory memory, which encompasses melody and harmony in addition to rhythm, has also been shown to be affected by ageing. In one study, it was shown that ageing differentially affects the various dimensions of auditory memory, whereby more musical training conferred a significant advantage on certain specific tasks (30). Overall, it was shown that younger participants (<25 years of age) outperformed their older counterparts (>60 years) in terms of short term memory for rhythmic sequences and long-term memory for novel sequences (30). Conversely, for previously memorized music, older adults displayed stronger activity in auditory regions. Therefore, there is clearly a difference in strategy, with older people relying more on auditory cortices, while younger people rely more on hippocampal areas (31). When it comes to encoding new melodies, older adults have reduced activity in regions relating to decision

making processes, related to novelty, and display reduced performance compared to younger adults. Such age-related differences in neural strategies for processing music underscore that brain function across the lifespan reflects both experiential learning and foundational developmental influences, making it essential to consider how epigenetic mechanisms shape these trajectories from the earliest stages.

Epigenetic Risk Factors in Neurodevelopment

Engaging with the multisensory stimulus that music represents can prompt structural and functional adaptations within the brain, and ultimately influence neural pathways and neuroplasticity (32). In this regard, it has been proposed that resilience, i.e., the ability of the brain to maintain or recover function after stress or insult, emerges from the lifelong interaction between biological predisposition and the environmental context. Accordingly, if resilience depends on the plastic potential of neural circuits, genetic variability, and environmental inputs (33), then music could be leveraged as a tool to promote resilience. Within this context, the Italian National Research Council (CNR) has developed LOTUS³ - a unifying initiative linking diverse research clusters focused on brain health, environmental exposure, and neuroplasticity. Directed by Professor Michela Fagiolini (CNR and Harvard Medical School), LOTUS³ unites research on molecular neuroscience, behavioral interventions, and environmental toxicology under a single vision, namely optimizing brain trajectories from early development to old age.

Music-Related Brain Changes in Ageing

Building on this framework of environmentally driven brain trajectories, music offers a compelling example of how specific experiences can modulate neural systems throughout adulthood and into older age. Accumulating evidence supports the role of musical engagement in enhancing brain plasticity, supporting cognitive reserve, and slowing structural decline. A comprehensive review by James et al. demonstrated broad cognitive protection through musical interventions, with the strongest effects observed in combined cognitive and physical training paradigms (34). Here, cognitive training paradigms include interventions targeting functions such as memory, attention, executive control, reasoning, and spatial navigation, often delivered through computerized tasks, games, language learning, or music-based activities; physical training paradigms include aerobic exercise and non-aerobic coordinative or sensorimotor activities such as walking, cycling, dancing, resistance training, tai chi, and balance-based exercise (34). Similarly, Marie et al. reported that both piano training and active music listening led to increases in cerebellar grey matter volume and auditory working memory, despite ongoing age-related atrophy (35). Of note, both piano training and active music listening had effects. Mack et al. found that one year of piano training enhanced cognitive flexibility and sustained attention in older adults (36), while Jünemann et al. demonstrated stabilization of white-matter integrity in the fornix—a key memory-related structure—

after six months of piano training (37). These converging results highlight the promising neuroprotective effects of music in ageing, with benefits from everyday cognitive functioning to structural brain integrity.

Musical engagement may also slow progression in pathological conditions such as Alzheimer's disease (AD). Despite profound impairment of episodic and semantic memory, individuals with AD often retain the ability to recognize familiar melodies and even learn new ones. AD patients have been shown capable of distinguishing familiar from novel melodies, detecting wrong notes, and singing familiar tunes accurately (38).

Elsewhere, it was reported that AD patients who learned sung lyrics had better word retention than spoken presentation, underscoring the mnemonic power of melodic structure (39). Moreover, music has been shown to enhance autobiographical recall: patients recalled life episodes more vividly when listening to familiar music, even when the pieces were unrelated to those experiences (40-42). Current thinking suggests that a separate musical memory exists that can exert positive effects, whereby learning and retention of musical information is mediated by brain networks distinct from those involved in other types of episodic and semantic memory (43,44). Moreover, the acquisition and recall of complex musical information could remain as intact cognitive activities, even in the presence of other memory impairment (45). Together, these findings suggest that neural systems supporting musical memory might be resistant to neurodegeneration until its latest stages (46,47).

4. Music-Based Interventions in Pathological Ageing

Music-based interventions represent a low-cost, accessible, and neurobiologically grounded approach to mitigating cognitive and emotional decline in ageing. By preserving hippocampal volume, stimulating dopaminergic circuits, and enhancing emotional well-being, music therapy and music-based interventions might complement physical and cognitive training strategies in the prevention and management of dementia.

Recent systematic reviews and meta-analyses indicate that music-based interventions may offer modest cognitive benefits for individuals with dementia, although findings remain mixed. Fusar-Poli et al. reported no overall significant effects of music therapy across outcomes, but subgroup analyses suggested a small positive effect of active interventions on global cognition (48). Similarly, Dorris et al. found that music-making produced a small yet statistically significant improvement in cognitive functioning among older adults with mild cognitive impairment or dementia (49). Despite these signals of benefit, the broader evidence base is constrained by substantial methodological weaknesses. Reviews highlight consistently small sample sizes, heterogeneous and often non-standardized intervention protocols, and variability in outcome measures, all of which limit interpretability (50-52). There remains a compelling need for rigorously designed, larger randomized controlled trials that incorporate well-defined and stage-appropriate intervention protocols,

assess both cognitive and positive emotional or social outcomes, and evaluate treatment duration and persistence of effects (51,53).

Beyond AD, multiple other studies have documented music's influence on physiological markers such as heart rate, heart-rate variability (HRV), respiration, blood pressure, and sleep quality. Krabs and colleagues demonstrated that both pleasant and unpleasant music, and even isochronous tones devoid of melody, reduce HRV compared to silence (54). The similarity of responses across emotional valences indicates that rhythm and temporal regularity play dominant roles in autonomic regulation. Orini et al. further confirmed that the tactus or beat alone is sufficient to elicit measurable autonomic responses, emphasizing entrainment as a key mechanism (55). Yet, as in the field of AD research, clinical application of music therapy faces methodological challenges, since the choice of tempo, familiarity with the music, the presence of lyrics, and the delivery mode (headphones, live, etc) are all factors that may influence outcomes. Furthermore, distinguishing between active music-making and passive listening is essential, as active participation engages motor and cognitive circuits that enhance plasticity and reward processing.

Among the potential benefits of active musical engagement, the capacity to synchronize movement to an external beat can be strategically harnessed for rehabilitation purposes. Rhythmic abilities may be thought to be characteristic of experts or musicians, but in fact, the capacity to couple movement to rhythm is very widespread and natural in the general

population (21,56). This ability is supported by dedicated neural circuitry, encompassing networks of cortical and subcortical regions whose interactions enable the temporal alignment of motor actions with rhythmic auditory cues (57,58). A particularly noteworthy point is that the neural regions engaged in rhythm processing are primarily those responsible for motor control, rather than areas specialized exclusively for rhythm. Even tasks that require only the perception or identification of a beat, without overt movement, elicit robust activation of motor circuits (59-61). This phenomenon reflects the strong functional coupling between rhythm perception and motor control, whereby auditory rhythmic input can stimulate motor networks even in the absence of actual movement. Building on these findings, a translational approach has emerged, moving from basic research to applied rehabilitation by developing theory-driven interventions (62).

A useful model for studying rhythm-based interventions in rehabilitation is Parkinson's disease (PD), where motor impairments and disrupted timing processes provide a clear framework for testing the effects of rhythmic stimulation (63,64). PD is a neurodegenerative disease that results from neuronal depletion in the substantia nigra and deregulation of the dopaminergic system with characteristic symptoms: akinesia (few initiated movements), bradykinesia (slow movements), rigidity, and tremor at rest. PD is also characterized by rhythm disturbances, often referred to as a general dysrhythmia(65), manifesting behaviorally as impaired rhythmic motor timing. For example, this is visible in a tendency to "hasten" during paced tapping when auditory cues are removed and

timing must be internally maintained. More recently, in a study by Puyjarinet et al. (66), it was found that general dysrhythmia is visible in patients with PD on the production side across different motor tasks such as finger tapping, oro-facial coordination, and gait, as well as in rhythm perception (e.g., in rhythm discrimination, the detection of tempo changes, and the detection of a misaligned beat). Notably, rhythm deficits in gait, vocal production, and manual motor control can be predicted by deficits in rhythmic perception. Another major challenge in PD is the presence of gait disturbances, which are associated with a high burden but respond poorly to standard pharmacological treatment, such as L-dopa (67). Evidence indicates, however, that rhythmic stimulation can significantly improve gait performance. Rhythmic auditory cueing, particularly when delivered through music, for example following the neurological music-therapy protocol named Rhythmic Auditory Stimulation (RAS), enables patients to walk faster, with longer steps and a more natural gait pattern (62,68-71). The effects are often immediate, but they typically dissipate once the stimulus is removed. Nevertheless, when cueing-based training is applied consistently over a period of one to two months, benefits can persist even in the absence of ongoing external stimulation (63,72,73).

5. Music- and Dance-Based Technological Advances for Rehabilitation

Music and dance occupy a unique space at the intersection of art, neuroscience, and technology. At the InfoMus-Casa Paganini Research Centre of the University of Genoa, Antonio Camurri and colleagues have developed a transdisciplinary framework that merges artistic practice, computational modeling, and human-centered design. The therapeutic applications of Casa Paganini's research extend from clinical rehabilitation to cultural well-being. In the "CARE HERE" project (74), Parkinson's patients used their bodies as 'brushes' to paint projected walls, with fluid motion generating visual and sonic feedback. This aesthetic and motivational approach quantitatively reduced movement hesitation, illustrating how beauty and play can enhance motor recovery.

The "DanzArTe" Emotional Well-being Technology, developed in collaboration with geriatricians, merges interactive sonification, visual art, and virtual agents to foster emotional and physical engagement among older adults. Participants reconstruct classical paintings through expressive movement; their gestures, analyzed in real time, 'reveal' the artwork as movement becomes fluid. Recent evaluations reported significant improvements in subjective well-being and resilience, validating the paradigm of aesthetic resonance, i.e. the emotional alignment between performer, movement, and sensory environment (75,76). By capturing the nuances of expressive gesture and translating them into multisensory feedback, music- and dance-based systems foster empathy, motivation, and neuroplastic recovery.

Another example of application of technology to enhance an aesthetic experience for rehabilitative purposes comes from a recent proof-of-concept study by Carlomagno and colleagues (77). They used immersive virtual reality (VR) combined with evocative, aesthetically engaging visual content (calming, pleasant landscapes) and music (culturally resonant songs) as a mini-intervention for 21 older individuals with varying levels of cognitive decline, including mild cognitive impairment and dementia. The primary goals were to assess whether this audiovisual intervention could enhance mood, evoke autobiographical memories, and improve overall emotional well-being, regardless of participants' cognitive status. Findings indicate that most participants experienced high immersion and predominantly positive emotional responses, with many reporting calmness and recollection of meaningful personal memories during the sessions. Moreover, self-rated mood significantly improved from before to after VR exposure. Some minor side effects—such as mild anxiety or physical discomfort from the headset—were noted, but overall, the experience was engaging and well received.

Beyond immersive art and VR, digital rehabilitation technologies, such as rhythmic serious games and adaptive mobile solutions (72), offer an alternative way to make rhythm-based training scalable, engaging, and personalized for people with PD. For example, serious games exploit gamification to increase motivation while enabling multisensory engagement, real-time monitoring, and individualized control of training parameters, in a format that is affordable and deployable on tablets or smartphones. For example, in a proof-of-concept study Puyjarinet et al.,

implemented a six-week home program (3 sessions/week) using a serious rhythmic game in which patients tapped in synchrony with music to gain points and progress (66). The intervention was associated with improvements in trained rhythmic functions (e.g., reduced tapping variability) and speech articulation, alongside improved beat perception that correlated with motor outcomes, consistent with the idea that strengthening a central rhythmic system can support broader motor function.

Overall, these exemplary studies illustrate how the fusion of art and technology can transform rehabilitation into an aesthetic experience, supporting and empowering patients across diverse functional domains (78).

6. The Role of Soundscapes in Healthy Ageing

The relationship between sound and health is paradoxical: sound can both injure and heal. At the intersection of musical composition and sound design, the notion of the 'soundscape', i.e. the total acoustic environment in which we are immersed, was first introduced in Raymond Murray Schafer's seminal work, "The Tuning of the World" (79). The soundscape has emerged as an important perspective in the domain of healthcare, notably to investigate how sonic environments might influence physiological and emotional well-being, especially among vulnerable and ageing populations.

In hospital settings, the sounds in the ambient acoustic environment, e.g., ventilators, footsteps, or distant voices, can alternately either comfort or distress patients. Studies in intensive care units have shown that acoustic insulation may paradoxically create environments that are ‘too silent’, leading to disorientation or anxiety (80). Recent research has shown that the use of artificial keynotes (i.e., soft, continuous background sounds) can restore a sense of spatial and temporal orientation. Conversely, signals such as alarms are often too intrusive. Customizing alarm tones to match staff preferences and attentional thresholds can reduce cognitive fatigue, an idea supported by research on auditory ergonomics (80-82). Other soundscape interventions in healthcare can range from environmental design to individualized therapy. For example, one study performed in a French long-term care facility for people with AD, experimented with diffused soundscapes tailored to specific spaces: church bells in communal areas, birdsong near gardens, and personalized music in patient rooms. These subtle sonic cues provided cognitive and emotional anchors, stimulating memory and reducing agitation (83). This work echoes findings by Voisin et al. (83) showing that familiar environmental sounds improve orientation and emotional stability in dementia care.

7. Future Directions and Emerging Opportunities

Future work should place greater emphasis on clarifying the mechanisms through which music may support healthy ageing. In particular, predictive

processing offers a useful framework for understanding how the balance between expectation and surprise in music may shape pleasure, movement, attention, and learning, and how these effects may differ according to musical experience, ageing trajectory, and neurological status. At the same time, further research should explore the potential of personalized and technology-supported interventions across the lifespan, including adaptive music training, home-based digital programmes, immersive multisensory environments, and rehabilitation strategies integrated with multidimensional geriatric care. Methodological priorities include larger and better characterized cohorts, greater harmonization of intervention protocols and outcome measures, longer follow-up, and multimodal designs combining behavioural, clinical, and neural markers to better define both efficacy and underlying mechanisms.

Conclusion

The proceedings of the "Sound of Longevity" meeting summarized here highlight the growing recognition of music as a powerful, multidimensional resource for promoting healthy ageing according to an interdisciplinary and proactive methodological approach (see Figure 2). Across biological, cognitive, emotional, and social domains, musical engagement, whether in the form of listening, active music-making, movement, technology-mediated musical experiences or carefully designed soundscapes, emerges as a low-cost, accessible, and person-centred intervention. Advances in neuroscience have begun to elucidate the neural mechanisms through

which music supports plasticity, resilience, and preserved function, even in the presence of neurodegenerative disease. At the same time, technological innovations now enable music- and dance-based interventions to be personalized, scalable, and integrated into clinical, home, and care settings.

Despite encouraging evidence, substantial methodological challenges remain, underscoring the need for rigorous, large-scale trials and standardized outcome measures. Future research should adopt a lifespan and systems-based perspective, embedding music within comprehensive geriatric assessment frameworks to fully harness its potential in supporting longevity, well-being, and dignity in ageing populations.

Data Availability

No original data were generated in this work.

Authors' contributions:


Conception and design: SM, AP

Writing of first draft: FE, WM, SM, AP

Critical revision and approval of final version for publication: FE, EB, ACa, ACe, SDB, MG, SK, WM, AS, CS, FT, PV, SM, AP.

ARTICLE IN PRESS

Figure 1. Leaflet of the meeting “The Sound of Longevity: Music and Technology for a Healthy Ageing”, Venice, June 3rd 2025.



PROGRAM

JUNE 3, 2025

- 14.30 **Welcome & Introduction**
Elvira Brattico, Stefania Maggi
- 15.00 **Multidimensional approach to ageing: technology and the music**
Alberto Pilotto
- 15.30 **Music in the brain across the lifespan**
Peter Vuust
- 16.00 **Epigenetic risk factors in neurodevelopment**
Michela Fagiolini
- 16.30 **Music-related brain changes in ageing**
Alois Saller
- 17.00 Coffee break
- 17.15 **Music-based interventions in pathological ageing**
Stefan Koelsch
- 17.45 **Rhythm for neurological rehabilitation**
Simone Dalla Bella
- 18.15 **Technological advancement for music- and dance-based rehabilitation**
Antonio Camurri
- 18.45 **Soundscape in the healthcare for aging people**
Andrea Cera
- 19.15 **Final remarks**
- 19.30 **Conclusions**

FACULTY

- Elvira Brattico**
Aarhus University, Denmark & University of Bari, Italy
- Antonio Camurri**
University of Genoa, Italy
- Andrea Cera**
IRCAM-Institut de Recherche et Coordination Acoustique/Music, Paris, France & Casapaganini_Infomus, University of Genoa, Italy
- Simone Dalla Bella**
University of Montreal, Canada
- Michela Fagiolini**
Neurosciences Institute, National Research Council-CNR, Italy
- Stefan Koelsch**
University of Bergen, Norway
- Stefania Maggi**
Neurosciences Institute, National Research Council-CNR
EICA President
- Alberto Pilotto**
E.O. Galliera Hospitals Genoa, Italy – EICA General Secretary
- Alois Saller**
A.O. University of Padua, Italy
- Peter Vuust**
Aarhus University & Royal Academy of Music Aarhus/Aalborg, Denmark
- ADDITIONAL PARTICIPANTS (in presence or online)**
- Fiona Ecarnot**
SINERGIES, Université Marie et Louis Pasteur, Besançon, France
- Wanda Morganti**
E.O. Galliera Hospitals, Genoa, Italy
- Fiorella Tonello, Marianna Noale, Federica Limongi, Zaira Romeo, Eleonora Macchia, Paola Siviero, Barbara Corso**
Neurosciences Institute, National Research Council-CNR, Italy

Figure 2: interdisciplinary approach to the role of music in promoting and preserving healthy ageing.



References

- [1] Menassa M, Stronks K, Khatmi F, Roa Diaz ZM, Espinola OP, Gamba M, Itodo OA, Buttia C, Wehrli F, Minder B, Velarde MR, Franco OH. Concepts and definitions of healthy ageing: a systematic review and synthesis of theoretical models. *EClinicalMedicine*. 2023; 56:101821
- [2] Ma G, Ma X. Music Intervention for older adults: Evidence Map of Systematic Reviews. *Medicine (Baltimore)*. 2023; 102:e36016
- [3] Raja S, Barry C, Upadhyay R, Alash R, O'Raghallaigh M, Hayes R, Romero-Ortuno R. Harmonious ageing: a narrative review of music therapy in the biomedical literature. *Eur Geriatr Med*. 2025; 16:461-72
- [4] Jespersen KV, Gebauer L, Vuust P. White Paper: Music interventions in health care. Center for Music in the Brain, Aarhus University, Denmark in collaboration with SoundFocus, Oticon Medical, WS Audiology and Danish Sound Cluster, Denmark. Available at: https://musicinthebrain.au.dk/fileadmin/Musicinthebrain/MusicIntheBrain_Whitepaper_2022_web.pdf [Access date: 21 January 2026]2022.
- [5] Sihvonen AJ, Sarkamo T, Leo V, Tervaniemi M, Altenmuller E, Soinila S. Music-based interventions in neurological rehabilitation. *Lancet Neurol*. 2017; 16:648-60
- [6] Brancatisano O, Baird A, Thompson WF. Why is music therapeutic for neurological disorders? The Therapeutic Music Capacities Model. *Neurosci Biobehav Rev*. 2020; 112:600-15
- [7] Grau-Sanchez J, Munte TF, Altenmuller E, Duarte E, Rodriguez-Fornells A. Potential benefits of music playing in stroke upper limb motor rehabilitation. *Neurosci Biobehav Rev*. 2020; 112:585-99

- [8] Zampino M, Polidori MC, Ferrucci L, O'Neill D, Pilotto A, Gogol M, Rubenstein L. Biomarkers of ageing in real life: three questions on ageing and the comprehensive geriatric assessment. *Geroscience*. 2022; 44:2611-22
- [9] Cruz-Jentoft AJ, Daragjati J, Fratiglioni L, Maggi S, Mangoni AA, Mattace-Raso F, Paccalin M, Polidori MC, Topinkova E, Ferrucci L, Pilotto A, Investigators MA. Using the Multidimensional Prognostic Index (MPI) to improve cost-effectiveness of interventions in multimorbid frail older persons: results and final recommendations from the MPI_AGE European Project. *Ageing Clin Exp Res*. 2020; 32:861-8
- [10] Mattace-Raso F, Pilotto A. The challenge of the multifaceted prognosis in the older people and the Multidimensional Prognostic Index. *Eur Geriatr Med*. 2021; 12:223-6
- [11] Pilotto A, Massone C, Iaccarino G, Genazzani A, Trompetto C, Arabia G, Morganti W, Seminerio E, Illario M, Castello L, Mori L, Pignolo L, Custureri R, Investigators MA. A Digital-Health Program Based on Comprehensive Geriatric Assessment for the Management of Older People at Their Home: Final Recommendations from the MULTIPLAT_AGE Network Project. *Healthcare (Basel)*. 2025; 13
- [12] Pilotto A, Morganti W, Barbagelata M, Seminerio E, Morelli S, Custureri R, Dini S, Senesi B, Prete C, Puleo G, Berutti Bergotto C, Vallone F, Custodero C, Camurri A, Group P-HPI. A transitional care program in a technologically monitored in-hospital facility reduces the length of hospital stay and improves multidimensional frailty in older patients: a Randomized Clinical Trial. *Ageing Clin Exp Res*. 2024; 36:160

- [13] Agres KR, Schaefer RS, Volk A, van Hooren S, Holzapfel A, Dalla Bella S, Müller M, de Witte M, Herremans D, Ramirez Melendez R, Neerincx M, Ruiz S, Meredith D, Dimitriadis T, Magee WL. Music, Computing, and Health: A Roadmap for the Current and Future Roles of Music Technology for Health Care and Well-Being. *Music & Science*. 2021; 4:2059204321997709
- [14] Tarr B, Launay J, Dunbar RI. Music and social bonding: "self-other" merging and neurohormonal mechanisms. *Front Psychol*. 2014; 5:1096
- [15] Stupacher J, Maes P-J, Witte M, Wood G. Music strengthens prosocial effects of interpersonal synchronization—If you move in time with the beat. *Netherlands: Elsevier Science*; 2017. p. 39-44.
- [16] Kasdan AV, Burgess AN, Pizzagalli F, Scartozzi A, Chern A, Kotz SA, Wilson SM, Gordon RL. Identifying a brain network for musical rhythm: A functional neuroimaging meta-analysis and systematic review. *Neurosci Biobehav Rev*. 2022; 136:104588
- [17] Kokal I, Engel A, Kirschner S, Keysers C. Synchronized drumming enhances activity in the caudate and facilitates prosocial commitment--if the rhythm comes easily. *PLoS One*. 2011; 6:e27272
- [18] Stupacher J, Mikkelsen J, Vuust P. Higher empathy is associated with stronger social bonding when moving together with music. *Psychol Music*. 2022; 50:1511-26
- [19] Stupacher J, Witek MAG, Vuoskoski JK, Vuust P. Cultural Familiarity and Individual Musical Taste Differently Affect Social Bonding when Moving to Music. *Sci Rep*. 2020; 10:10015

- [20] Dalla Bella S, Janaqi S, Benoit CE, Farrugia N, Begel V, Verga L, Harding EE, Kotz SA. Unravelling individual rhythmic abilities using machine learning. *Sci Rep*. 2024; 14:1135
- [21] Sowinski J, Dalla Bella S. Poor synchronization to the beat may result from deficient auditory-motor mapping. *Neuropsychologia*. 2013; 51:1952-63
- [22] Witek MA, Clarke EF, Wallentin M, Kringelbach ML, Vuust P. Syncopation, body-movement and pleasure in groove music. *PLoS One*. 2014; 9:e94446
- [23] Matthews TE, Witek MAG, Lund T, Vuust P, Penhune VB. The sensation of groove engages motor and reward networks. *Neuroimage*. 2020; 214:116768
- [24] Witek MAG, Liu J, Kuubertzie J, Yankyera AP, Adzei S, Vuust P. A Critical Cross-cultural Study of Sensorimotor and Groove Responses to Syncopation Among Ghanaian and American University Students and Staff. *Music Perception*. 2020; 37:278-97
- [25] Klarlund M, Brattico E, Pearce M, Wu Y, Vuust P, Overgaard M, Du Y. Worlds apart? Testing the cultural distance hypothesis in music perception of Chinese and Western listeners. *Cognition*. 2023; 235:105405
- [26] Stupacher J, Matarrelli B, Cozzoli D, Ventura M, Montinaro F, de Gennaro L, Vuust P, Brattico E. Individuals with substance use disorders experience an increased urge to move to complex music. *Proc Natl Acad Sci U S A*. 2025; 122:e2502656122

- [27] Pando-Naude V, Matthews TE, Hojlund A, Jakobsen S, Ostergaard K, Johnsen E, Garza-Villarreal EA, Witek MAG, Penhune V, Vuust P. Dopamine dysregulation in Parkinson's disease flattens the pleasurable urge to move to musical rhythms. *Eur J Neurosci*. 2024; 59:101-18
- [28] Matthews TE, Stupacher J, Vuust P. The Pleasurable Urge to Move to Music Through the Lens of Learning Progress. *J Cogn*. 2023; 6:55
- [29] Koelsch S, Vuust P, Friston K. Predictive Processes and the Peculiar Case of Music. *Trends Cogn Sci*. 2019; 23:63-77
- [30] Fernández-Rubio G, Olsen ER, Klarlund M, Mallon O, Carlomagno F, Vuust P, Kringelbach ML, Brattico E, Bonetti L. Investigating the impact of age on auditory short-term, long-term, and working memory. *Psychology of Music*. 2024; 52:187-98
- [31] Bonetti L, Risgaard Olsen E, Carlomagno F, Serra E, Szabo SA, Klarlund M, Andersen MH, Frausing L, Vuust P, Brattico E, Kringelbach ML, Fernandez-Rubio G. Working Memory Predicts Long-Term Recognition of Auditory Sequences: Dissociation Between Confirmed Predictions and Prediction Errors. *Scand J Psychol*. 2025;
- [32] Kunikullaya UK, Pranjic M, Rigby A, Pallas-Ferrer I, Anand H, Kunnivil R, Jaschke AC. The molecular basis of music-induced neuroplasticity in humans: A systematic review. *Neurosci Biobehav Rev*. 2025; 175:106219
- [33] Fagiolini M, Hensch TK. Inhibitory threshold for critical-period activation in primary visual cortex. *Nature*. 2000; 404:183-6
- [34] James CE, Muller DM, Muller CAH, Van De Looij Y, Altenmuller E, Kliegel M, Van De Ville D, Marie D. Randomized controlled trials of non-

pharmacological interventions for healthy seniors: Effects on cognitive decline, brain plasticity and activities of daily living-A 23-year scoping review. *Heliyon*. 2024; 10:e26674

[35] Marie D, Muller CAH, Altenmuller E, Van De Ville D, Junemann K, Scholz DS, Kruger THC, Worschech F, Kliegel M, Sinke C, James CE. Music interventions in 132 healthy older adults enhance cerebellar grey matter and auditory working memory, despite general brain atrophy. *Neuroimage Rep*. 2023; 3:100166

[36] Mack M, Marie D, Worschech F, Kruger THC, Sinke C, Altenmuller E, James CE, Kliegel M. Effects of a 1-year piano intervention on cognitive flexibility in older adults. *Psychol Ageing*. 2025; 40:218-35

[37] Junemann K, Marie D, Worschech F, Scholz DS, Grouiller F, Kliegel M, Van De Ville D, James CE, Kruger THC, Altenmuller E, Sinke C. Six Months of Piano Training in Healthy Elderly Stabilizes White Matter Microstructure in the Fornix, Compared to an Active Control Group. *Front Ageing Neurosci*. 2022; 14:817889

[38] Hsieh S, Hornberger M, Piguet O, Hodges JR. Neural basis of music knowledge: evidence from the dementias. *Brain*. 2011; 134:2523-34

[39] Moussard A, Bigand E, Belleville S, Peretz I. Music as an aid to learn new verbal information in Alzheimer's disease. US: University of California Press; 2012. p. 521-31.

[40] El Haj M, Clément S, Fasotti L, Allain P. Effects of music on autobiographical verbal narration in Alzheimer's disease. *Journal of Neurolinguistics*. 2013; 26:691-700

- [41] Foster NA, Valentine ER. The effect of auditory stimulation on autobiographical recall in dementia. *Exp Ageing Res.* 2001; 27:215-28
- [42] Irish M, Cunningham CJ, Walsh JB, Coakley D, Lawlor BA, Robertson IH, Coen RF. Investigating the enhancing effect of music on autobiographical memory in mild Alzheimer's disease. *Dement Geriatr Cogn Disord.* 2006; 22:108-20
- [43] Schulkind MD. Is memory for music special? *Ann N Y Acad Sci.* 2009; 1169:216-24
- [44] Ren Y, Brown TI. Beyond the ears: A review exploring the interconnected brain behind the hierarchical memory of music. *Psychon Bull Rev.* 2024; 31:507-30
- [45] Finke C, Esfahani NE, Ploner CJ. Preservation of musical memory in an amnesic professional cellist. *Curr Biol.* 2012; 22:R591-2
- [46] Cuddy LL, Duffin JM, Gill SS, Brown CL, Sikka R, Vanstone AD. Memory for melodies and lyrics in Alzheimer's disease. *Music Perception.* 2012; 29:479-91
- [47] Vanstone AD, Sikka R, Tangness L, Sham R, Garcia A, Cuddy LL. Episodic and semantic memory for melodies in Alzheimer's disease. *Music Perception.* 2012; 29:501-7
- [48] Fusar-Poli L, Bieleninik Ł, Brondino N, Chen XJ, Gold C. The effect of music therapy on cognitive functions in patients with dementia: a systematic review and meta-analysis. *Ageing Ment Health.* 2018; 22:1097-106

- [49] Dorris JL, Neely S, Terhorst L, VonVille HM, Rodakowski J. Effects of music participation for mild cognitive impairment and dementia: A systematic review and meta-analysis. *J Am Geriatr Soc.* 2021; 69:2659-67
- [50] Lam HL, Li WTV, Laher I, Wong RY. Effects of Music Therapy on Patients with Dementia-A Systematic Review. *Geriatrics (Basel).* 2020; 5
- [51] van der Steen JT, van der Wouden JC, Methley AM, Smaling HJA, Vink AC, Bruinsma MS. Music-based therapeutic interventions for people with dementia. *Cochrane Database Syst Rev.* 2025; 3:Cd003477
- [52] Grau-Sanchez J, Jamey K, Paraskevopoulos E, Dalla Bella S, Gold C, Schlaug G, Belleville S, Rodriguez-Fornells A, Hackney ME, Sarkamo T. Putting music to trial: Consensus on key methodological challenges investigating music-based rehabilitation. *Ann N Y Acad Sci.* 2022; 1518:12-24
- [53] Moreno-Morales C, Calero R, Moreno-Morales P, Pintado C. Music Therapy in the Treatment of Dementia: A Systematic Review and Meta-Analysis. *Front Med (Lausanne).* 2020; 7:160
- [54] Krabs RU, Enk R, Teich N, Koelsch S. Autonomic effects of music in health and Crohn's disease: the impact of isochronicity, emotional valence, and tempo. *PLoS One.* 2015; 10:e0126224
- [55] Orini M, Bailon R, Enk R, Koelsch S, Mainardi L, Laguna P. A method for continuously assessing the autonomic response to music-induced emotions through HRV analysis. *Med Biol Eng Comput.* 2010; 48:423-33
- [56] Dalla Bella S, Foster NEV, Laflamme H, Zagala A, Melissa K, Komeilipoor N, Blais M, Rigoulot S, Kotz SA. Mobile version of the

Battery for the Assessment of Auditory Sensorimotor and Timing Abilities (BAASTA): Implementation and adult norms. *Behav Res Methods*. 2024; 56:3737-56

[57] Steele CJ, Penhune VB. Specific increases within global decreases: a functional magnetic resonance imaging investigation of five days of motor sequence learning. *J Neurosci*. 2010; 30:8332-41

[58] Zatorre RJ, Chen JL, Penhune VB. When the brain plays music: auditory-motor interactions in music perception and production. *Nat Rev Neurosci*. 2007; 8:547-58

[59] Schwartz M, Kotz SA. A dual-pathway neural architecture for specific temporal prediction. *Neurosci Biobehav Rev*. 2013; 37:2587-96

[60] Schwartz M, Tavano A, Schroger E, Kotz SA. Temporal aspects of prediction in audition: cortical and subcortical neural mechanisms. *Int J Psychophysiol*. 2012; 83:200-7

[61] Schwartz M, Rothermich K, Kotz SA. Functional dissociation of pre-SMA and SMA-proper in temporal processing. *Neuroimage*. 2012; 60:290-8

[62] Dalla Bella S. Music and movement: Towards a translational approach. *Neurophysiol Clin*. 2018; 48:377-86

[63] Benoit CE, Dalla Bella S, Farrugia N, Obrig H, Mainka S, Kotz SA. Musically cued gait-training improves both perceptual and motor timing in Parkinson's disease. *Front Hum Neurosci*. 2014; 8:494

[64] Bella SD, Benoit CE, Farrugia N, Keller PE, Obrig H, Mainka S, Kotz SA. Gait improvement via rhythmic stimulation in Parkinson's disease is linked to rhythmic skills. *Sci Rep*. 2017; 7:42005

- [65] Tolleson CM, Dobolyi DG, Roman OC, Kanoff K, Barton S, Wylie SA, Kubovy M, Claassen DO. Dysrhythmia of timed movements in Parkinson's disease and freezing of gait. *Brain Res.* 2015; 1624:222-31
- [66] Puyjarinet F, Begel V, Geny C, Driss V, Cuartero MC, De Cock VC, Pinto S, Dalla Bella S. At-Home Training With a Rhythmic Video Game for Improving Orofacial, Manual, and Gait Abilities in Parkinson's Disease: A Pilot Study. *Front Neurosci.* 2022; 16:874032
- [67] Grabli D, Karachi C, Welter ML, Lau B, Hirsch EC, Vidailhet M, Francois C. Normal and pathological gait: what we learn from Parkinson's disease. *J Neurol Neurosurg Psychiatry.* 2012; 83:979-85
- [68] Ghai S, Ghai I, Schmitz G, Effenberg AO. Effect of rhythmic auditory cueing on parkinsonian gait: A systematic review and meta-analysis. *Sci Rep.* 2018; 8:506
- [69] McIntosh GC, Brown SH, Rice RR, Thaut MH. Rhythmic auditory-motor facilitation of gait patterns in patients with Parkinson's disease. *J Neurol Neurosurg Psychiatry.* 1997; 62:22-6
- [70] Rochester L, Hetherington V, Jones D, Nieuwboer A, Willems AM, Kwakkel G, Van Wegen E. The effect of external rhythmic cues (auditory and visual) on walking during a functional task in homes of people with Parkinson's disease. *Arch Phys Med Rehabil.* 2005; 86:999-1006
- [71] Thaut MH, McIntosh GC, Rice RR, Miller RA, Rathbun J, Brault JM. Rhythmic auditory stimulation in gait training for Parkinson's disease patients. *Mov Disord.* 1996; 11:193-200
- [72] Cochen De Cock V, Dotov D, Damm L, Lacombe S, Ihalainen P, Picot MC, Galtier F, Lebrun C, Giordano A, Driss V, Geny C, Garzo A,

Hernandez E, Van Dyck E, Leman M, Villing R, Bardy BG, Dalla Bella S. BeatWalk: Personalized Music-Based Gait Rehabilitation in Parkinson's Disease. *Front Psychol.* 2021; 12:655121

[73] Nombela C, Hughes LE, Owen AM, Grahn JA. Into the groove: can rhythm influence Parkinson's disease? *Neurosci Biobehav Rev.* 2013; 37:2564-70

[74] Camurri A, Mazzarino B, Volpe G, Morasso PG, Priano F, Re C. Application of multimedia techniques in the physical rehabilitation of Parkinson's patients. *Comput Animat Virtual Worlds.* 2003; 14:269-78

[75] Barbagelata M, Morganti W, Seminerio E, Camurri A, Ghisio S, Loro M, Puleo G, Dijk B, Nolasco I, Costantini C, Cera A, Senesi B, Ferrari N, Canepa C, Custodero C, Pilotto A. Resilience improvement through a multicomponent physical and cognitive intervention for older people: the DanzArTe emotional well-being technology project. *Ageing Clin Exp Res.* 2024; 36:72

[76] Camurri A, Seminerio E, Morganti W, Canepa C, Ferrari N, Ghisio S, Cera A, Coletta P, Barbagelata M, Puleo G, Nolasco I, Costantini C, Senesi B, Pilotto A. Development and validation of an art-inspired multimodal interactive technology system for a multi-component intervention for older people: a pilot study. *Front Comput Sci.* 2023; 5:1290589

[77] Carlomagno F, Bevilacqua V, Brunetti A, Sibilano E, Delussi M, Lippolis M, Diomede R, Brattico E. Proof-of-Concept Study on the Use of Virtual Reality with Evocative and Aesthetic Content for Elderly Individuals with Cognitive Decline. *Applied Sciences.* 2025; 15:4627

- [78] Brattico E, Varankaitė U. Aesthetic empowerment through music. *Musicae Scientiae*. 2019; 23:285-303
- [79] Schafer RM. *The Tuning of the World*. New York: Knopf; 1977.
- [80] Louwers G, Gommers D, van der Heide EM, Pont S, Ozcan E. Tranquil or desolate? A mixed-methods investigation of patient sound experiences, needs and emotions in single patient ICU rooms. *Intensive Crit Care Nurs*. 2025; 89:104031
- [81] Ozcan E, van Egmond R. The effect of visual context on the identification of ambiguous environmental sounds. *Acta Psychol (Amst)*. 2009; 131:110-9
- [82] Louwers G, Pont S, Gommers D, van der Heide E, Ozcan E. Sonic ambiances through fundamental needs: An approach on soundscape interventions for intensive care patients. *J Acoust Soc Am*. 2024; 156:2376-94
- [83] Voisin F, Bidotti A, Mourey F, editors. *Designing Soundscapes for Alzheimer's Disease Care, with Preliminary Clinical Observations. Perception, Representations, Image, Sound, Music*; 2021 2021; Cham: Springer International Publishing.