



A non-local semilinear eigenvalue problem

Giovanni Franzina¹ · Danilo Licheri²

Received: 21 July 2022 / Revised: 13 October 2022 / Accepted: 13 October 2022 /
Published online: 16 November 2022
© The Author(s) 2022

Abstract

We prove that positive solutions of the *fractional Lane–Emden equation* with homogeneous Dirichlet boundary conditions satisfy pointwise estimates in terms of the best constant in Poincaré’s inequality on all open sets, and are isolated in L^1 on smooth bounded ones, whence we deduce the isolation of the first *non-local semilinear eigenvalue*.

Keywords Eigenvalues · Constrained critical points · Lane–Emden equation

Mathematics Subject Classification 35P30 · 49R05

1 Introduction

This paper concerns a semilinear problem for the fractional Laplace operator with homogeneous Dirichlet boundary conditions in N -dimensional Euclidean spaces with applications to a model for non-local filtration in a porous medium. We recall that, given $s \in (0, 1)$, the s -Laplacian of a smooth function u on \mathbb{R}^N is defined, up to a normalisation constant depending only on N and s , by the formula

$$(-\Delta)^s u(x) = \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^N \setminus B_\varepsilon(x)} \frac{u(x) - u(y)}{|x - y|^{N+2s}} dy. \quad (1.1)$$

✉ Giovanni Franzina
giovanni.franzina@cnr.it

Danilo Licheri
d.licheri2@studenti.unica.it

¹ Istituto per le Applicazioni del Calcolo “M. Picone”, Consiglio Nazionale delle Ricerche, Via dei Taurini 19, 00185 Rome, Italy

² Dipartimento di Matematica e Informatica, Università degli Studi di Cagliari, Via Ospedale 72, 09124 Cagliari, Italy

The integral in (1.1) is usually multiplied by the quantity $4^s \Gamma(\frac{N}{2} + s) / (\pi^{N/2} |\Gamma(-s)|)$, which has a precise degenerate behaviour both as $s \rightarrow 0^+$ and as $s \rightarrow 1^-$. The specific normalisation choice has no bearing for the matter of this paper and will be omitted.

By classical spectral theory in Hilbert spaces, it is known that the eigenvalue problem

$$(-\Delta)^s u = \lambda u$$

in an open set $\Omega \subset \mathbb{R}^N$, with Dirichlet conditions $u = 0$ in the complement $\mathbb{R}^N \setminus \Omega$, has non trivial solutions for a discrete set of real numbers λ , which either is empty or consists of an unbounded non-decreasing sequence of *eigenvalues*. The corresponding *eigenfunctions* are the stationary points of the double integral

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy \tag{1.2}$$

subject to an $L^2(\Omega)$ -constraint.

The variational problem under an $L^q(\Omega)$ -constraint, with $q \neq 2$, leads one to a different non-local semilinear elliptic boundary value problem, formally

$$\begin{cases} (-\Delta)^s u = \lambda \|u\|_{L^q(\Omega)}^{2-q} |u|^{q-2} u, & \text{in } \Omega, \\ u = 0, & \text{in } \mathbb{R}^N \setminus \Omega. \end{cases} \tag{1.3}$$

Any fixed solution u of (1.3), if multiplied by a specific constant depending on u , solves the *fractional Lane–Emden equation*

$$(-\Delta)^s u = |u|^{q-2} u, \text{ in } \Omega, \tag{1.4}$$

with $u = 0$ in $\mathbb{R}^N \setminus \Omega$.

The largest lower bound for the collection $\mathfrak{S}(\Omega, s, q)$ of all positive numbers λ for which (1.3) admits a non-trivial solution is called the *first q -semilinear s -eigenvalue*

$$\lambda_1(\Omega, s, q) = \inf_{\varphi \in C_0^\infty(\Omega)} \left\{ \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\varphi(x) - \varphi(y))^2}{|x - y|^{N+2s}} dx dy : \int_{\Omega} |\varphi|^q dx = 1 \right\}. \tag{1.5}$$

In some cases, for example whenever Ω has finite N -dimensional volume, the embedding of $\mathcal{D}_0^{s,2}(\Omega)$ into $L^q(\Omega)$ is compact, which assures the infimum to be achieved.

For $q \in (1, 2)$, in fact, a necessary and *sufficient* condition that the embedding be compact is that it be continuous, see [19, Theorem 1.3]. Hence, we have the following existence and uniqueness result.

Theorem A *Let $N \geq 1$, $s \in (0, 1)$, $q \in (1, 2)$, let $\Omega \subset \mathbb{R}^N$ be open, and let $\lambda_1(\Omega, s, q) > 0$. Up to a multiplicative constant, there exists a unique eigenfunction achieving the minimum in (1.5). The first eigenfunction has constant sign, and the first eigenvalue is the unique one admitting eigenfunctions with this property.*

The proof of Theorem A follows standard methods, cf. [6]. Its conclusion implies the uniqueness of *positive least energy solutions* of (1.4), i.e., positive solutions of the fractional Lane–Emden equation, under homogeneous Dirichlet boundary conditions, that minimise the energy functional

$$\frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\varphi(x) - \varphi(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{q} \int_{\Omega} |\varphi|^q dx. \tag{1.6}$$

Thus, for $q \in (1, 2)$, to every open set Ω with $\lambda_1(\Omega, s, q) > 0$ we can uniquely associate the positive least energy solution $w_{\Omega, s, q}$, also called the *fractional Lane–Emden density of Ω* . In fact, the definition can be given for arbitrary open sets in \mathbb{R}^N , see Section 5 for details. Incidentally, even if uniqueness generally fails to hold in the superhomogeneous case, one can however show that the simplicity property of Theorem A holds also for $q > 2$ smaller than a suitable threshold depending on Ω , see Proposition 3.5.

Remarkably, in analogy with the local case (cf. [8]), a negative power of the fractional Lane–Emden density of Ω appears as a singular weight in a sort of Hardy inequality:

$$\int_{\Omega} \frac{u^2}{w_{\Omega, s, q}^{2-q}} dx \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy, \quad \text{for all } u \in C_0^\infty(\Omega). \tag{1.7}$$

We refer to Proposition 5.1 for more details about (1.7). A better known Hardy-type inequality in the fractional setting would involve the distance to the boundary, instead:

$$\int_{\Omega} \frac{u(x)^2}{\text{dist}(x, \partial\Omega)^{2s}} dx \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy, \quad \text{for all } u \in C_0^\infty(\Omega). \tag{1.8}$$

Inequality (1.8) always holds, e.g., on bounded Lipschitz sets (see Section 5).

For $q \in (1, 2)$, (1.7) implies the strict stability of the fractional Lane–Emden density as a critical point of the energy (1.6). In that respect, it is noteworthy that for this special solution the following estimates hold on any (possibly unbounded) open set supporting a Poincaré inequality.

Theorem B *Let $q \in (1, 2)$ and assume that $\lambda_1(\Omega, s, 2) > 0$. Then, $w_{\Omega, s, q} \in L^\infty(\Omega)$ and there exists a constant C , depending only on N, s and q , such that*

$$\|w_{\Omega, s, q}\|_{L^\infty(\Omega)}^{2-q} \leq C \lambda_1(\Omega, s, 2)^{-1}. \tag{1.9}$$

Conversely, for all $q \in (1, 2)$, if $w_{\Omega, s, q} \in L^\infty(\Omega)$, then $\lambda_1(\Omega, s, 2) \geq \|w_{\Omega, s, q}\|_{L^\infty(\Omega)}^{q-2}$.

This statement is the non-local counterpart of [8, Proposition 4.3]. Note that the qualitative consequence of estimate (1.9) is valid irrespective of the dimension, at variance with what happens with the more difficult problem of a non-decreasing and *convex* non-linearity; in fact, for such a problem, in the local case, stable solutions are

bounded up to dimension 9, as shown in [12], whereas in higher dimensions examples of singular solutions can be made, see [12].

From inequalities (1.7) and (1.8), thanks to fractional Hopf’s lemma, we can also infer the local uniqueness in $L^1(\Omega)$ for positive solutions of fractional Lane–Emden equation (1.4); this means that the positive least energy solution $w_{\Omega,s,q}$ of (1.4) is isolated in $\mathcal{D}_0^{s,2}(\Omega)$ with respect to the topology of the convergence in $L^1(\Omega)$. We refer to Proposition 6.1 for a more precise statement. By a strategy borrowed from [5], where the result was first proved in the local case, we draw the following consequence.

Theorem C *Let $N \geq 1$, $s \in (0, 1)$, $q \in (1, 2)$ and let $\Omega \subset \mathbb{R}^N$ a bounded open set with $C^{1,1}$ boundary. Then, $\lambda_1(\Omega, s, q)$ is isolated, i.e., there exist no sequence of q -semilinear s -eigenvalues converging to it.*

Little more is known about higher eigenvalues, except that they form a closed set that does not accumulate to $\lambda_1(\Omega, s, q)$. It is indeed possible to assemble an unbounded sequence of q -semilinear s -eigenvalues by means of standard critical point theory (see Remark 3.1; see also [26, Proposition 9] for the linear case) but it is not known if that gives a complete description of the q -semilinear s -spectrum, nor is it known if the latter is a discrete set.

Put in perspective, the simplicity (Theorem A) and the isolation (Theorem C) of the first q -semilinear s -eigenvalue, paired with the global estimates (Theorem B), have implications on the long-time asymptotics for the initial-boundary value problem

$$\begin{cases} \partial_t v + (-\Delta)^s (|v|^{m-1} v) = 0, & \text{in } \Omega \times (0, T), \\ v = 0, & \text{in } (\mathbb{R}^N \setminus \Omega) \times (0, T), \\ v = v_0, & \text{in } \Omega \times \{0\}, \end{cases}$$

where $m = 1/(q - 1)$. We refer the reader to [31] for this *fractional porous media equation*. We hope to return to this topic in the future, while in this paper we limit our attention to the elliptic problem.

Plan of the paper In Section 2, after framing our problem in appropriate function spaces we introduce the fractional semilinear eigenvalue problem and the non-local Lane–Emden density. More details on the former are provided in Section 3, and various properties of the latter are discussed in Section 4. The preliminary results are used to prove (1.7) in Section 5, where (1.8) is also proved. Then, Section 6 is devoted to the isolation of positive solutions of the non-local Lane–Emden equation; eventually, all the partial results are used in Section 7 to prove Theorems A, B, and C.

2 Framework and (pseudo) differential equations

Throughout this paper, we fix an integer $N \geq 1$, a real number $s \in (0, 1)$ and an open set $\Omega \subset \mathbb{R}^N$. The square root of

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy \tag{2.1}$$

is a norm on the vector space $C_0^\infty(\Omega)$. The metric completion of this space is denoted, here and henceforth, by $\mathcal{D}_0^{s,2}(\Omega)$.

Remark 2.1 (*Analogies and differences with other spaces*) Except for the special case $s = \frac{1}{2}$, if Ω is bounded with Lipschitz boundary, then $\mathcal{D}_0^{s,2}(\Omega)$ coincides with the closure $H_0^s(\Omega)$ of $C_0^\infty(\Omega)$ in the Sobolev–Slobodeckij space $H^s(\Omega)$ of all $u \in L^2(\Omega)$ such that

$$[u]_{H^s(\Omega)}^2 := \int_\Omega \int_\Omega \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy < +\infty.$$

In fact, in that case (see [10, Appendix B]), the “censored” Sobolev norm $\|u\|_{L^2(\Omega)} + [u]_{H^s(\Omega)}$ is equivalent to that defined by

$$\|u\|_{L^2(\Omega)} + \left(\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy \right)^{\frac{1}{2}}$$

and the latter is equivalent to the norm in $\mathcal{D}_0^{s,2}(\Omega)$, because Lipschitz sets support a Poincaré-type inequality. On the contrary, if $\partial\Omega$ is not Lipschitz regular, then the existence of functions $u \in H^s(\Omega)$ for which the integral

$$\int_\Omega \int_{\mathbb{R}^N \setminus \Omega} \frac{u(x)^2}{|x - y|^{N+2s}} dx dy$$

diverges cannot be ruled out. If Ω is bounded and Lipschitz, then $\mathcal{D}_0^{s,2}(\Omega)$ coincides with the Hilbert space $X_0^s(\Omega) = \{u \in H^s(\mathbb{R}^N) : u = 0 \text{ a.e. in } \mathbb{R}^N \setminus \Omega\}$ considered in [22].

For a general open set, it is not true that all the elements of $\mathcal{D}_0^{s,2}(\Omega)$ are functions; $\mathcal{D}_0^{s,2}(\Omega)$ is not even a distribution space, in general: see, e.g., [14, 21]. A restriction that clears off this difficulty is to consider open sets Ω supporting a Sobolev-type inequality, on which $\mathcal{D}_0^{s,2}(\Omega)$ is a function space; namely, assuming that the infimum in (1.5) is a positive number.

2.1 Semilinear fractional spectrum

We denote by 2_s^* the fractional Sobolev conjugate exponent, defined by $2N/(N - 2s)$ if $2s < N$ and $+\infty$ otherwise.

Definition 2.1 (*Semilinear fractional eigenvalues*) For $q \in (1, 2_s^*)$, we consider the constrained critical points of the double integral (2.1) along the submanifold

$$\left\{ u \in \mathcal{D}_0^{s,2}(\Omega) : \int_\Omega |u|^q dx = 1 \right\}. \tag{2.2}$$

We call *q-semilinear s-eigenvalues* the corresponding constrained critical values. Their collection is denoted by $\mathfrak{S}(\Omega, s, q)$, and is said to be the *q-semilinear s-spectrum* of Ω .

Clearly, (1.5) is the largest lower bound for $\mathfrak{S}(\Omega, s, q)$, and it is its minimum whenever the variational problem (1.5) has a solution. The restriction $q < 2_s^*$ in Definition 2.1 is natural because for $q > 2_s^*$ loss of compactness occur regardless of the properties of Ω . If $0 < s < N/2$, in the borderline case $q = 2_s^*$ the infimum in (1.5) is independent of Ω , and gives the best constant in Sobolev inequality, that reads as

$$\mathcal{S}(N, s) \|v\|_{L^{2_s^*}(\Omega)}^2 \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(v(x) - v(y))^2}{|x - y|^{N+2s}} dx dy, \quad \text{for all } v \in C_0^\infty(\mathbb{R}^N). \tag{2.3}$$

By Lagrange’s multipliers rule, the *q-semilinear s-eigenvalues* are those positive real numbers λ for which the equation

$$(-\Delta)^s u = \lambda \|u\|_{L^q(\Omega)}^{2-q} |u|^{q-2} u \tag{2.4}$$

has a non-trivial solution $u \in \mathcal{D}_0^{s,2}(\Omega)$ in the weak sense, viz.

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))(\varphi(x) - \varphi(y))}{|x - y|^{N+2s}} dx dy = \lambda \|u\|_{L^q(\Omega)}^{2-q} \int_{\Omega} |u|^{q-2} u \varphi dx, \tag{2.5}$$

for all $\varphi \in \mathcal{D}_0^{s,2}(\Omega)$.

2.2 Fractional Lane–Emden equation

Given an open set $\mathcal{U} \subset \mathbb{R}^N$, we will say a *weak supersolution* (resp., *subsolution*) of the latter in \mathcal{U} any function $u \in \mathcal{D}_0^{s,2}(\mathcal{U})$ such that

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))(\varphi(x) - \varphi(y))}{|x - y|^{N+2s}} dx dy \geq \int_{\mathcal{U}} |u|^{q-2} u \varphi dx, \quad (\text{resp., } \leq) \tag{2.6}$$

for all non-negative $\varphi \in \mathcal{D}_0^{s,2}(\mathcal{U})$. A function that is both a weak supersolution and a weak subsolution in \mathcal{U} will be called a *weak solution* in \mathcal{U} . Clearly, the weak solutions of (1.4) are the critical points on $\mathcal{D}_0^{s,2}(\Omega)$ of the free energy (1.6).

Definition 2.2 (*Fractional Lane–Emden densities*) Let $q \in (1, 2)$ and assume that $\lambda_1(\Omega, s, q) > 0$. We denote by $w_{\Omega,s,q}$ the unique solution of the variational problem

$$\min_{\varphi \in \mathcal{D}_0^{s,2}(\Omega)} \left\{ \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\varphi(x) - \varphi(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{q} \int_{\Omega} \varphi^q dx : \varphi \geq 0 \text{ a.e. in } \Omega \right\}, \tag{2.7}$$

and we call it the *(s, q)–Lane–Emden density* of Ω .

Remark 2.2 By [19, Theorem 1.3], $\lambda_1(\Omega, s, q) > 0$ implies the compactness of the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^q(\Omega)$; then, any minimising sequence for (2.7) is easily seen to be bounded in $\mathcal{D}_0^{s,2}(\Omega)$, so it converges, up to relabelling, weakly in $\mathcal{D}_0^{s,2}(\Omega)$ and strongly in $L^q(\Omega)$. Also, the constraint $\varphi \geq 0$ is convex. Thus, solutions of (2.7) exist by direct methods in calculus of variations. As for their uniqueness, minimisers of the even functional (1.6) cannot change sign by Lemma 7.1, and thence constrained minimisers are non-negative minimisers of the free energy (1.6). Then, we conclude by the uniqueness of non-negative weak solutions of (1.4) (see Remark 4.1).

3 The fractional semilinear spectral problem

Next proposition provides quantitative L^∞ -bounds for q -semilinear s -eigenfunctions u corresponding to $\lambda \in \mathfrak{S}(\Omega, s, q)$ in terms of the $L^q(\Omega)$ -norm of u and of the eigenvalue λ . For this standard result, in the proof we limit ourselves to check that Moser-type iterations such as those in Appendix to [8] can be repeated in this framework, too.

Proposition 3.1 *Let $q \in (1, 2_s^*)$ and assume the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^q(\Omega)$ to be compact. Let $\lambda \in \mathfrak{S}(\Omega, s, q)$ and let $u \in \mathcal{D}_0^{s,2}(\Omega)$ be a corresponding q -semilinear s -eigenfunction. Then*

$$\|u\|_{L^\infty(\Omega)} \leq C_1(N, s, q)\lambda^{\frac{2_s^*}{2(2_s^*-q)}} \|u\|_{L^q(\Omega)}, \quad \text{if } 2_s^* < +\infty, \tag{3.1a}$$

$$\|u\|_{L^\infty(\Omega)} \leq C_2(N, s, q, |\Omega|)\lambda \|u\|_{L^q(\Omega)}, \quad \text{if } 2_s^* = +\infty. \tag{3.1b}$$

Proof With no loss of generality, we may assume that $u > 0$. Fix $\beta > 1$ and $M > 0$. By [11, Lemma A.2] with $p = 2, a = u(x), b = u(y)$ and $g(t) = (t \wedge M)^\beta$, we get

$$\begin{aligned} & \frac{2\beta}{\beta + 1} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{\left((u(x) \wedge M)^{\frac{\beta+1}{2}} - (u(y) \wedge M)^{\frac{\beta+1}{2}} \right)^2}{|x - y|^{N+2s}} dx dy \\ & \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y)) \left((u(x) \wedge M)^\beta - (u(y) \wedge M)^\beta \right)}{|x - y|^{N+2s}} dx dy. \end{aligned} \tag{3.2}$$

Here and henceforth, we are using the following notation:

$$a \wedge b := \min\{a, b\}, \quad a \vee b := \max\{a, b\}.$$

The choice $\varphi = (u \wedge M)^\beta$ in (2.5) implies that the right integral in (3.2) does not exceed

$$\lambda \|u\|_{L^q(\Omega)}^{2-q} \int_{\Omega} u^{q-1} (u \wedge M)^\beta dx.$$

Case $N > 2s$ By the compactness of the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^q(\Omega)$ and by a density argument, Sobolev inequality (2.3) holds with $v = (u \wedge M)^{\frac{\beta+1}{2}}$. Thus, the left hand side in (3.2) is at least

$$S(N, s) \frac{2\beta}{\beta + 1} \left(\int_{\Omega} (u \wedge M)^{\frac{\beta+1}{2} 2_s^*} dx \right)^{\frac{2}{2_s^*}}.$$

As $M > 0$ was arbitrary, by the material above we deduce that

$$S(N, s) \left(\int_{\Omega} u^{\frac{\beta+1}{2} 2_s^*} dx \right)^{\frac{2}{2_s^*}} \leq \lambda \|u\|_{L^q(\Omega)}^{2-q} \frac{\beta + 1}{2\beta} \int_{\Omega} u^{\beta+q-1} dx. \tag{3.3}$$

If $1 < q < 2$, by arguing as in the second part of the proof of [7, Proposition 2.5] we see that (3.3) implies (3.1). If instead $2 \leq q < 2_s^*$, then, by Hölder’s inequality we have

$$\int_{\Omega} u^{\beta+q-1} dx \leq \|u\|_{L^q(\Omega)}^{q-2} \left(\int_{\Omega} u^{\frac{\beta+1}{2} q} dx \right)^{\frac{2}{q}},$$

and thence it follows that

$$S(N, s) \left(\int_{\Omega} u^{\frac{\beta+1}{2} 2_s^*} dx \right)^{\frac{2}{2_s^*}} \leq \lambda \frac{\beta + 1}{2\beta} \left(\int_{\Omega} u^{\frac{\beta+1}{2} q} dx \right)^{\frac{2}{q}},$$

which leads one to (3.1) again, thanks to the iteration scheme in first part of the proof of [7, Proposition 2.5].

Case $N = 1$ and $\frac{1}{2} < s < 1$ In this case, the conclusion is an immediate consequence of fractional Morrey’s embedding, see [9, Corollary 2.7].

Case $N = 1$ and $s = \frac{1}{2}$ The obvious fact in this borderline case is that solutions have bounded mean oscillation. To prove they are also bounded, we first focus on exponents $q \in (1, 2]$. By the second statement in [19, Lemma 2.3] with $p = 2$, $N = 1$ and $r = 2q$,

$$C_1 \left(\int_{\Omega} \varphi^{2q} dx \right)^{\frac{2}{q}} \leq \left(\int_{\Omega} \varphi^q dx \right)^{\frac{2}{q}} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{(\varphi(x) - \varphi(y))^2}{|x - y|^2} dx dy \tag{3.4}$$

holds, in particular, with $\varphi = (u \wedge M)^{\frac{\beta+1}{2}}$, for all $M > 0$. The constant $C_1 > 0$ depends only on q and s . Then, by (3.2), arguing as done in the previous case we get

$$C_2(s, q) \left(\int_{\Omega} u^{\frac{\beta+1}{2} 2q} dx \right)^{\frac{2}{2q}} \leq \lambda \|u\|_{L^q(\Omega)}^{2-q} \frac{\beta + 1}{2\beta} \int_{\Omega} u^{\beta+q-1} dx.$$

Hence, we arrive at the desired conclusion by arguing as done after equation (13) in [7], with minor changes (just replace 2^* by $2q$.)

In order to deal with the exponents $q > 2$, we take $\sigma \in (\frac{1}{4}, \frac{1}{2})$ with $\frac{1}{2} - \sigma$ so small that the Sobolev conjugate $2_\sigma^* = 2/(1 - 2\sigma)$ exceeds $2q$ and we observe that, for all $\varphi \in \mathcal{D}_0^{s,2}(\Omega)$,

$$C_3 \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{(\varphi(x) - \varphi(y))^2}{|x - y|^{1+2\sigma}} dx dy \leq \left(\int_{\Omega} \varphi^2 dx \right)^{2(1-2\sigma)} \left(\int_{\mathbb{R}} \int_{\mathbb{R}} \frac{(\varphi(x) - \varphi(y))^2}{|x - y|^2} dx dy \right)^{4\sigma},$$

where C_3 is an absolute constant. This follows by a homogeneity argument based on the obvious remark that

$$\iint_{|y-x|<1} \frac{(\varphi(x) - \varphi(y))^2}{|x - y|^{1+2\sigma}} dx dy \leq \iint_{|y-x|<1} \frac{(\varphi(x) - \varphi(y))^2}{|x - y|^2} dx dy,$$

and

$$\iint_{|y-x|\geq 1} \frac{(\varphi(x) - \varphi(y))^2}{|x - y|^{1+2\sigma}} dx dy \leq 2 \int_{\Omega} \varphi(x)^2 \int_{|y-x|\geq 1} \frac{dy}{|x - y|^{1+2\sigma}} dx \leq \frac{2}{\sigma} \int_{\Omega} \varphi^2 dx.$$

Recalling that $2 < 2q < 2_\sigma^*$, by interpolation we also have

$$\left(\int_{\Omega} \varphi^{2q} dx \right)^{\frac{1}{2q}} \leq \left(\int_{\Omega} \varphi^2 dx \right)^{\frac{\theta}{2}} \left(\int_{\Omega} \varphi^{2_\sigma^*} dx \right)^{\frac{1-\theta}{2_\sigma^*}},$$

where $\theta \in (0, 1)$. Then, by Sobolev inequality (2.3) with σ instead of s and by Hölder’s inequality, we have again (3.4), but with a constant different from C_1 , depending only on Ω, s and q .

In conclusion, we can take $\varphi = (u \wedge M)^{\frac{\beta+1}{2}}$ and argue as done for the exponents in the range $(1, 2]$ to get the desired estimate also in the case $q > 2$. □

The following elementary proposition contains a general property of the first semilinear fractional eigenvalue.

Proposition 3.2 *Let $q \in (1, 2_s^*)$ and assume the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^q(\Omega)$ to be compact. Then, the infimum in (1.5) is a minimum. Moreover, any minimiser is either a strictly positive or a strictly negative function.*

Proof The existence of a minimiser is an immediate consequence of the direct methods in the calculus of variations. The fact that it must have constant sign follows by Lemma 7.1. Then, the last statement follows by the strong minimum principle of Proposition 7.1. □

Besides the first eigenvalue (1.5), higher eigenvalues also exist. In fact, it is straightforward to check that the squared norm (1.2) in $\mathcal{D}_0^{s,2}(\Omega)$ satisfies the Palais–Smale condition. Hence, in view of [29, Theorem 5.7], $\mathfrak{S}(\Omega, q, s)$ is an infinite set. More precisely, for all $n \in \mathbb{N}$ we denote by $\mathfrak{I}_n(\Omega, s, q)$ the collection of all subsets A of

$$\left\{ u \in \mathcal{D}_0^{s,2}(\Omega) : \int_{\Omega} |u|^q dx = 1 \right\} \tag{3.5}$$

that are symmetric and compact in $\mathcal{D}_0^{s,2}(\Omega)$ and satisfy the following property; for every $k < n$, there exist no odd and continuous mapping from A to $\mathbb{R}^k \setminus \{0\}$. We can rephrase last property saying that the *Krasnoselskii’s genus* of A is larger than or equal to n . Then, by setting

$$\lambda_n(\Omega, s, q) = \inf_{A \in \mathfrak{X}_n(\Omega, s, q)} \max_{u \in A} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy, \tag{3.6}$$

one defines an unbounded non-decreasing sequence of q -semilinear s -eigenvalues.

Remark 3.1 In general, $\mathfrak{S}(\Omega, s, q)$ is closed. Indeed, if a sequence $(\lambda_j)_{j \in \mathbb{N}} \subset \mathfrak{S}(\Omega, s, q)$ converges to a positive number λ , there is a corresponding sequence of q -semilinear s -eigenfunctions [obtained by renormalisation in $L^q(\Omega)$] which has constant $L^q(\Omega)$ -norm and converging norm in $\mathcal{D}_0^{s,2}(\Omega)$. By uniform convexity, some subsequence is converging strongly to a limit u in $L^q(\Omega)$, and this implies that u is a q -semilinear s -eigenfunction corresponding to λ .

3.1 The sub-homogeneous case

We recall two properties of $\lambda_1(\Omega, s, q)$ for $q \leq 2$.

Proposition 3.3 *Let $q \in (1, 2]$ and assume that $\lambda_1(\Omega, s, q) > 0$. If $\lambda \in \mathfrak{S}(\Omega, s, q)$ and u is a corresponding eigenfunction, then $u \geq 0$ a.e. in Ω implies $\lambda = \lambda_1(\Omega, s, q)$.*

Proof By assumption, the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^q(\Omega)$ is continuous. Then, since $q \in (1, 2]$, by Gagliardo–Nirenberg interpolation inequality (see [19, Lemma 2.3]) it is also compact. Thus, the assumptions of Proposition 3.2 are valid.

Let $v \in \mathcal{D}_0^{s,2}(\Omega)$ be a first eigenfunction, and assume that $v > 0$ a.e. in Ω . Then, let $\lambda \in \mathfrak{S}(\Omega, s, q)$, let u be a corresponding eigenfunction, and assume that $u \geq 0$ a.e. in Ω , as well. This implies $u > 0$ a.e. in Ω by the strong minimum principle of Proposition 7.1. Being free to multiply by constants, we shall also assume both u and v to have unit norm in $L^q(\Omega)$.

Fix $\varepsilon > 0$ and write $u_\varepsilon = u + \varepsilon$. For every $x, y \in \mathbb{R}^N$, by [6, Proposition 4.2] with $p = 2$,

$$(u(x) - u(y)) \left(\frac{v(x)^q}{u_\varepsilon(x)^{q-1}} - \frac{v(y)^q}{u_\varepsilon(y)^{q-1}} \right) \leq |v(x) - v(y)|^q |u(x) - u(y)|^{2-q}.$$

Multiplying by the kernel $|x - y|^{N+2s} = |x - y|^{N\frac{q}{2} + sq + N(1 - \frac{q}{2}) + s(2 - q)}$ and integrating yields

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u(x) - u(y)}{|x - y|^{N+2s}} \left(\frac{v(x)^q}{u_\varepsilon(x)^{q-1}} - \frac{v(y)^q}{u_\varepsilon(y)^{q-1}} \right) dx dy \leq \lambda_1(\Omega, s, q)^{\frac{q}{2}} \lambda^{\frac{2-q}{2}},$$

where we also used Hölder’s inequality with exponents $\frac{2}{q}$ and $\frac{2}{2-q}$, the equations satisfied by u and v , and their normalisation in $L^q(\Omega)$. Since $\varphi = v^q / u_\varepsilon^{q-1}$ is admissible

in (2.5), we have

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u(x) - u(y)}{|x - y|^{N+2s}} \left(\frac{v(x)^q}{u_\varepsilon(x)^{q-1}} - \frac{v(y)^q}{u_\varepsilon(y)^{q-1}} \right) dx dy = \lambda \int_{\Omega} u(x)^{q-1} \frac{v(x)^q}{(u(x) + \varepsilon)^{q-1}} dx.$$

Therefore, for every $\varepsilon > 0$ we end up with inequality

$$\lambda \int_{\Omega} u(x)^{q-1} \frac{v(x)^q}{(u(x) + \varepsilon)^{q-1}} dx \leq \lambda_1(\Omega, s, q)^{\frac{q}{2}} \lambda^{\frac{2-q}{2}}. \tag{3.7}$$

Since $u > 0$ a.e. in Ω , applying Fatou’s lemma and dividing λ out we arrive at

$$1 = \int_{\Omega} v(x)^q dx \leq \left(\frac{\lambda_1(\Omega, s, q)}{\lambda} \right)^{\frac{q}{2}},$$

which gives $\lambda \leq \lambda_1(\Omega, s, q)$. The definition of $\lambda_1(\Omega, s, q)$ gives the opposite inequality. □

Proposition 3.4 *Let $q \in (1, 2]$ and assume that $\lambda_1(\Omega, s, q) > 0$. Then, $\lambda_1(\Omega, s, q)$ is simple, i.e., all the corresponding eigenfunctions are mutually proportional.*

Proof Let u and v be first eigenfunctions. By Proposition 7.1, we may assume that $u, v > 0$. Nor is there any restriction in assuming both u and v to have unit norm in $L^q(\Omega)$. For all $t \in [0, 1]$, consider the function $\xi_t : \Omega \rightarrow \mathbb{R}^2$ defined by $\xi_t(x) = (t^{1/q}u(x), (1 - t)^{1/q}v(x))$. Let $\|\cdot\|_{\ell^q}$ denote the ℓ^q -norm in \mathbb{R}^2 . Then, the convexity of $\tau \mapsto |\tau|^{2/q}$ implies

$$\|\xi_t(x) - \xi_t(y)\|_{\ell^q}^2 \leq t(u(x) - u(y))^2 + (1 - t)(v(x) - v(y))^2, \quad \text{for all } x, y \in \Omega. \tag{3.8a}$$

Also, for every $t \in [0, 1]$, set $\sigma_t(x) = \|\xi_t(x)\|_{\ell^q}$ for $x \in \Omega$ and $\sigma_t(x) = 0$ for $x \in \mathbb{R}^N \setminus \Omega$. Then

$$(\sigma_t(x) - \sigma_t(y))^2 = (\|\xi_t(x)\|_{\ell^q} - \|\xi_t(y)\|_{\ell^q})^2, \quad \text{for all } x, y \in \Omega. \tag{3.8b}$$

Hence, by triangle inequality, $\sigma_t \in \mathcal{D}_0^{s,2}(\Omega)$ with the estimate

$$\begin{aligned} \iint_{\mathbb{R}^{2N}} \frac{(\sigma_t(x) - \sigma_t(y))^2}{|x - y|^{N+2s}} dx dy &\leq t \iint_{\mathbb{R}^{2N}} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy + (1 - t) \\ &\quad \iint_{\mathbb{R}^{2N}} \frac{(v(x) - v(y))^2}{|x - y|^{N+2s}} dx dy. \end{aligned}$$

The normalisation in $L^q(\Omega)$ of u and of v implies that the right hand side in the latter equals $\lambda_1(\Omega, s, q)$. On the other hand, the left hand side is larger than or equal to $\lambda_1(\Omega, s, q)$, because

$$\int_{\Omega} \sigma_t(x)^q dx = \int_{\Omega} \|\xi_t(x)\|_{\ell^q}^q dx = t \int_{\Omega} u(x)^q dx + (1 - t) \int_{\Omega} v(x)^q dx = 1.$$

Thus, σ_t is admissible for the minimisation problem that defines $\lambda_1(\Omega, s, q)$. Therefore, for every $t \in [0, 1]$, the previous integral inequality is an equality. As a consequence, the pointwise identity

$$(\sigma_t(x) - \sigma_t(y))^2 = t(u(x) - u(y))^2 + (1 - t)(v(x) - v(y))^2$$

holds for all $t \in [0, 1]$ and for a.e. $x, y \in \Omega$. In view of (3.8), the latter yields the equality case in triangle inequality

$$|\|\xi_t(x)\|_{\ell^q} - \|\xi_t(y)\|_{\ell^q}| \leq \|\xi_t(x) - \xi_t(y)\|_{\ell^q},$$

which occurs if and only if there exists $\alpha(x, y) \in \mathbb{R}$ with $\xi_t(x) = \alpha(x, y)\xi_t(y)$. Owing to the definition of ξ_t , it follows that $u(x) = \alpha(x, y)u(y)$ and $v(x) = \alpha(x, y)v(y)$. In conclusion, for a.e. x, y , we have $\frac{u(x)}{v(x)} = \frac{u(y)}{v(y)}$ and this concludes the proof. \square

3.2 The super-homogeneous case

Following the proof of [7, Proposition 4.3] about an analogous property in the local case, we show that the first eigenvalue on Ω is simple also in the super-homogeneous case $q > 2$, for all q up to a suitable threshold (depending on Ω). For this purpose, we first discuss the continuous dependence of $\lambda_1(\Omega, s, q)$ on q with a method used in [1, Lemma 4] to derive monotonicity of semilinear eigenvalues with respect to q in the local case; here we limit our attention to the right continuity at $q = 2$, which can be proved also by different methods (see [3, Lemma 2.1].)

Lemma 3.1 *We have*

$$\lim_{q \rightarrow 2^+} \lambda_1(\Omega, s, q) = \lambda_1(\Omega, s, 2).$$

Proof By [19, Corollary 1.2], we have $\lambda_1(\Omega, s, 2) > 0$ if and only if

$$\lambda_1(\Omega, s, q) > 0, \quad \text{for every } q \in [2, 2_s^*). \tag{3.9}$$

Hence, we can assume that (3.9) holds, otherwise the conclusion is obvious. Therefore,

$$\sup_{v \in C_0^\infty(\Omega)} \left\{ \int_{\Omega} |v|^q dx : \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(v(x) - v(y))^2}{|x - y|^{N+2s}} dx dy = 1 \right\} = \lambda_1(\Omega, s, q)^{-\frac{q}{2}}, \tag{3.10}$$

for all $q \in [2, 2_s^*)$, which can be seen by a straightforward homogeneity argument. Since

$$\frac{d^2}{dq^2} \int_{\Omega} |v|^q dx q = \int_{\{v \neq 0\}} |v|^q (\log |v|)^2 dx \geq 0, \quad \text{for all } q > 1,$$

the left hand side of (3.10), as a function of q , is the pointwise supremum of a family of lower semicontinuous convex functions on $(1, 2_s^*)$. Thus, $q \mapsto \lambda_1(\Omega, s, q)^{-q/2}$ is

continuous on $[2, 2_s^*)$, and hence so it is $q \mapsto \lambda_1(\Omega, s, q)$ on $[2, 2_s^*)$, by composition. \square

Proposition 3.5 *Assume that the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^2(\Omega)$ is compact. Then, there exists $q_\Omega \in (2, 2_s^*)$ such that $\lambda_1(\Omega, s, q)$ is simple for all $q \in (2, q_\Omega)$.*

Proof Let $(q_n)_{n \in \mathbb{N}}$ be a decreasing sequence converging to 2 and let $(u_n)_{n \in \mathbb{N}}$ and $(v_n)_{n \in \mathbb{N}}$ be sequences in $\mathcal{D}_0^{s,2}(\Omega)$ such that, for all $n \in \mathbb{N}$, equation (2.5) holds with $\lambda = \lambda_1(\Omega, s, q_n)$ both for $u = u_n$ and for $u = v_n$. By Proposition 3.2, we may assume u_n and v_n to be positive functions, nor does it cause any loss of generality assuming them to have unit $L^{q_n}(\Omega)$ -norm. Then, by using themselves as test functions in their own equations, in view of Lemma 3.1 we see that

$$\begin{aligned} \lim_{n \rightarrow \infty} \iint_{\mathbb{R}^{2N}} \frac{(u_n(x) - u_n(y))^2}{|x - y|^{N+2s}} dx dy &= \lim_{n \rightarrow \infty} \iint_{\mathbb{R}^{2N}} \frac{(v_n(x) - v_n(y))^2}{|x - y|^{N+2s}} dx dy \\ &= \lambda_1(\Omega, s, 2). \end{aligned}$$

Also, because, by assumption, the infimum that defines $\lambda_1(\Omega, s, 2)$ is achieved, we have

$$\lambda_1(\Omega, s, 2) = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\bar{u}(x) - \bar{u}(y))^2}{|x - y|^{N+2s}} dx dy$$

for an appropriate function $\bar{u} \in \mathcal{D}_0^{s,2}(\Omega)$ with unit norm in $L^2(\Omega)$.

By Proposition 3.4, \bar{u} is uniquely determined; hence, from the assumption that the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^2(\Omega)$ is compact, we infer that both $(u_n)_{n \in \mathbb{N}}$ and $(v_n)_{n \in \mathbb{N}}$ converge to \bar{u} strongly in $\mathcal{D}_0^{s,2}(\Omega)$ and pointwise a.e. in Ω , by using the last two identities in display and the fact that, for any given $\gamma > 2$, owing to Proposition 3.1 we have

$$\|u_n - \bar{u}\|_{L^\gamma(\Omega)} \leq c \|u_n - \bar{u}\|_{L^{\frac{2}{\gamma}}(\Omega)}^{\frac{2}{\gamma}}, \quad \|v_n - \bar{u}\|_{L^\gamma(\Omega)} \leq c \|v_n - \bar{u}\|_{L^{\frac{2}{\gamma}}(\Omega)}^{\frac{2}{\gamma}}$$

for a constant $c > 0$ independent of n .

As $q_n > 2$, by Proposition 3.1 there exists $C > 0$, depending only on the data, with

$$w_n := (q_n - 1) \int_0^1 [tu_n + (1 - t)v_n]^{q_n - 2} dt \leq C. \tag{3.11}$$

The latter appears as a weight in the equation for $\psi_n = \|u_n - v_n\|_{L^2(\Omega)}^{-1} (u_n - v_n)$, viz.

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\psi_n(x) - \psi_n(y))(\varphi(x) - \varphi(y))}{|x - y|^{N+2s}} dx dy = \lambda_1(\Omega, s, q_n) \int_{\Omega} w_n \psi_n \varphi dx, \tag{3.12}$$

for all $\varphi \in \mathcal{D}_0^{s,2}(\Omega)$. After choosing $\varphi = \psi_n$ in (3.12), in view of (3.11) we see that $(\psi_n)_{n \in \mathbb{N}}$ is bounded in $\mathcal{D}_0^{s,2}(\Omega)$. Thus, by assumption, a subsequence (not relabelled)

converges to some limit ψ , weakly in $\mathcal{D}_0^{s,2}(\Omega)$ and strongly in $L^2(\Omega)$. Then, ψ is bound to have unit norm in $L^2(\Omega)$, in particular $\psi \neq 0$.

We claim that

$$w_n \rightarrow 1, \quad \text{in } L^2_{\text{loc}}(\Omega). \tag{3.13}$$

Thence, recalling also Lemma 3.1, by passing to the limit in (3.12) we arrive at

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\psi(x) - \psi(y))(\varphi(x) - \varphi(y))}{|x - y|^{N+2s}} dx dy = \lambda_1(\Omega, s, 2) \int_{\Omega} \psi \varphi dx, \tag{3.14}$$

for all $\varphi \in \mathcal{D}_0^{s,2}(\Omega)$, i.e., ψ is a non-trivial first eigenfunction. By Proposition 3.4, it follows that either $\psi = \bar{u}$ or $\psi = -\bar{u}$. On the other hand, we can plug in $\varphi = \psi_n^\pm$ into (3.12) and deduce from (3.11), for n large enough, that

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\psi_n^\pm(x) - \psi_n^\pm(y))^2}{|x - y|^{N+2s}} dx dy \leq 2C\lambda_1(\Omega, s, 2) \int_{\Omega} |\psi_n^\pm|^2 dx.$$

We argue by contradiction and we assume that $u_n \neq v_n$, for all $n \in \mathbb{N}$. Hence, both $\Omega_n^+ = \{u_n > v_n\}$ and $\Omega_n^- = \{u_n < v_n\}$ must have non-zero measure, because u_n and v_n have the same $L^q(\Omega)$ -norm. Then, we can estimate from below the left hand side to get

$$|\Omega_n^\pm|^{\frac{2s}{N}} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\psi_n^\pm(x) - \psi_n^\pm(y))^2}{|x - y|^{N+2s}} dx dy \geq C' \int_{\Omega} |\psi_n^\pm|^2 dx$$

where C' depends only on N and s ; indeed, if Ω_n^\pm has infinite measure, then the latter is trivial; otherwise, we can deduce it from the definition of $\lambda_1(\Omega, s, 2)$, its scaling properties and the fractional Faber–Krahn inequality [10, Theorem 3.5]. Combining the upper and the lower bound yields $\inf_{n \in \mathbb{N}} |\Omega_n^\pm| > 0$, which is inconsistent with the pointwise convergence of ψ_n to its constant sign limit ψ . Indeed, by Egorov’s Theorem, the pointwise a.e. convergence implies its uniform convergence out of an exceptional set of arbitrarily small measure; since the limit function is either positive or negative, it follows that $|\Omega_n^+| \wedge |\Omega_n^-|$ converges to zero as $n \rightarrow \infty$.

Thus, we are left with proving the claim (3.13). To do so, we consider a bounded open set $\Omega' \Subset \Omega$ and observe that

$$\begin{aligned} \int_{\Omega'} (w_n - 1)^2 dx &\leq \int_{\Omega'} \int_0^1 \left[(q_n - 1)[tu_n + (1 - t)v_n]^{q_n - 2} - 1 \right]^2 dt dx \\ &\leq 2 \int_{\Omega'} \int_0^1 \left[(q_n - 2)^2 \left([tu_n + (1 - t)v_n]^{q_n - 2} \right)^2 \right. \\ &\quad \left. + \left([tu_n + (1 - t)v_n]^{q_n - 2} - 1 \right)^2 \right] dt dx \\ &\leq 2C^2 |\Omega'| + 2 \int_{\Omega'} \int_0^1 \left(|tu_n + (1 - t)v_n|^{q_n - 2} - 1 \right)^2 dt dx. \end{aligned}$$

By the pointwise convergence a.e. in Ω of both u_n and v_n to \bar{u} and by (3.11), the latter implies that $w_n \rightarrow 1$ in $L^2(\Omega')$ by dominated convergence theorem. Since Ω' was arbitrary, that entails (3.13), as desired. \square

4 Fractional Lane–Emden densities

In this section we always limit our attention to exponents $q \in (1, 2)$ and we prove some properties of the fractional Lane–Emden density of Ω . We recall that in this paper the function $w_{\Omega,s,q}$ is introduced in Definition 2.2, under the assumption that $\lambda_1(\Omega, s, q) > 0$, as a non-negative weak solution of (1.4) (see also Remark 2.2).

Remark 4.1 Equation (1.4) has indeed a unique non-negative weak solution; by Proposition 3.3, any such function is a non-negative q -semilinear s -eigenfunction with $L^q(\Omega)$ -norm equal to $\lambda_1(\Omega, s, q)^{\frac{1}{q-2}}$, whence the uniqueness by Proposition 3.4.

Proposition 4.1 *Let $q \in (1, 2)$, let Ω_1 and Ω_2 be bounded open sets and, for $i \in \{1, 2\}$, let w_i be the fractional Lane–Emden density $w_{\Omega_i,s,q}$ on Ω_i . Then*

$$\Omega_1 \subset \Omega_2 \implies w_1 \leq w_2.$$

Proof Let us write $w_i = w_{\Omega_i,s,q}$ in Ω_i and $w_i = 0$ in $\mathbb{R}^N \setminus \Omega_i$, for $i \in \{1, 2\}$. The inequality

$$(a \vee b - c \vee d)^2 - (a - c)^2 \leq (b - d)^2 - (a \wedge b - c \wedge d)^2,$$

with $a = w_1(x)$, $b = w_2(x)$, $c = w_1(y)$ and $d = w_2(y)$ entails the submodularity property

$$\begin{aligned} & \frac{1}{2} \iint_{\mathbb{R}^{2N}} \frac{((w_1 \vee w_2)(x) - (w_1 \vee w_2)(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{2} \iint_{\mathbb{R}^{2N}} \frac{(w_1(x) - w_1(y))^2}{|x - y|^{N+2s}} dx dy \\ & \leq \frac{1}{2} \iint_{\mathbb{R}^{2N}} \frac{(w_2(x) - w_2(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{2} \iint_{\mathbb{R}^{2N}} \frac{((w_1 \wedge w_2)(x) - (w_1 \wedge w_2)(y))^2}{|x - y|^{N+2s}} dx dy. \end{aligned}$$

By minimality of w_1 , we also have

$$\begin{aligned} & \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(w_1(x) - w_1(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{q} \int_{\Omega_1} w_1^q dx \\ & \leq \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{((w_1 \wedge w_2)(x) - (w_1 \wedge w_2)(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{q} \int_{\Omega_1} (w_1 \wedge w_2)^q dx. \end{aligned}$$

Then, taking into account the integral identity

$$\frac{1}{q} \int_{\Omega_1} w_1^q dx - \frac{1}{q} \int_{\Omega_2} (w_1 \vee w_2)^q dx = \frac{1}{q} \int_{\Omega_1} (w_1 \wedge w_2)^q dx - \frac{1}{q} \int_{\Omega_2} w_2^q dx$$

and summing up, gives

$$\begin{aligned} & \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{((w_1 \vee w_2)(x) - (w_1 \vee w_2)(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{q} \int_{\Omega_2} (w_1 \vee w_2)^q dx \\ & \leq \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(w_2(x) - w_2(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{q} \int_{\Omega_2} w_2^q dx. \end{aligned}$$

Hence, by the minimality property of w_2 , we infer that $w_2 = w_1 \vee w_2$, as desired. \square

We can extend Definition 2.2 to the case $\lambda_1(\Omega, s, q) = 0$, as done in the local case (see [8].)

Definition 4.1 Let $q \in (1, 2)$. Then, we set

$$w_{\Omega,s,q}(x) = \lim_{r \rightarrow +\infty} w_{\Omega \cap B_r,s,q}(x), \quad \text{for all } x \in \Omega. \tag{4.1}$$

and we continue to call $w_{\Omega,s,q}$ the (s, q) -Lane–Emden density of Ω .

By Proposition 4.1, the limit (4.1) always exists, so that the definition is well posed. The following lemma assures its consistency with Definition 2.2.

Lemma 4.1 Let $q \in (1, 2)$ and assume that $\lambda_1(\Omega, s, q) > 0$. For every $r > 0$, we set $w_r(x) = w_{\Omega \cap B_r,s,q}(x)$ if $x \in B_r$ and $w_r(x) = 0$ otherwise. Then, w_r converge pointwise to $w_{\Omega,s,q}$ as $r \rightarrow +\infty$.

Proof As $r \rightarrow +\infty$, the (s, q) -Lane–Emden density w_r on $\Omega \cap B_r$ converges to an appropriate function $\bar{w} \leq w_{\Omega,s,q}$. By minimality, for every given $\varphi \in C_0^\infty(\Omega)$ there exists $R_\varphi > 0$ such that, for all $r \geq R_\varphi$, we have

$$\begin{aligned} & \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(w_r(x) - w_r(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{q} \int_{\Omega \cap B_r} w_r^q dx \\ & \leq \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\varphi(x) - \varphi(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{q} \int_{\Omega \cap B_r} |\varphi|^q dx. \end{aligned} \tag{4.2}$$

Note that the equation for w_r is (2.5) with $\Omega \cap B_r$ in place of Ω , $u = w_r$ and $\lambda = \|w_r\|_{L^q(\Omega \cap B_r)}^{q-2}$. Testing with $\varphi = w_r$ the equation for w_r , we get

$$\begin{aligned} & \iint_{\mathbb{R}^{2N}} \frac{(w_r(x) - w_r(y))^2}{|x - y|^{N+2s}} dx dy \\ & = \int_{\Omega \cap B_r} w_r^q dx \leq \lambda_1(\Omega, s, q)^{\frac{q}{2}} \left[\iint_{\mathbb{R}^{2N}} \frac{(w_r(x) - w_r(y))^2}{|x - y|^{N+2s}} dx dy \right]^{\frac{q}{2}}, \end{aligned}$$

where in the second inequality we also used that $w_r = 0$ in $\Omega \setminus B_r$. Since $q < 2$, we deduce that w_r converges to \bar{w} weakly in $\mathcal{D}_0^{s,2}(\Omega)$ and strongly in $L^q(\Omega)$. Thus,

passing to the limit as $r \rightarrow \infty$ in (4.2), we obtain

$$\begin{aligned} \frac{1}{2} \iint_{\mathbb{R}^{2N}} \frac{(\bar{w}(x) - \bar{w}(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{q} \int_{\Omega} \bar{w}^q dx &\leq \frac{1}{2} \iint_{\mathbb{R}^{2N}} \frac{(\varphi(x) - \varphi(y))^2}{|x - y|^{N+2s}} dx dy \\ &\quad - \frac{1}{q} \int_{\Omega} |\varphi|^q dx, \end{aligned}$$

for all $\varphi \in C_0^\infty(\Omega)$, which by uniqueness implies that $\bar{w} = w_{\Omega,s,q}$. □

Following [15], for all $w \in \mathcal{D}_0^{s,2}(\mathbb{R}^N)$ and for all $x_0 \in \mathbb{R}^N$, we set

$$\text{Tail}(w, x_0, \rho) = \rho^{2s} \int_{\mathbb{R}^N \setminus B_\rho(x_0)} \frac{|w(x)|}{|x - x_0|^{N+2s}} dx.$$

The only difference between next proposition and [15, Theorem 1.1] is in that we consider an equation with a non-zero right hand-side; some obvious changes in the proof of [15] give the estimate for the non-homogeneous equation. A similar estimate holds for non-homogeneous equations with data in L^γ , $\gamma > N/s$, but that is not relevant to our case.

Proposition 4.2 *Let $\mathcal{U} \subset \mathbb{R}^N$ be an open set, $x_0 \in \mathcal{U}$, $\delta \in (0, 1]$, $0 < r < \text{dist}(x, \partial\mathcal{U})$, $f \in L^\infty(\mathcal{U})$ and let $w \in \mathcal{D}_0^{s,2}(\mathbb{R}^N)$ be a non-negative weak sub-solution of $(-\Delta)^s w = f$ in \mathcal{U} , i.e., $w \geq 0$ in \mathcal{U} and*

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(w(x) - w(y))(\varphi(x) - \varphi(y))}{|x - y|^{N+2s}} dx dy \leq \int_{\mathcal{U}} f w \varphi dx,$$

for all non-negative $\varphi \in C_0^\infty(\mathcal{U})$. Then

$$\text{ess sup}_{B_{r/2}(x_0)} w \leq C \left[\delta \text{Tail}(w, x_0, r/2) + \delta r^{2s} \|f\|_{L^\infty(\mathcal{U})} + \left(\frac{r^{N-2s}}{\delta} \right)^{\frac{N}{4s}} \left(\int_{B_r(x_0)} w^2 dx \right)^{\frac{1}{2}} \right],$$

where the constant depends only on N and s .

5 Functional inequalities with special singular weights

In the present section, we introduce a couple of Hardy-type inequalities. In the following proposition, we see that Lane–Emden inequalities (see [8, Section 3]) are valid also in the non-local case.

Proposition 5.1 *Let $q \in (1, 2)$ and $u \in C_0^\infty(\Omega)$. Then*

$$\int_{\Omega} \frac{u^2}{w_{\Omega,s,q}^{\frac{2-q}{q}}} dx \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy, \tag{5.1}$$

with the agreement that the left integrand be 0 at all points where $w_{\Omega,s,q} = +\infty$.

Proof We first prove (5.1) in the special case of a bounded open set. We write $w = w_{\Omega,s,q}$, and we take $\varepsilon > 0$. By Proposition 3.1, $w \in \mathcal{D}_0^{s,2}(\Omega) \cap L^\infty(\Omega)$. Hence, so does $(w + \varepsilon)^{-1}$, because $t \mapsto (t + \varepsilon)^{-1}$ is a Lipschitz function on $(0, \infty)$. Then, by [4, Lemma 2.4] we can plug $\varphi = u^2/(w + \varepsilon)$ into the equation for w and get

$$\int_{\Omega} w(x)^{q-1} \frac{u(x)^2}{w(x) + \varepsilon} dx = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{w(x) - w(y)}{|x - y|^{N+2s}} \left(\frac{u(x)^2}{w(x) + \varepsilon} - \frac{u(y)^2}{w(y) + \varepsilon} \right) dx dy$$

for all $\varepsilon > 0$. In view of [6, Proposition 4.2], and recalling that $w > 0$ a.e. in Ω , by Fatou’s lemma it follows that

$$\int_{\Omega} \frac{u^2}{w^{2-q}} dx \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy.$$

For the general case, we take $R > 0$ so large that the support of u is contained in B_r for all $r \geq R$. For all such radii r , by the material above we have

$$\int_{\Omega \cap B_r} \frac{u^2}{w_r^{2-q}} dx \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy,$$

where w_r is the (s, q) -Lane–Emden density of $\Omega \cap B_r$. In view of Definition 4.1, by Fatou’s lemma we get the conclusion passing to the limit as $r \rightarrow \infty$. \square

The more familiar Hardy-type inequality of next proposition implies some restriction on the domain. The assumption made below is not optimal, though; for instance, a uniform exterior cone condition is also a valid assumption; more generally, for the statement to hold true it would be sufficient that no boundary point belong to the measure-theoretic interior of Ω , see [13]. The boundedness assumption is not essential, either: see [16, Theorem 1.1], or also [28, Theorem 1.10].

Proposition 5.2 *Let $s \in (0, 1)$ and let $\Omega \subset \mathbb{R}^N$ be an open bounded Lipschitz set. Then, for all $u \in C_0^\infty(\Omega)$,*

$$\int_{\Omega} \frac{u(x)^2}{\text{dist}(x, \partial\Omega)^{2s}} dx \leq C \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy \tag{5.2}$$

for a constant $C > 0$ depending only on Ω .

Before proving Proposition 5.2, we make a brief comment on (5.2). When it comes to fractional Hardy inequalities, there are a number of variants of the same statement. A stronger one just involves the Sobolev–Slobodeckij seminorm $[u]_{H^s(\Omega)}$ in the right hand side (instead of taking integrals on the whole of \mathbb{R}^N), implying various restrictions both on Ω and on s : for a more detailed account on the topic, we refer to [13, 16–18, 28]. Here, incidentally, in view of Remark 2.1 we may point out the following.

Corollary 5.1 *If $s \in (0, 1)$ and $2s \neq N$, then, under the assumptions of Proposition 5.2,*

$$\int_{\Omega} \frac{u(x)^2}{\text{dist}(x, \partial\Omega)^{2s}} dx \leq C \left(\int_{\Omega} \int_{\Omega} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy + \int_{\Omega} u^2 dx \right)$$

for all $u \in C_0^\infty(\Omega)$.

Proof of Proposition 5.2 By assumption, Ω is a bounded Lipschitz set. Hence, it satisfies the uniform exterior cone condition, *i.e.*, that there exists $\ell > 0$ and a cone K , with given aperture, such that every boundary point ξ is the vertex of a cone K_ξ isometric to K that satisfies $K_\xi \cap B_\ell(\xi) \subset \mathbb{R}^N \setminus \Omega$. For, see *e.g.* [20, Theorem 1.2.2.2].

For ease of notation, we write $\delta(x) := \text{dist}(x, \partial\Omega)$. For all $x \in \Omega$ with $\delta(x) \geq \ell$, we can pick $\xi_x \in \partial\Omega$ with minimum distance to x and we have $|x - y| \leq |\xi_x - y| + \delta(x) \leq 2\delta(x)$ for all $y \in K_{\xi_x} \cap B_\ell(\xi_x)$, whence it follows that

$$\int_{K_{\xi_x} \cap B_\ell(\xi_x)} \frac{dy}{|x - y|^{N+2s}} \geq (2\delta(x))^{-(N+2s)} |K_{\xi_x} \cap B_\ell(\xi_x)| \geq \frac{\theta \ell^N \delta(x)^{-2s}}{2^{N+2s} D^N N},$$

where $\theta = \mathcal{H}^{N-1}(K \cap \partial B_1(0))$ and D is the diameter of Ω .

The inequality $|x - y| \leq |\xi_x - y| + \delta(x)$ holds also for all points $x \in \Omega$ with $\delta(x) \leq \ell$, and we infer that

$$\begin{aligned} \int_{K_{\xi_x} \cap B_\ell(\xi_x)} \frac{dy}{|x - y|^{N+2s}} &\geq \int_0^{\delta(x)} \frac{\theta \rho^{N-1} d\rho}{(\rho + \delta)^{N+2s}} \\ &= \frac{\theta}{N\delta(x)^{2s}} \int_0^1 \frac{dt}{(1 + t^{1/N})^{N+2s}} \geq \frac{\theta \delta(x)^{-2s}}{2^{N+2s} N}. \end{aligned}$$

Since for all $x \in \Omega$ we have $K_{\xi_x} \cap B_\ell(\xi_x) \subset \mathbb{R}^N \setminus \Omega$, it follows that

$$\int_{\mathbb{R}^N \setminus \Omega} \frac{dy}{|x - y|^{N+2s}} \geq \frac{\theta \delta(x)^{-2s}}{2^{N+2s} N} \left(\frac{\ell^N}{D^N} \wedge 1 \right).$$

That gives the desired conclusion, because we have

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} = \int_{\Omega} \int_{\Omega} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} + 2 \int_{\Omega} u(x)^2 \int_{\mathbb{R}^N \setminus \Omega} \frac{dy}{|x - y|^{N+2s}},$$

for all $u \in C_0^\infty(\Omega)$. □

Remark 5.1 For the use we shall make of Proposition 5.2, we do not need to pay much attention to the explicit value of the constant $C > 0$. For sure, the proof presented implies a very rough estimate of the optimal (unknown) constant.

Remark 5.2 By density, the inequality of Corollary 5.1 holds for all functions that belong to $\mathcal{D}_0^{s,2}(\Omega)$. Given $p \in (1, \infty)$, a similar inequality, with suitable adjustments to the exponents, is valid for functions in the homogeneous fractional Sobolev space $\mathcal{D}_0^{s,p}(\Omega)$ defined as the completion of $C_0^\infty(\Omega)$ with respect to

$$\left(\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{\frac{1}{p}}$$

and this can be seen by minor changes in the proof presented here. This variant was considered, for example, in [4], where the authors provide a constant that works on convex open sets, with stable asymptotic behaviour as $s \nearrow 1$.

6 Local in L^1 uniqueness for fractional Lane–Emden positive solutions

The following proposition is the non-local counterpart of [5, Proposition 4.1].

Proposition 6.1 *Let $q \in (1, 2)$ and assume that the weighted space*

$$L^2(\Omega, w_{\Omega,s,q}^{q-2}) = \left\{ u \in L^1_{\text{loc}}(\Omega) : \int_{\Omega} w_{\Omega,s,q}^{q-2} u^2 dx < +\infty \right\} \tag{6.1}$$

contains $\mathcal{D}_0^{s,2}(\Omega)$ with compact embedding. Then, every critical point of

$$\frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy - \frac{1}{q} \int_{\Omega} |u|^q dx$$

must satisfy $\|u - w_{\Omega,s,q}\|_{L^1(\Omega)} \geq \delta$, where $\delta > 0$ depends only on s, q, Ω and N .

Proof We will prove a contrapositive statement: if a sequence $(u_n)_{n \in \mathbb{N}}$, consisting of weak solutions of the fractional Lane–Emden equation (1.4), converges to $w := w_{\Omega,s,q}$ in $L^1(\Omega)$, then

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\psi(x) - \psi(y))^2}{|x - y|^{N+2s}} dx dy \leq (q - 1) \int_{\Omega} w^{q-2} \psi^2 dx, \tag{6.2}$$

for some $\psi \in \mathcal{D}_0^{s,2}(\Omega) \setminus \{0\}$. Note that (6.2) is in contradiction with Proposition 5.1, because $q < 2$.

By setting $Q_n = (w^{q-1} - |u_n|^{q-2}u_n) / (w - u_n)$ at all points where $w \neq u_n$ and $Q_n = 0$ elsewhere, the weak equation for the difference $w - u_n$ takes the form

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{((w - u_n)(x) - (w - u_n)(y))(\varphi(x) - \varphi(y))}{|x - y|^{N+2s}} dx dy = \int_{\Omega} Q_n (w - u_n) \varphi dx,$$

for $\varphi \in \mathcal{D}_0^{s,2}(\Omega)$; the choice $\varphi = t_n^{-1}(w - u_n)$, where $t_n = \int_{\Omega} w^{q-2}(w - u_n)^2 dx$, gives

$$\frac{1}{t_n} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{((w - u_n)(x) - (w - u_n)(y))^2}{|x - y|^{N+2s}} dx dy = \int_{\Omega} Q_n \left(\frac{w - u_n}{\sqrt{t_n}} \right)^2 dx. \tag{6.3}$$

By [5, Lemma A.1], we have the following pointwise bound

$$0 \leq Q_n(x) \leq 2^{2-q} w^{q-2}(x), \quad \text{for all } x \in \Omega, \tag{6.4}$$

and, by construction, that prevents the right integral in (6.3) from exceeding the constant 2^{2-q} . Therefore, setting $\psi_n = (w - u_n)/\sqrt{t_n}$ defines a bounded sequence in $\mathcal{D}_0^{s,2}(\Omega)$, which clearly has unit norm in the weighted space (6.1).

By assumption, we deduce that ψ_n converges weakly in $\mathcal{D}_0^{s,2}(\Omega)$ and strongly in the weighted space (6.1) to a non-zero limit ψ . Thus, by (6.3), we can write

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\psi_n(x) - \psi_n(y))^2}{|x - y|^{N+2s}} dx dy = \int_{\Omega} Q_n (\psi_n^2 - \psi^2) dx + \int_{\Omega} Q_n \psi^2 dx. \tag{6.5}$$

The convergence of the sequence ψ_n implies

$$\limsup_{n \rightarrow \infty} \int_{\Omega} Q_n (\psi_n^2 - \psi^2) dx \leq 0 \tag{6.6}$$

because, by the pointwise bound (6.4) and by Hölder’s inequality, we have

$$\begin{aligned} \int_{\Omega} Q_n (\psi_n^2 - \psi^2) dx &\leq 2^{2-q} \left(\int_{\Omega} w^{q-2} (\psi_n - \psi)^2 dx \right)^{\frac{1}{2}} \times \\ &\quad \times \left[\left(\int_{\Omega} w^{q-2} \psi_n^2 dx \right)^{\frac{1}{2}} + \left(\int_{\Omega} w^{q-2} \psi^2 dx \right)^{\frac{1}{2}} \right]. \end{aligned}$$

In order to deal with the second integral in the right hand side of (6.5), we would better handle the pointwise limit behaviour of Q_n . Since

$$w^{q-1} - |u_n|^{q-2} u_n = - \int_0^1 \frac{d}{dt} \left[|w + t(u_n - w)|^{q-2} (w + t(u_n - w)) \right] t dt$$

for every $x \in \Omega$, we have

$$Q_n(x) \leq (q - 1) \int_0^1 f_n(x, t) dt, \quad \text{where } f_n(x, t) = |(1 - t)w(x) + tu_n(x)|^{q-2}. \tag{6.7}$$

The Hölder continuity of $\tau \mapsto \tau^{2-q}$ and the convexity of $\tau \mapsto \tau^{q-2}$ imply

$$||a + t(b - a)|^{q-2} - a^{q-2}| \leq \frac{t^{2-q}}{a^{2-q}} (b - a)^{2-q} \left[(1 - t)a^{q-2} + tb^{q-2} \right]$$

for all $t \in [0, 1]$ and for all $a, b > 0$. Then, at all points x where $u_n(x) > 0$, we have

$$\sup_{t \in [0,1]} |f_n(x, t) - w^{2-q}(x)| \leq \frac{|w(x) - u_n(x)|^{2-q}}{w(x)^{2-q}} \left[(1-t)w(x)^{q-2} + tu_n(x)^{q-2} \right]. \tag{6.8}$$

As u_n converges to w in $L^1(\Omega)$, a subsequence (not relabelled) also converges point-wise a.e. in Ω . In view of (6.8), that assures the uniform convergence of $f_n(x, \cdot)$ to the constant $w(x)^{2-q}$ for all x out of a negligible set, so that

$$\lim_{n \rightarrow \infty} \int_0^1 f_n(x, t) dt = w(x)^{2-q}, \quad \text{for a.e. } x \in \Omega. \tag{6.9}$$

From (6.7) and (6.9) we infer that

$$\limsup_{n \rightarrow \infty} Q_n \psi^2 \leq (q - 1)w^{2-q} \psi^2, \quad \text{a.e. in } \Omega.$$

Hence and from (6.4), by Fatou’s Lemma for the sequence $2^{2-q}w^{q-2} - Q_n \psi^2$, we arrive at

$$\limsup_{n \rightarrow \infty} \int_{\Omega} Q_n \psi^2 dx \leq (q - 1) \int_{\Omega} w^{2-q} \psi^2 dx. \tag{6.10}$$

Inserting (6.6) and (6.10) in the identity (6.5) and using the lower semicontinuity of the left hand side of (6.5) with respect to the weak convergence in $\mathcal{D}_0^{s,2}(\Omega)$, we arrive at (6.2), as desired. \square

Remark 6.1 In view of Proposition 5.1, the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^2(\Omega, w_{\Omega,s,q}^{q-2})$ is continuous, for example, on all open sets with finite volume. The stronger requirement that it be compact may be met under higher regularity assumptions on $\partial\Omega$.

Lemma 6.1 *Let $q \in (1, 2)$, let $\Omega \subset \mathbb{R}^N$ be a bounded open set with $C^{1,1}$ boundary and let $v \in C_0^\infty(\Omega)$. Then*

$$\int_{\Omega} w_{\Omega,s,q}^{q-2} v^2 dx \leq \left(\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(v(x) - v(y))^2}{|x - y|^{N+2s}} dx dy \right)^{\frac{2-q}{2}} \|v\|_{L^2(\Omega)}^q. \tag{6.11}$$

Proof By Hopf’s lemma for the fractional Laplacian (see [24, Lemma 7.3]) we have a constant $C > 0$, only depending on Ω, N, q and s , such that

$$w_{\Omega,s,q}(x) \geq C \text{dist}(x, \partial\Omega)^s. \tag{6.12}$$

Incidentally, we mention that the more precise asymptotic boundary behaviour

$$w_{\Omega,s,q} \asymp \text{dist}(\cdot, \partial\Omega)^s$$

is known: for the semilinear equation we refer to Theorem 6.4 and the following remarks in [2] (alternatively, see [25] for the linear equation with a bounded right hand

side, which is also relevant to our case thanks to Proposition 3.1). Since $q \in (1, 2)$, by Hölder’s inequality with exponents $\frac{2}{2-q}$ and $\frac{2}{q}$ we have

$$\int_{\Omega} \text{dist}(x, \partial\Omega)^{s(q-2)} v^2 dx \leq \left(\int_{\Omega} \frac{v^2}{\text{dist}(x, \partial\Omega)^{2s}} dx \right)^{\frac{2-q}{2}} \left(\int_{\Omega} v^2 dx \right)^{\frac{q}{2}}. \tag{6.13}$$

Then, by (6.12), Proposition 5.2 and (6.13), we improve the fractional Lane–Emden inequality (5.1) to (6.11). □

The conclusion of the previous lemma assures compactness for the weighted embedding. Thus, we end this section with the remark that the isolation of fractional Lane–Emden densities holds, for example, on open sets with smooth boundary.

Proposition 6.2 *Let $q \in (1, 2)$ and let $\Omega \subset \mathbb{R}^N$ be a bounded open set with $C^{1,1}$ boundary. Then, the conclusion of Proposition 6.1 holds.*

Proof By assumption, $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^2(\Omega)$ is compact; this and Lemma 6.1 imply the compactness of the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^2(\Omega, w_{\Omega,s,q}^{q-2})$, too. □

7 Proof of the main results

7.1 Proof of Theorem A

Because $q \in (1, 2)$, the assumption $\lambda_1(\Omega, s, q) > 0$ implies the compactness of the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^q(\Omega)$ (see [19, Theorem 1.3]). Then, a first eigenfunction exists by Proposition 3.2. Also, Proposition 3.4 entails uniqueness up to proportionality, and the last statement is true by Proposition 3.3. □

7.2 Proof of Theorem B

Let us write $w = w_{\Omega,s,q}$. The last statement is a consequence of Proposition 5.1. Then, we assume that $\lambda_1(\Omega, s, 2) > 0$ and we prove the following fact: there exists a constant $C_2(N, s)$, that only depends on N and s , such that

$$\|w\|_{L^\infty(\Omega)}^{2-q} \lambda_1(\Omega, s, 2) \leq C_1(N, s, q) \tag{7.1}$$

holds with a suitable constant $C_1(N, s, q)$, depending only on N, s and q , provided that

$$\|w\|_{L^\infty(\Omega)}^{2-q} \geq C_2(N, s). \tag{7.2}$$

That fact would imply $\|w_{\Omega,s,q}\|_{L^\infty(\Omega)}^{2-q} \lambda_1(\Omega, s, 2) \leq \max\{C_2 \lambda_1(\Omega, s, 2), C_1\}$, whence we would infer (1.9) by a scaling argument, because for all $t > 0$ we have

$$\|w_{t\Omega,s,q}\|_{L^\infty(t\Omega)}^{2-q} = t^{2s} \|w_{\Omega,s,q}\|_{L^\infty(\Omega)}^{2-q}, \quad \lambda_1(t\Omega, s, 2) = t^{-2s} \lambda_1(\Omega, s, 2).$$

In order to prove that, for appropriate choices of constants, (7.2) implies (7.1) as desired, we follow the lines of the proof of [30, Theorem 9]. Since the L^∞ -norm is lower semicontinuous with respect to the pointwise (monotone) convergence and the first eigenvalue $\lambda_1(\cdot, s, 2)$ is monotone non-increasing with respect to set inclusion, in order to prove the claim we may assume Ω to be smooth and bounded, up to an approximation argument. So, by arguing under this assumption, in view of Proposition 3.1 we will assume w to belong to $L^\infty(\Omega)$ and to achieve its maximum at an interior point, that we may consider to be the origin in \mathbb{R}^N up to an unessential translation.

We now identify w with the function that agrees with w in Ω and equals zero everywhere else and we claim that w is a weak subsolution of the fractional Lane–Emden equation (1.4) in \mathbb{R}^N . To see this, we fix a non-negative function $\eta \in C_0^\infty(\mathbb{R}^N)$ and, for every $\varepsilon > 0$, we take a monotone non-decreasing Lipschitz continuous function $H_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$, with $H_\varepsilon(u) = 0$ for all $u \leq 0$ and $H_\varepsilon(u) = 1$ for all $u \geq \varepsilon$. Then

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{w(x) - w(y)}{|x - y|^{N+2s}} [H_\varepsilon(w(x))\eta(x) - H_\varepsilon(w(y))\eta(y)] dx dy = \int_{\Omega} w^{q-1} H_\varepsilon(w)\eta dx \tag{7.3}$$

because of the weak equation for w with $H_\varepsilon(w)\eta$ as a test function. To handle the left hand side of (7.3), we write the identity $2(a\xi - b\zeta) = (a + b)(\xi - \zeta) + (a - b)(\xi + \zeta)$ with $a = \eta(x)$, $b = \eta(y)$, $\xi = H_\varepsilon(w(x))$ and $\zeta = H_\varepsilon(w(y))$. After multiplying the result by $w(x) - w(y)$ and integrating against the singular kernel on $\Omega \times \Omega$, we see that

$$\begin{aligned} & 2 \int_{\Omega} \int_{\Omega} \frac{w(x) - w(y)}{|x - y|^{N+2s}} [H_\varepsilon(w(x))\eta(x) - H_\varepsilon(w(y))\eta(y)] dx dy \\ &= \int_{\Omega} \int_{\Omega} \frac{w(x) - w(y)}{|x - y|^{N+2s}} (\eta(x) + \eta(y)) [H_\varepsilon(w(x)) - H_\varepsilon(w(y))] dx dy \tag{7.4} \\ &+ \int_{\Omega} \int_{\Omega} \frac{w(x) - w(y)}{|x - y|^{N+2s}} (\eta(x) - \eta(y)) [H_\varepsilon(w(x)) + H_\varepsilon(w(y))] dx dy. \end{aligned}$$

Notice that the first integral in the right hand side of (7.4) is non-negative, due to the monotonicity of the function H_ε . Thus

$$\begin{aligned} & \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{w(x) - w(y)}{|x - y|^{N+2s}} [H_\varepsilon(w(x))\eta(x) - H_\varepsilon(w(y))\eta(y)] dx dy \\ & \geq \frac{1}{2} \int_{\Omega} \int_{\Omega} \frac{w(x) - w(y)}{|x - y|^{N+2s}} (\eta(x) - \eta(y)) [H_\varepsilon(w(x)) + H_\varepsilon(w(y))] dx dy \tag{7.5} \\ & + 2 \int_{\Omega} \int_{\mathbb{R}^N \setminus \Omega} \frac{w(x)}{|x - y|^{N+2s}} [H_\varepsilon(w(x))\eta(x) - H_\varepsilon(w(y))\eta(y)] dx dy. \end{aligned}$$

By dominated convergence theorem, the limit as $\varepsilon \rightarrow 0^+$ in (7.3) and (7.5) gives

$$\begin{aligned} & \int_{\Omega} \int_{\Omega} \frac{w(x) - w(y)}{|x - y|^{N+2s}} (\eta(x) - \eta(y)) dx dy + 2 \int_{\Omega} \int_{\mathbb{R}^N \setminus \Omega} \frac{w(x)\eta(x)}{|x - y|^{N+2s}} dx dy \\ & \leq \int_{\Omega} w^{q-1} \eta dx, \end{aligned}$$

and that proves the claim.

We let r be a positive radius, that will be chosen later, and we take a cut-off $\zeta \in C_0^\infty(\Omega)$ from the ball $B_{r/2}$ to B_r , with $|\nabla\zeta| \leq \frac{2}{r}$. Since w is a weak subsolution of (1.4), the localised Caccioppoli estimate of [11, Proposition 3.5], with $F = w^{q-1}$, $p = 2$, $\beta = 1$, $\delta = 0$, $L = 1$ and $\Omega' = B_r$, gives

$$\int_{B_r} \int_{B_r} \frac{(w(x)\zeta(x) - w(y)\zeta(y))^2}{|x - y|^{N+2s}} dx dy \leq C_3(N, s) \left(w(0)^q r^N + w(0)^2 r^{N-2s} \right), \tag{7.6}$$

where $C_3(N, s) > 0$ depends only on N and s . Moreover, by the fact that $w \in L^\infty(\Omega)$,

$$\int_{B_r} \int_{\mathbb{R}^N \setminus B_r} \frac{(w(x)\zeta(x) - w(y)\zeta(y))^2}{|x - y|^{N+2s}} dx dy \leq C_4(N, s) w(0)^2 r^{N-2s}, \tag{7.7}$$

where $C_4(N, s) > 0$ depends only on N and s . Also, by Proposition 4.2 we have

$$\int_{B_r} w^2 dx \geq C_5(N, s, q) r^N \left(w(0) - \delta \text{Tail}(w, 0, r/2) - \delta r^{2s} w(0) \right)^2,$$

where the constant $C_5(N, s, q)$ depends only on N, s and q and $\delta \in (0, 1]$ is a parameter that we can take as small as we wish. By combining the latter with (7.6) and (7.7), for δ smaller than an appropriate $\delta_0(N, s) \in (0, 1]$, we obtain

$$\lambda_1(\Omega, s, 2) \leq C_6(N, s, q) \left(w(0)/2 - \delta r^{2s} w(0)^{q-2} \right)^{-2} \left(w(0)^q + w(0)^2 r^{-2s} \right), \tag{7.8}$$

where we set $C_6 = 2(C_3 + C_4)C_5^{-1}$ and we used the fact that the function $w\zeta$ is an admissible competitor for the infimum that defines the constant $\lambda_1(\Omega, s, 2)$. Then, we take $\delta \leq 2^{-q} \wedge \delta_0(N, s)$. Hence, with the choice

$$r = \left(\frac{1}{2} w(0) \right)^{\frac{2-q}{2s}},$$

we have $w(0) - \delta r^{2s} w(0)^{q-2} \geq w(0)/4$, and (7.8) gives (7.1) with $C_1 = 16(1 + 2^{2-q})C_6$. □

Remark 7.1 If $\lambda_1(\Omega, s, q) > 0$, we have the stronger estimate

$$\|w_{\Omega, s, q}\|_{L^\infty(\Omega)} \leq \mathcal{C} \lambda_1(\Omega, s, q)^{-\gamma}$$

where the constant $\mathcal{C} > 0$ and the exponent $\gamma > 0$ depends only on N, s and q . Indeed, $w_{\Omega, s, q}$ is the first q -semilinear s -eigenfunction with $L^q(\Omega)$ -norm $\lambda_1(\Omega, s, q)^{\frac{1}{q-2}}$, and the estimate follows at once by Proposition 3.1. In fact, this estimate can also be seen as a particular case of the more general one of Theorem B. Indeed, the positivity of the greatest lower bound $\lambda_1(\Omega, s, 2)$ for the spectrum of the fractional (linear) s -Laplacian is, by definition, equivalent to the continuity of the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^2(\Omega)$. Domains with this property are not necessarily bounded, nor are they required to

have finite measure; also, an open set Ω may support a Sobolev–Poincaré inequality that makes $\lambda_1(\Omega, s, 2)$ strictly positive even if $\lambda_1(\Omega, s, q) = 0$ for all $q \in (1, 2)$ (examples are provided by bounded cylinders $\omega \times (-M, M)$, with $M > 0$ and $\omega \subset \mathbb{R}^{N-1}$.) Conversely, given any $q \in (1, 2)$, the fact that $\lambda_1(\Omega, s, q) > 0$ implies that $\lambda_1(\Omega, s, 2) > 0$, too; in fact, it implies that the embedding $\mathcal{D}_0^{s,2}(\Omega) \hookrightarrow L^2(\Omega)$ is compact, by interpolation: see [19, Lemma 2.3].

7.3 Proof of Theorem C

Arguing by contradiction, we assume that a sequence $(\lambda_n)_{n \in \mathbb{N}} \subset \mathfrak{S}(\Omega, s, q)$ converges to $\lambda_1(\Omega, s, q)$. For each λ_n , we pick an eigenfunction u_n with unit norm in $L^q(\Omega)$. That defines a bounded sequence in $\mathcal{D}_0^{s,2}(\Omega)$, due to equation (2.5) with $\lambda = \lambda_n$ and $u = \varphi = u_n$. Then, by possibly passing to a subsequence, we may assume that u_n converges weakly in $\mathcal{D}_0^{s,2}(\Omega)$ and strongly in $L^q(\Omega)$ to a limit function u with unit norm in $L^q(\Omega)$. Hence, by passing to the limit as $n \rightarrow \infty$ in (2.5) with $u = u_n$ and $\lambda = \lambda_n$, it is easily seen that u is a first q -semilinear s -eigenfunction. Owing to Theorem A, up to changing everywhere sign to each element of the sequence, $u > 0$ and its multiple $w = \lambda_1(\Omega, s, q)^{\frac{1}{q-2}} u$ is the fractional Lane–Emden density of Ω . Moreover, each function $v_n = \lambda_1(\Omega, s, q)^{\frac{1}{q-2}} u_n$ is a weak solution of the fractional Lane–Emden equation (1.4); yet, by construction, v_n converges to $w_{\Omega, s, q}$ in $\mathcal{D}_0^{s,2}(\Omega)$, in contradiction with Proposition 6.2. \square

Acknowledgements The authors are grateful to Lorenzo Brasco for fruitful discussions, improving the quality of the paper, for pointing attention to the Hardy-type inequality, and for suggesting the approximation trick used in Theorem B; an anonymous referee is acknowledged for pointing out the relevance of the sub-modularity of squared norms. This material is based upon work supported by the Swedish Research Council under Grant No. 2016-06596 while G. Franzina was in residence at Institut Mittag-Leffler in Djursholm, Sweden during the Research Program “Geometric Aspects of Nonlinear Partial Differential Equations” in 2022. The research of D. Licheri is supported by the University of Cagliari, Italy and the Fondazione di Sardegna under the Grant No. CUP:F72F20000200007.

Declarations

Conflict of interest The authors have no relevant financial or non-financial interests to disclose, and declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Appendix A

The following one is an obvious consequence of inequality $(|a| - |b|)^2 \leq (a - b)^2$, that is strict if and only if $ab < 0$.

Lemma 7.1 *For all $u \in \mathcal{D}_0^{s,2}(\Omega)$, we have*

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(|u(x)| - |u(y)|)^2}{|x - y|^{N+2s}} dx dy \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy \tag{A.1}$$

with strict inequality unless either $u \geq 0$ or $u \leq 0$ a.e. in Ω .

Incidentally, we observe that for all functions $u \in \mathcal{D}_0^{s,2}(\Omega)$ the following identity also holds:

$$\begin{aligned} & \iint_{\mathbb{R}^{2N}} \frac{(|u(x)| - |u(y)|)^2}{|x - y|^{N+2s}} dx dy \\ &= \iint_{\mathbb{R}^{2N}} \frac{((u \wedge 0)(x) - (u \wedge 0)(y))^2}{|x - y|^{N+2s}} dx dy \\ &+ \iint_{\mathbb{R}^{2N}} \frac{((u \vee 0)(x) - (u \vee 0)(y))^2}{|x - y|^{N+2s}} dx dy \\ &+ 4 \int_{\{u(x)>0\}} \int_{\{u(y)<0\}} \frac{u(x)u(y)}{|x - y|^{N+2s}} dx dy. \end{aligned}$$

Sine the last integral is negative, the previous Lemma can also be deduced from the submodularity of the squared seminorm, see [23, Lemma 3] with $v = 0$.

The following form of the minimum principle for supersolutions is well known: e.g., see [27, Lemma 6]. For convenience of the reader, we present the proof of its statement in the setting relevant to the present manuscript, and we point out that Ω is not required to be connected.

Proposition 7.1 *Let $u \in \mathcal{D}_0^{s,2}(\Omega)$ satisfy*

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))(\varphi(x) - \varphi(y))}{|x - y|^{N+2s}} dx dy \geq 0 \text{ for all non-negative } \varphi \in C_0^\infty(\Omega)$$

and assume that $u \geq 0$ a.e. in Ω . Then, either $u = 0$ a.e. in Ω or $u > 0$ a.e. in Ω .

Proof By [6, Theorem A.1], $u > 0$ in each connected component where it is not identically zero. Then, we argue as in the proof of [11, Proposition 2.6] and we prove a contrapositive statement: if $u \equiv 0$ in a connected component Ω_0 of Ω , then, by assumption, for all $\varphi \in C_0^\infty(\Omega_0) \setminus \{0\}$ such that $\varphi \geq 0$,

$$\int_{\Omega \setminus \Omega_0} \int_{\Omega_0} \frac{u(x)\varphi(y)}{|x - y|^{n+2s}} dx dy - \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))(\varphi(x) - \varphi(y))}{|x - y|^{n+2s}} dx dy \leq 0$$

which, by Fubini's theorem, implies $u = 0$ a.e. in $\Omega \setminus \Omega_0$, hence a.e. in Ω . □

References

1. Anello, G., Faraci, F., Iannizzotto, A.: On a problem of Huang concerning best constants in Sobolev embeddings. *Ann. Mat. Pura Appl.* **194**(3), 767–779 (2015)
2. Bonforte, M., Figalli, A., Vázquez, J.L.: Sharp boundary behaviour of solutions to semilinear nonlocal elliptic equations. *Calc. Var. Partial Differ. Equ.* **57**(2), 1–43 (2018). (**Article No. 57**)
3. Brasco, L.: On principal frequencies and isoperimetric ratios in convex sets. *Ann. Fac. Sci. Toulouse Math.* (6) **29**(4), 977–1005 (2020)
4. Brasco, L., Cinti, E.: On fractional Hardy inequalities in convex sets. *Discrete Contin. Dyn. Syst.* **38**(8), 4019–4040 (2018)
5. Brasco, L., De Philippis, G., Franzina, G.: Positive solutions to the sublinear Lane–Emden equation are isolated. *Commun. Partial Differ. Equ.* **46**(10), 1940–1972 (2021)
6. Brasco, L., Franzina, G.: Convexity properties of Dirichlet integrals and Picone-type inequalities. *Kodai Math. J.* **37**(3), 769–799 (2014)
7. Brasco, L., Franzina, G.: An overview on constrained critical points of Dirichlet integrals. *Rend. Semin. Mat. Univ. Politec. Torino* **78**(2), 7–50 (2020)
8. Brasco, L., Franzina, G., Ruffini, B.: Schrödinger operators with negative potentials and Lane–Emden densities. *J. Funct. Anal.* **274**(6), 1825–1863 (2018)
9. Brasco, L., Gómez-Castro, D., Vázquez, J.L.: Characterisation of homogeneous fractional Sobolev spaces. *Calc. Var. Partial Differ. Equ.* **60**(2), 60 (2021)
10. Brasco, L., Lindgren, E., Parini, E.: The fractional Cheeger problem. *Interfaces Free Bound.* **16**(3), 419–458 (2014)
11. Brasco, L., Parini, E.: The second eigenvalue of the fractional p -Laplacian. *Adv. Calc. Var.* **9**(4), 323–355 (2016)
12. Cabré, X., Figalli, A., Ros-Oton, X., Serra, J.: Stable solutions to semilinear elliptic equations are smooth up to dimension 9. *Acta Math.* **224**(2), 187–252 (2020)
13. Cheng, Z.Q., Song, R.: Hardy inequality for censored stable processes. *Tohoku Math. J. (2)* **55**(3), 439–450 (2003)
14. Deny, J., Lions, J.L.: Les espaces du type de Beppo Levi. *Ann. Inst. Fourier (Grenoble)* **5**, 305–370 (1954)
15. Di Castro, A., Kuusi, T., Palatucci, G.: Local behavior of fractional p -minimizers. *Ann. Inst. Henri Poincaré Anal. Non Linéaire* **33**(5), 1279–1299 (2016)
16. Dyda, B.: A fractional order Hardy inequality. *Ill. J. Math.* **48**(2), 575–588 (2004)
17. Dyda, B., Vähäkangas, A.V.: A framework for fractional Hardy inequalities. *Ann. Acad. Sci. Fenn. Math.* **39**, 675–689 (2014)
18. Dyda, B., Vähäkangas, A.V.: Characterizations for fractional Hardy inequality. *Adv. Calc. Var.* **8**(2), 173–182 (2015)
19. Franzina, G.: Non-local torsion functions and embeddings. *Appl. Anal.* **98**(10), 1811–1826 (2019)
20. Grisvard, P.: *Elliptic Problems in Nonsmooth Domains*. Pitman, London (1985)
21. Hörmander, L., Lions, J.L.: Sur la complétion par rapport à une intégrale de Dirichlet. *Math. Scand.* **4**(2), 259–270 (1956)
22. Molica Bisci, G., Rădulescu, V.D., Servadei, R.: *Variational Methods for Nonlocal Fractional Problems*. Encyclopedia of Mathematics and Its Applications 162, Cambridge University Press, Cambridge (2016)
23. Palatucci, G., Savin, O., Valdinoci, E.: Local and global minimizers for a variational energy involving a fractional norm. *Ann. Mat. Pura Appl.* **192**(4), 673–718 (2013)
24. Ros-Oton, X.: Nonlocal elliptic equations in bounded domains: a survey. *Publ. Mat.* **60**(1), 3–26 (2016)
25. Ros-Oton, X., Serra, J.: The Dirichlet problem for the fractional Laplacian: regularity up to the boundary. *J. Math. Pures Appl.* **101**(3), 275–302 (2014)
26. Servadei, R., Valdinoci, E.: Variational methods for non-local operators of elliptic type. *Discrete Contin. Dyn. Syst.* **33**(5), 2105–2137 (2013)
27. Servadei, R., Valdinoci, E.: Weak and viscosity solutions of the fractional Laplace equation. *Publ. Mat.* **58**(1), 133–154 (2014)
28. Sk, F.: Characterization of fractional Sobolev–Poincaré and (localized) Hardy inequalities (2022, Preprint)
29. Struwe, M.: *Variational Methods. Applications to Nonlinear Partial Differential Equations and Hamiltonian Systems*. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*, vol. 34. Springer, Berlin (2008)

30. van der Berg, M., Bucur, D.: On the torsion function with Robin or Dirichlet boundary conditions. *J. Funct. Anal.* **266**(3), 1647–1666 (2014)
31. Vázquez, J.L.: Recent progress in the theory of nonlinear diffusion with fractional Laplacian operators. *Discrete Contin. Dyn. Syst. S* **7**(4), 857–885 (2014)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.